

LOW COMPLEXITY SOLUTIONS OF THE ALLEN-CAHN EQUATION ON THREE-SPHERES

ROBERT HASLHOFER, MOHAMMAD N. IVAKI

ABSTRACT. In this short note, we prove that on the three-sphere with any bumpy metric there exist at least four solutions of the Allen-Cahn equation with spherical interface and index at most two. The proof combines several recent results from the literature.

1. INTRODUCTION

In this short note, we consider the Allen-Cahn equation,

$$(1.1) \quad -\Delta_g u = \frac{1}{\varepsilon^2} (u - u^3),$$

on S^3 endowed with any bumpy metric g . The Allen-Cahn equation has its origin as a model for phase transitions [AC79] and has received a lot of attention especially due to its close connections with the theory of minimal surfaces, see e.g., the surveys [T08, S09, P12].

In a beautiful recent paper [GG], Gaspar-Guaraco proved that on any closed manifold the number of solutions of (1.1) goes to infinity as $\varepsilon \rightarrow 0$. Most of these solutions naturally have high complexity, i.e., large index and interfaces of high genus. In this note, we address the remaining question of finding solutions with low complexity, i.e., solutions with interface of genus zero and low index, and prove:

Theorem 1.2. *Consider the Allen-Cahn equation (1.1) on S^3 endowed with any bumpy metric g . Then for any small enough $\varepsilon > 0$ there exist at least four solutions with spherical interface and index at most two. More precisely, exactly one of the following alternatives holds:*

- (1) *There are at least two stable solutions with spherical interface, and at least four index-one solutions with spherical interface.*

R.H. has been partially supported by NSERC grant RGPIN-2016-04331 and a Connaught New Researcher Award. M.I. has been supported by a Marsden Postdoctoral Fellowship. Both authors thank the Fields Institute for providing an excellent research environment during the thematic program on Geometric Analysis.

- (2) *There are no stable solutions with spherical interface, at least two index-one solutions with spherical interface, and at least two index-two solutions with spherical interface.*

The two cases are illustrated in Figure 1 and Figure 2 for the scenarios that (S^3, g) looks like a dumbbell and an elongated ellipsoid, respectively.




FIGURE 1. Solutions on a dumbbell




FIGURE 2. Solutions on elongated ellipsoids

The solutions transition from $u_\varepsilon \approx -1$ to $u_\varepsilon \approx +1$ along the spherical interface, and the transition is modeled on the function $\tanh(\frac{t}{\sqrt{2\varepsilon}})$, where t denotes the signed distance from the interface.

We recall that solutions of (1.1) are critical points of the energy functional

$$(1.3) \quad E_\varepsilon[u] = \int_M \frac{\varepsilon}{2} |\nabla u|^2 + \frac{1}{4\varepsilon} (1 - u^2)^2 \, \text{dvol}_g$$

The index of a critical point is defined as the number of directions in which the second variation of the functional is negative.

We also recall that a Riemannian metric g on a manifold M is called bumpy if (M, g) does not contain any immersed closed minimal hypersurface with a non-trivial Jacobi field. By a result of White [W91, W17] bumpy metrics are generic in the sense of Baire.

To prove Theorem 1.2, we combine several results from the literature, including in particular the gluing construction from Pacard-Ritore [PR03], the existence result for minimal two-spheres from Haslhofer-Ketover [HK], and the index estimate from Chodosh-Mantoulidis [CM]. An interesting feature is that the proof relies on some methods from other fields, in particular, mean curvature flow with surgery, that have not been used before in the analysis of the Allen-Cahn equation.

2. THE PROOF

Proof of Theorem 1.2. We first consider the case that there exists at least one solution u_ε of (1.1) which is stable and has spherical interface (for every small enough ε). Having a spherical interface, means that for $\varepsilon \rightarrow 0$ (after passing to a subsequence), the associated varifolds

$$(2.1) \quad V_\varepsilon(\phi) = \frac{3}{4} \int \varepsilon |\nabla u_\varepsilon(x)|^2 \phi(x, T_x \{u_\varepsilon = u_\varepsilon(x)\}) \, \text{dvol}_g(x)$$

converge to an embedded minimal two-sphere Σ , possibly with multiplicity. It follows from classical theory [T95, TW12] (see also [H]) that Σ is a stable critical point of the area functional. Thus, by a result of Ketover-Liokumovich [KL, Thm. 1.6] each of the two balls $B^{\ell/r}$ in S^3 bounded by Σ contains in its interior an index-one minimal two-sphere $\Sigma^{\ell/r}$. We can now apply the main theorem of Pacard-Ritore [PR03] (see also [P12]) to get solutions $u_\varepsilon^{\ell/r}$ with spherical interface $\Sigma^{\ell/r}$. Since for the Pacard-Ritore solutions $u_\varepsilon^{\ell/r}$ the associated varifolds $V_\varepsilon^{\ell/r}$ converge to $\Sigma^{\ell/r}$ with multiplicity one, it follows from Chodosh-Mantoulidis [CM, Sec. 6] (see also [dPKW13, Sec. 9] for closely related computations) that the index is actually continuous under the limit, i.e., for ε small enough we get

$$(2.2) \quad \text{ind}(u_\varepsilon^{\ell/r}) = \text{ind}(\Sigma^{\ell/r}) = 1,$$

where the possibility of nullity is ruled out by the bumpyness assumption. Together with the obvious fact that whenever u is a solution of (1.1) then $-u$ also is a solution, this finishes the proof in case (1).

Consider now the case that (1.1) admits no stable solutions with spherical interface. Suppose towards a contradiction that (S^3, g) contains a stable embedded minimal two-sphere Σ . Then, arguing as above, for ε small enough the associated Pacard-Ritore solutions would satisfy

$$(2.3) \quad \text{ind}(u_\varepsilon) = \text{ind}(\Sigma) = 0,$$

contradicting our assumption that (1.1) has no stable solutions with spherical interface. Hence, by a classical result of Simon-Smith [SS82] (see also [MN12, Lem. 3.5] and [HK, Lem. 3.2]) there exists an embedded minimal two-sphere $\Sigma^1 \subset (S^3, g)$, which is obtained by one parameter min-max, realizes the 1-width, and has index one (since it can't be stable). By a recent result from Haslhofer-Ketover [HK, Thm. 1.4], which has been established using combined efforts from mean curvature flow with surgery [BH, HK17, BHH] and min-max theory [KMN], there also exists an embedded minimal two-sphere $\Sigma^2 \subset (S^3, g)$ which has index two. Arguing as above, for ε small enough the associated Pacard-Ritore solutions satisfy

$$(2.4) \quad \text{ind}(\pm u_\varepsilon^1) = \text{ind}(\Sigma^1) = 1,$$

and

$$(2.5) \quad \text{ind}(\pm u_\varepsilon^2) = \text{ind}(\Sigma^2) = 2.$$

This establishes case (2), and thus finishes the proof of the theorem. \square

REFERENCES

- [AC79] S. Allen, J. W. Cahn, *A microscopic theory for antiphase boundary motion and its application to antiphase domain coarsening*, Acta. Metall. 27 (1979), 1084–1095.
- [BH] S. Brendle, G. Huisken, *Mean curvature flow with surgery of mean convex surfaces in three-manifolds* J. Eur. Math. Soc. (to appear).
- [BHH] R. Buzano, R. Haslhofer, O. Hershkovits, *The moduli space of two-convex embedded spheres*, arXiv:1607.05604, 2016.
- [CM] O. Chodosh, C. Mantoulidis, *Minimal surfaces and the Allen-Cahn equation on 3-manifolds: index, multiplicity, and curvature estimates*, arXiv:1803.02716, 2018.
- [dPKW13] M. del Pino, M. Kowalczyk, J. Wei, *Entire solutions of the Allen-Cahn equation and complete embedded minimal surfaces of finite total curvature in \mathbb{R}^3* . J. Differential Geom. 93 (2013), no. 1, 67–131.
- [GG] P. Gaspar, M. Guaraco, *The Allen-Cahn equation on closed manifolds*, arXiv:1608.06575, 2016.
- [H] F. Hiesmayr, *Spectrum and index of two-sided Allen-Cahn minimal hypersurfaces*, arXiv:1704.07738, 2017.
- [HK17] R. Haslhofer, B. Kleiner, *Mean curvature flow with surgery*, Duke Math. J. 166 (2017), no. 9, 1591–1626.

- [HK] R. Haslhofer, D. Ketover, *Minimal two-spheres in three-spheres*, arXiv:1708.06567, 2017.
- [KL] D. Ketover, Y. Liokumovich, *On the existence of unstable minimal Heegaard surfaces*, arXiv:1709.09744, 2017.
- [KMN] D. Ketover, F. Marques, A. Neves, *The catenoid estimate and its geometric applications*, arXiv:1601.04514, 2016.
- [MN12] F. Marques, A. Neves, *Rigidity of min-max minimal spheres in three-manifolds*, Duke Math. J. 161 (2012), no. 14, 2725–2752.
- [PR03] F. Pacard, M. Ritore, *From constant mean curvature hypersurfaces to the gradient theory of phase transitions*, J. Differential Geom. 64 (2003), no. 3, 359–423.
- [P12] F. Pacard, *The role of minimal surfaces in the study of the Allen-Cahn equation*, Lecture notes from the Santalo Summer School Geometric Analysis, University of Granada, Spain, 2012.
- [S09] O. Savin, *Phase transitions, minimal surfaces and a conjecture of De Giorgi*, Current developments in mathematics (2009), 59–113.
- [SS82] F. Smith, *On the existence of embedded minimal 2-spheres in the 3-sphere endowed with an arbitrary Riemannian metric*, Phd thesis, Supervisor: Leon Simon, University of Melbourne, 1982.
- [T95] Y. Tonegawa, *On stable critical points for a singular perturbation problem*, Comm. Anal. Geom. 13 (2005), no. 2, 439–459.
- [T08] Y. Tonegawa, *Applications of geometric measure theory to two-phase separation problems*, Sugaku Expositions, 21 (2008), 178–196.
- [TW12] Y. Tonegawa, N. Wickramasekera, *Stable phase interfaces in the van der Waals-Cahn-Hilliard theory*, J. Reine Angew. Math. 668 (2012), 191–210.
- [W91] B. White, *The space of minimal submanifolds for varying Riemannian metrics*, Indiana Univ. Math. J. 40 (1991), no. 1, 161–200.
- [W17] B. White, *On the bumpy metrics theorem for minimal submanifolds*, Amer. J. Math. 139 (2017), no. 4, 1149–1155.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, 40
ST GEORGE STREET, TORONTO, ON M5S 2E4, CANADA

E-mail: roberth@math.toronto.edu, m.ivaki@utoronto.ca