

V-FILTRATIONS AND MINIMAL EXPONENTS FOR LOCAL COMPLETE INTERSECTIONS

QIANYU CHEN, BRADLEY DIRKS, MIRCEA MUSTĂŢĂ, AND SEBASTIÁN OLANO

ABSTRACT. We define and study a notion of *minimal exponent* for a local complete intersection subscheme Z of a smooth complex algebraic variety X , extending the invariant defined by Saito in the case of hypersurfaces. Our definition is in terms of the Kashiwara-Malgrange V -filtration associated to Z . We show that the minimal exponent describes how far the Hodge filtration and order filtration agree on the local cohomology $\mathcal{H}_Z^r(\mathcal{O}_X)$, where r is the codimension of Z in X . We also study its relation to the Bernstein-Sato polynomial of Z . Our main result describes the minimal exponent of a higher codimension subscheme in terms of the invariant associated to a suitable hypersurface; this allows proving the main properties of this invariant by reduction to the codimension 1 case. A key ingredient for our main result is a description of the Kashiwara-Malgrange V -filtration associated to *any* ideal (f_1, \dots, f_r) in terms of the microlocal V -filtration associated to the hypersurface defined by $\sum_{i=1}^r f_i y_i$.

1. INTRODUCTION

Let X be a smooth, irreducible, complex algebraic variety. If Z is a nonempty hypersurface in X , then the *minimal exponent* $\tilde{\alpha}(Z)$ of Z (written also as $\tilde{\alpha}(f)$ if Z is defined by $f \in \mathcal{O}_X(X)$) is an important invariant of the singularities of Z introduced by Saito [Sai94]. When Z has isolated singularities, it can be described via asymptotic expansions of integrals along vanishing cycles and it was studied extensively in the 80s, see for example [Var82], [Ste85], and [Loe84]; in this setting, it has been known as *complex singularity index* or *Arnold exponent* of Z . In general, it is defined as the negative of the largest root of the *reduced Bernstein-Sato polynomial* of Z (with the convention that it is ∞ if this polynomial is 1, which is the case if and only if Z is smooth). By results of Kollár [Kol97] and Lichtin [Lic89], it is known that the minimal exponent refines an important invariant of singularities in birational geometry, the *log canonical threshold* $\text{lct}(X, Z)$; more precisely, we always have $\text{lct}(X, Z) = \min\{\tilde{\alpha}(Z), 1\}$. Our main goal in this paper is to introduce and study a generalization of the minimal exponent to the case when Z is locally a complete intersection in X .

Before giving the definition in the general context, we recall the connection between the minimal exponent of hypersurfaces and two important \mathcal{D} -module theoretic constructions associated to Z , the Hodge filtration on the local cohomology $\mathcal{H}_Z^1(\mathcal{O}_X)$ of \mathcal{O}_X along Z and the Kashiwara-Malgrange V -filtration associated to Z . Recall that if Z is any closed subscheme of X , the local cohomology sheaves $\mathcal{H}_Z^q(\mathcal{O}_X)$ underlie mixed Hodge modules in the sense of Saito's theory [Sai90]. In particular, they carry a Hodge filtration: this is an increasing filtration by coherent \mathcal{O}_X -modules which is compatible with the order filtration on the sheaf

2020 *Mathematics Subject Classification.* 14F10, 14B05, 14J17.

Q.C. was partially supported by NSF grant DMS-1952399, B.D. was partly supported by NSF grant DMS-2001132 and NSF-MSPRF grant DMS-2303070, and M.M. was partially supported by NSF grants DMS-2301463 and DMS-1952399.

\mathcal{D}_X of differential operators on X . If Z is a reduced hypersurface in X , then the only nonzero local cohomology is $\mathcal{H}_Z^1(\mathcal{O}_X) = \mathcal{O}_X(*Z)/\mathcal{O}_X$ (where $\mathcal{O}_X(*Z)$ is the sheaf of rational functions with poles along Z). In this case it is known that for every $k \geq 0$ we have

$$F_k \mathcal{H}_Z^1(\mathcal{O}_X) \subseteq O_k \mathcal{H}_Z^1(\mathcal{O}_X) := \mathcal{O}_X((k+1)Z)/\mathcal{O}_X$$

and Saito showed in [Sai16] that

$$(1) \quad F_k \mathcal{H}_Z^1(\mathcal{O}_X) = O_k \mathcal{H}_Z^1(\mathcal{O}_X) \text{ for all } k \leq p \text{ if and only if } \tilde{\alpha}(Z) \geq p+1.$$

Using a refinement of this result to a setting involving twists by rational multiples of Z , as well as properties of Hodge filtrations, it was shown in [MP20] that one can extend the known properties of the Arnold exponent to arbitrary hypersurface singularities.

The proof of (1) makes use of results about V -filtrations. Let us briefly recall this notion, due to Malgrange [Mal83] and Kashiwara [Kas83], in the more general context that is relevant to this paper. Working locally, let us suppose that Z is a closed subscheme of X defined by the ideal generated by nonzero regular functions $f_1, \dots, f_d \in \mathcal{O}_X(X)$. If $\iota: X \hookrightarrow X \times \mathbf{A}^d$ is the graph embedding associated to $\mathbf{f} = (f_1, \dots, f_d)$, that is, $\iota(x) = (x, f_1(x), \dots, f_d(x))$, then the V -filtration is a decreasing filtration $(V^\gamma B_{\mathbf{f}})_{\gamma \in \mathbf{Q}}$ on

$$B_{\mathbf{f}} := \iota_* \mathcal{O}_X = \bigoplus_{\beta \in \mathbf{Z}_{\geq 0}^d} \mathcal{O}_X \partial_t^\beta \delta_{\mathbf{f}},$$

indexed by rational numbers, and characterized by a few properties (for details, see Section 2). In the case of one function, the V -filtration plays an important role in the theory of mixed Hodge modules. We also recall that in the case when we only have one function $g \in \mathcal{O}_X(X)$, Saito introduced in [Sai94] a related filtration, the *microlocal V -filtration* $(V^\gamma \tilde{B}_g)_{\gamma \in \mathbf{Q}}$ on

$$\tilde{B}_g = \bigoplus_{j \in \mathbf{Z}} \mathcal{O}_X \partial_t^j \delta_g.$$

If Z is the hypersurface defined by g , then the minimal exponent $\tilde{\alpha}(Z)$ is described as follows:

$$\tilde{\alpha}(Z) = \sup\{\gamma > 0 \mid \delta_g \in V^\gamma \tilde{B}_g\}$$

(see [Sai16, (1.3.4)]). In terms of the usual V -filtration, this says that if q is a nonnegative integer and $\gamma \in (0, 1]$ is a rational number, then

$$(2) \quad \tilde{\alpha}(Z) \geq q + \gamma \text{ if and only if } \partial_t^q \delta_g \in V^\gamma B_g.$$

Suppose now that Z is a closed subscheme of X that is a local complete intersection, of pure codimension $r \geq 1$. We define the minimal exponent $\tilde{\alpha}(Z)$ such that the analogue of formula of (2) holds in this setting. Working locally, we may assume that Z is defined by the ideal generated by $f_1, \dots, f_r \in \mathcal{O}_X(X)$. In this case, we put

$$(3) \quad \tilde{\alpha}(Z) = \begin{cases} \sup\{\gamma > 0 \mid \delta_{\mathbf{f}} \in V^\gamma B_{\mathbf{f}}\}, & \text{if } \delta_{\mathbf{f}} \notin V^r B_{\mathbf{f}}; \\ \sup\{r-1+q+\gamma \mid \partial_t^\beta \delta_{\mathbf{f}} \in V^{r-1+\gamma} B_{\mathbf{f}} \text{ for } |\beta| \leq q\}, & \text{if } \delta_{\mathbf{f}} \in V^r B_{\mathbf{f}}, \end{cases}$$

where in the latter case, the supremum is over all nonnegative integers q and all rational numbers $\gamma \in (0, 1]$ with the property that $\partial_t^\beta \delta_{\mathbf{f}} \in V^{r-1+\gamma} B_{\mathbf{f}}$ for all $\beta = (\beta_1, \dots, \beta_r) \in \mathbf{Z}_{\geq 0}^r$, with $\beta_1 + \dots + \beta_r \leq q$. In fact, the supremum in the definition is a maximum unless $\tilde{\alpha}(Z) = \infty$ (which we show is the case if and only if Z is smooth). We note that by [BMS06, Theorem 1], which describes the multiplier ideals of Z in terms of $V^\bullet B_{\mathbf{f}}$, we have

$$\text{lct}(X, Z) = \min\{\tilde{\alpha}(Z), r\}.$$

We note that $\tilde{\alpha}(Z)$ does depend on the ambient variety and not just on Z . Whenever X is not understood from the context, we write $\tilde{\alpha}(X, Z)$ in order to avoid confusion. However, the dependence is easy to understand: the difference $\tilde{\alpha}(X, Z) - \dim(X)$ only depends on Z (see Proposition 4.14).

In order to prove the basic properties of the minimal exponent for local complete intersections, we describe it as the minimal exponent of a hypersurface. Arguing locally, we may again assume that Z has pure codimension r in X and it is defined in X by the ideal generated by $f_1, \dots, f_r \in \mathcal{O}_X(X)$. We consider $Y = X \times \mathbf{A}^r$, with coordinates y_1, \dots, y_r on \mathbf{A}^r , and let $U = X \times (\mathbf{A}^r \setminus \{0\})$.

Theorem 1.1. *With the above notation, if $g = \sum_{i=1}^r f_i y_i \in \mathcal{O}_Y(Y)$, then*

$$\tilde{\alpha}(Z) = \tilde{\alpha}(g|_U).$$

The proof of Theorem 1.1 relies on a general result of independent interest describing the V -filtration associated to $f_1, \dots, f_d \in \mathcal{O}_X(X)$ (without any complete intersection assumption) in terms of the microlocal V -filtration associated to $g = \sum_{i=1}^d f_i y_i \in \mathcal{O}_X(X)[y_1, \dots, y_d]$; see Theorem 3.3 for the precise statement. Another application of this connection is a relation between b -functions corresponding to f_1, \dots, f_d and microlocal b -functions corresponding to g . This greatly extends the main result of [Mus22], which says that the Bernstein-Sato polynomial $b_Z(s)$ of Z is equal to the reduced Bernstein-Sato polynomial $b_g(s)/(s+1)$ of g .

The description in Theorem 1.1, together with the results on minimal exponents of hypersurfaces from [MP20], allow us to obtain similar results for local complete intersections. In order to state these, it is convenient to use a local version of the minimal exponent. If Z is a local complete intersection in X as above and $x \in Z$ is a point, then we put $\tilde{\alpha}_x(Z) := \max_{V \ni x} \tilde{\alpha}(V, Z \cap V)$, where the maximum is over the open neighborhoods V of x in X .

Theorem 1.2. *Let X be a smooth, irreducible, n -dimensional complex algebraic variety and let Z be a local complete intersection closed subscheme of X , of pure codimension r .*

- i) *If H is a smooth hypersurface in X that contains no irreducible component of Z and $Z_H = Z \cap H \hookrightarrow H$, then for every $x \in Z_H$, we have*

$$\tilde{\alpha}_x(H, Z_H) \leq \tilde{\alpha}_x(X, Z).$$

- ii) *Given a smooth morphism $\mu: X \rightarrow T$ such that for every $t \in T$, $Z_t := Z \cap \mu^{-1}(t) \hookrightarrow X_t = \mu^{-1}(t)$ has pure codimension r , then the following hold:*

- ii₁) *For every $\alpha \in \mathbf{Q}_{>0}$, the set*

$$\{x \in Z \mid \tilde{\alpha}_x(X_{\mu(x)}, Z_{\mu(x)}) \geq \alpha\}$$

is open in Z .

- ii₂) *There is an open subset T_0 of T such that for every $t \in T_0$ and $x \in Z_t$, we have*

$$\tilde{\alpha}_x(X_t, Z_t) = \tilde{\alpha}_x(X, Z).$$

In particular, the set $\{\tilde{\alpha}_x(X_{\mu(x)}, Z_{\mu(x)}) \mid x \in Z\}$ is finite. Moreover, if $s: T \rightarrow X$ is a section of μ such that $s(T) \subseteq Z$, then the set $\{t \in T \mid \tilde{\alpha}_{s(t)}(X_t, Z_t) \geq \alpha\}$ is open in T for every $\alpha \in \mathbf{Q}_{>0}$.

- iii) *If $x \in Z$ is a point defined by the ideal \mathfrak{m}_x and the ideal defining Z at x is contained in \mathfrak{m}_x^k , for some $k \geq 2$, then*

$$\tilde{\alpha}_x(Z) \leq \frac{n}{k}.$$

Another main result of the paper says that the minimal exponent controls the behavior of the Hodge filtration on local cohomology. Recall that if Z is locally a complete intersection of pure codimension r , the only nontrivial local cohomology of the structure sheaf is $\mathcal{H}_Z^r(\mathcal{O}_X)$, and if Z is defined by f_1, \dots, f_r , then

$$\mathcal{H}_Z^r(\mathcal{O}_X) = \mathcal{O}_X[1/f_1 \cdots f_r] / \sum_{i=1}^r \mathcal{O}_X[1/f_1 \cdots \widehat{f_i} \cdots f_r].$$

The Hodge filtration on this mixed Hodge module was studied in [MP22]. There is another natural filtration, the *order filtration* (or *Ext filtration*) given by

$$O_k \mathcal{H}_Z^r(\mathcal{O}_X) = \{u \in \mathcal{H}_Z^r(\mathcal{O}_X) \mid I_Z^{k+1} u = 0\},$$

where I_Z is the ideal defining Z . For every $k \geq 0$ we have $F_k \mathcal{H}_Z^r(\mathcal{O}_X) \subseteq O_k \mathcal{H}_Z^r(\mathcal{O}_X)$ and if equality holds for $k = p$, then it holds for all k , with $0 \leq k \leq p$. The *singularity level* $p(Z)$ of the Hodge filtration on $\mathcal{H}_Z^r(\mathcal{O}_X)$ is

$$p(Z) = \sup\{k \geq 0 \mid F_k \mathcal{H}_Z^r(\mathcal{O}_X) = O_k \mathcal{H}_Z^r(\mathcal{O}_X)\},$$

with the convention that this is -1 if the above set is empty. With this notation, we prove

Theorem 1.3. *If X is a smooth, irreducible, complex algebraic variety and Z is a local complete intersection closed subscheme of X , of pure codimension r , then*

$$p(Z) = \max\{\lfloor \tilde{\alpha}(Z) \rfloor - r, -1\}.$$

In particular, by combining Theorems 1.3 and 1.2, we see that the invariant $p(Z)$ satisfies analogous properties to those in Theorem 1.2. This was already shown in [MP22, Section 9] by different methods. For an application of Theorem 1.3 to an Inversion-of-Adjunction type statement, see Corollary 5.2. The main ingredient in the proof of Theorem 1.3 is the description of the Hodge filtration on $\mathcal{H}_Z^r(\mathcal{O}_X)$ in terms of the V -filtration on $B_{\mathbf{f}}$. This relies on the interplay between the Hodge filtration and the V -filtration for filtered \mathcal{D} -modules that underlie mixed Hodge modules. In the case of one function, this is built into the definition of Hodge modules. However, the case of several functions is more subtle and has only recently been elucidated in [CD23]. Using these results, we give the following description of the Hodge filtration on local cohomology:

Theorem 1.4. *If X is a smooth, irreducible, complex algebraic variety and $f_1, \dots, f_r \in \mathcal{O}_X(X)$ define a complete intersection closed subscheme Z of pure codimension r , then for every $p \geq 0$, we have*

$$F_p \mathcal{H}_Z^r(\mathcal{O}_X) = \left\{ \left[\sum_{|\alpha| \leq p} \frac{\alpha_1! \cdots \alpha_r! h_\alpha}{f_1^{\alpha_1+1} \cdots f_r^{\alpha_r+1}} \mid \sum_{|\alpha| \leq p} h_\alpha \partial_t^\alpha \delta_{\mathbf{f}} \in V^r B_{\mathbf{f}} \right] \right\}.$$

One interesting question that remains open is the precise relation between $\tilde{\alpha}(Z)$ and the Bernstein-Sato polynomial of Z . We recall that for an arbitrary closed subscheme of the smooth variety X , one can define a Bernstein-Sato polynomial $b_Z(s) \in \mathbf{Q}[s]$, extending the classical notion from the case of hypersurfaces (see [BMS06]). As in the classical case, all its roots are negative rational numbers, with the largest root being $-\text{lct}(X, Z)$. It is easy to see that if Z is a (nonempty) local complete intersection of pure codimension r , then $(s+r)$ divides $b_Z(s)$, see Proposition 6.1 below. By analogy with the definition of the minimal exponent in the case of hypersurfaces, we define $\tilde{\gamma}(Z)$ to be the negative of the largest root of $b_Z(s)/(s+r)$ (with the convention that this is infinite if $b_Z(s)/(s+r) = 1$).

Question 1.5. If Z is locally a complete intersection in the smooth irreducible variety X , of pure codimension r , do we have $\tilde{\alpha}(Z) = \tilde{\gamma}(Z)$?

Note that in light of Theorem 1.3, a positive answer to Question 1.5 would provide a positive answer to [MP22, Conjecture 9.11], relating the Hodge filtration on $\mathcal{H}_Z^r(\mathcal{O}_X)$ and the invariant $\tilde{\gamma}(Z)$. We can prove the following relation between the two invariants:

Theorem 1.6. *With the notation in Question 1.5, we have $\tilde{\alpha}(Z) \geq \tilde{\gamma}(Z)$ and*

$$\min \{ \tilde{\alpha}(Z), r+1 \} = \min \{ \tilde{\gamma}(Z), r+1 \}.$$

We recall that by [BMS06, Theorem 4], under the assumptions of Theorem 1.6, the subscheme Z has rational singularities if and only if $\tilde{\gamma}(Z) > r$. By combining Theorems 1.3 and 1.6, we obtain the following result, which gives a positive answer to [MP22, Conjecture 8.4].

Corollary 1.7. *If X is a smooth, irreducible variety and Z is a local complete intersection closed subscheme of X , of pure codimension r , then Z has rational singularities if and only if $\tilde{\alpha}(Z) > r$. In particular, if $F_1 \mathcal{H}_Z^r(\mathcal{O}_X) = O_1 \mathcal{H}_Z^r(\mathcal{O}_X)$, then Z has rational singularities.*

We note that since the first version of this paper was written, a positive answer to Question 1.5 has been given in [Dir23].

Outline of the paper. In Section 2 we review the basic facts about V -filtrations and b -functions. The following section is devoted to the result relating the V -filtration associated to f_1, \dots, f_d and the microlocal V -filtration associated to $\sum_{i=1}^d f_i y_i$. In Section 4 we introduce the minimal exponent of a local complete intersection subscheme, prove the description in Theorem 1.1, as well as various general properties of this invariant, including the ones in Theorem 1.2. In Section 5 we relate the minimal exponent to the Hodge filtration on local cohomology, proving Theorems 1.3 and 1.4. Finally, in the last section, we discuss the connection with the Bernstein-Sato polynomial and prove Theorem 1.6.

Acknowledgments. We would like to thank Mihnea Popa and Christian Schnell for many discussions related to the subject of this work. We are also grateful to Karl Schwede for providing some useful references and to the anonymous referee for the comments on a previous version of this article.

2. REVIEW OF V -FILTRATIONS

In this section we recall the definition and some basic properties of V -filtrations. For details, we refer to [Kas83], [BMS06, Section 1], and [Sai88, Section 3.1]. Let X be a fixed smooth, irreducible, complex algebraic variety. Recall that \mathcal{D}_X denotes the sheaf of differential operators on X . In this paper all \mathcal{D} -modules will be *left* \mathcal{D} -modules. For general facts about \mathcal{D} -modules, we refer to [HTT08].

Given nonzero regular functions $f_1, \dots, f_d \in \mathcal{O}_X(X)$, we denote by $\mathfrak{a} \subseteq \mathcal{O}_X$ the ideal (f_1, \dots, f_d) and by Z the closed subscheme of X defined by \mathfrak{a} . We consider the graph embedding

$$\iota: X \hookrightarrow W = X \times \mathbf{A}^d, \quad \iota(x) = (x, f_1(x), \dots, f_d(x))$$

and the \mathcal{D} -module theoretic push-forward $B_{\mathbf{f}} = \iota_+(\mathcal{O}_X)$ (we denote by \mathbf{f} the d -tuple (f_1, \dots, f_d)). We denote the standard coordinates on \mathbf{A}^d by t_1, \dots, t_d and use multi-index notation, so for $\beta = (\beta_1, \dots, \beta_d) \in \mathbf{Z}_{\geq 0}^d$, we put $t^\beta = t_1^{\beta_1} \cdots t_d^{\beta_d}$ and $\partial_t^\beta = \partial_{t_1}^{\beta_1} \cdots \partial_{t_d}^{\beta_d}$. We also put $\beta! = \prod_i \beta_i!$ and $|\beta| = \sum_i \beta_i$. Finally, we consider $s_i = -\partial_{t_i} t_i$ for $1 \leq i \leq d$ and $s = \sum_{i=1}^d s_i$.

It is convenient to consider $B_{\mathbf{f}}$ as an \mathcal{R} -module on X , where $\mathcal{R} = \mathcal{D}_X \langle t_1, \dots, t_d, \partial_{t_1}, \dots, \partial_{t_d} \rangle$. The general description of \mathcal{D} -module push-forward via closed immersions gives

$$B_{\mathbf{f}} = \bigoplus_{\beta \in \mathbf{Z}_{\geq 0}^d} \mathcal{O}_X \partial_t^\beta \delta_{\mathbf{f}},$$

where the actions of \mathcal{O}_X and ∂_{t_i} are the obvious ones, while the actions of $D \in \text{Der}_{\mathbf{C}}(\mathcal{O}_X)$ and of the t_i are given by

$$(4) \quad D \cdot h \partial_t^\beta \delta_{\mathbf{f}} = D(h) \partial_t^\beta \delta_{\mathbf{f}} - \sum_{i=1}^d D(f_i) h \partial_t^{\beta+e_i} \delta_{\mathbf{f}} \quad \text{and} \quad t_i \cdot h \partial_t^\beta \delta_{\mathbf{f}} = f_i h \partial_t^\beta \delta_{\mathbf{f}} - \beta_i h \partial_t^{\beta-e_i} \delta_{\mathbf{f}},$$

where e_1, \dots, e_d is the standard basis of \mathbf{Z}^d . We will also consider on $B_{\mathbf{f}}$ the *Hodge filtration*¹ given by

$$F_p B_{\mathbf{f}} = \bigoplus_{|\beta| \leq p} \mathcal{O}_X \partial_t^\beta \delta_{\mathbf{f}}.$$

It is sometimes convenient to consider the larger \mathcal{R} -module $B_{\mathbf{f}}^+$ corresponding to the push-forward $\iota_+(\mathcal{O}_X[1/f_1 \cdots f_d])$, namely

$$B_{\mathbf{f}}^+ = \bigoplus_{\beta \in \mathbf{Z}_{\geq 0}^d} \mathcal{O}_X[1/f_1 \cdots f_d] \partial_t^\beta \delta_{\mathbf{f}}.$$

We may also write the elements of $B_{\mathbf{f}}^+$ in terms of the operators s_1, \dots, s_d , as follows. Let us put $Q_m(x) = (-1)^m m! \binom{x}{m} = x(x-1) \cdots (x-m+1) \in \mathbf{C}[x]$ for $m \in \mathbf{Z}_{\geq 0}$ and

$$Q_\beta(s_1, \dots, s_d) = \prod_{i=1}^d Q_{\beta_i}(s_i) \in \mathbf{C}[s_1, \dots, s_d] \quad \text{for} \quad \beta = (\beta_1, \dots, \beta_d) \in \mathbf{Z}_{\geq 0}^d.$$

It follows from Lemma 7.1 that we can write $\partial_{t_i}^{\beta_i} = Q_{\beta_i}(s_i) t_i^{-\beta_i}$ and since $t_i^{-\beta_i} \delta_{\mathbf{f}} = \frac{1}{f_i^{\beta_i}} \delta_{\mathbf{f}}$ in $B_{\mathbf{f}}^+$, we have

$$(5) \quad \sum_{\beta} h_{\beta} \partial_t^\beta \delta_{\mathbf{f}} = \sum_{\beta} h_{\beta} \frac{Q_{\beta}(s_1, \dots, s_d)}{f_1^{\beta_1} \cdots f_d^{\beta_d}} \delta_{\mathbf{f}},$$

where $h_{\beta} \in \mathcal{O}_X[1/f_1 \cdots f_d]$ for all $\beta \in \mathbf{Z}_{\geq 0}^d$ (with only finitely many nonzero).

Since the polynomials $Q_{\beta}(s_1, \dots, s_d)$, with $\beta = (\beta_1, \dots, \beta_d)$ running over $\mathbf{Z}_{\geq 0}^d$, give a basis of $\mathbf{C}[s_1, \dots, s_d]$ over \mathbf{C} , it is easy to see that if we put $\mathbf{f}^{\mathbf{s}} = f_1^{s_1} \cdots f_d^{s_d}$, then we have an isomorphism of \mathcal{R} -modules

$$(6) \quad B_{\mathbf{f}}^+ \simeq \mathcal{O}_X[1/f_1 \cdots f_d, s_1, \dots, s_d] \mathbf{f}^{\mathbf{s}},$$

that maps $\partial_t^\beta \delta_{\mathbf{f}}$ to $\frac{Q_{\beta}(s_1, \dots, s_d)}{f_1^{\beta_1} \cdots f_d^{\beta_d}} \mathbf{f}^{\mathbf{s}}$. Note that a derivation $D \in \text{Der}_{\mathbf{C}}(\mathcal{O}_X)$ acts on $\mathbf{f}^{\mathbf{s}}$ in the expected way:

$$D \cdot \mathbf{f}^{\mathbf{s}} = \sum_{i=1}^d \frac{s_i D(f_i)}{f_i} \mathbf{f}^{\mathbf{s}}.$$

We also note that the action of t_i on the left-hand side of (6) corresponds on the right-hand side to the automorphism that maps s_i to $s_i + 1$, and similarly, the action of $s_i =$

¹It is often the case that one shifts this filtration so that what we denote by $F_p B_{\mathbf{f}}$ is considered to be $F_{p+d} B_{\mathbf{f}}$.

$-\partial_{t_i} t_i$ on the left-hand side of (6) corresponds on the right-hand side to multiplication by s_i . We sometimes tacitly use this isomorphism to denote an element of $B_{\mathbf{f}}^+$ by $P(s_1, \dots, s_d) \mathbf{f}^{\mathbf{s}}$, for some $P \in \mathcal{O}_X[1/f_1 \cdots f_d, s_1, \dots, s_d]$. Note that it follows from (5) that if we write $P(s_1, \dots, s_d) = \sum_{\beta} g_{\beta} Q_{\beta}(s_1, \dots, s_d)$, with $g_{\beta} \in \mathcal{O}_X[1/f_1 \cdots f_d]$, then $P(s_1, \dots, s_d) \mathbf{f}^{\mathbf{s}} \in B_{\mathbf{f}}$ if and only if $g_{\beta} \in \mathcal{O}_X \cdot \frac{1}{f_1^{\beta_1} \cdots f_d^{\beta_d}}$ for all $\beta \in \mathbf{Z}_{\geq 0}^d$.

We now turn to the V -filtration. On \mathcal{R} we consider the decreasing filtration

$$V^m \mathcal{R} = \bigoplus_{|\alpha| - |\beta| \geq m} \mathcal{D}_X t^{\alpha} \partial_t^{\beta}$$

for $m \in \mathbf{Z}$. It is then clear that

$$V^0 \mathcal{R} = \mathcal{D}_X \langle t_i, t_i \partial_{t_j} \mid i, j \rangle \quad \text{and} \quad V^1 \mathcal{R} = \sum_{i=1}^d t_i \cdot V^0 \mathcal{R} = \sum_{i=1}^d V^0 \mathcal{R} \cdot t_i.$$

The V -filtration on $B_{\mathbf{f}}$ is a decreasing, exhaustive filtration $(V^{\gamma} B_{\mathbf{f}})_{\gamma}$ indexed by rational numbers, which is *discrete and left-continuous*² and satisfies the following properties:

- i) Every $V^{\gamma} B_{\mathbf{f}}$ is a coherent $V^0 \mathcal{R}$ -submodule of $B_{\mathbf{f}}$.
- ii) $t_i \cdot V^{\gamma} B_{\mathbf{f}} \subseteq V^{\gamma+1} B_{\mathbf{f}}$ and $\partial_{t_i} \cdot V^{\gamma} B_{\mathbf{f}} \subseteq V^{\gamma-1} B_{\mathbf{f}}$ for all $i \leq d$ and $\gamma \in \mathbf{Q}$.
- iii) $V^1 \mathcal{R} \cdot V^{\gamma} B_{\mathbf{f}} = V^{\gamma+1} B_{\mathbf{f}}$ if $\gamma \gg 0$.
- iv) The action of $s + \gamma$ on $\text{Gr}_V^{\gamma}(B_{\mathbf{f}})$ is nilpotent for all $\gamma \in \mathbf{Q}$.

Here we put $\text{Gr}_V^{\gamma}(B_{\mathbf{f}}) = V^{\gamma} B_{\mathbf{f}} / V^{>\gamma} B_{\mathbf{f}}$, where $V^{>\gamma} B_{\mathbf{f}} = \bigcup_{\beta > \gamma} V^{\beta} B_{\mathbf{f}}$.

By the theory of Kashiwara [Kas83], extending a result of Malgrange [Mal83], there is a unique such V -filtration. Uniqueness follows by easy arguments, while existence is a deeper statement.

Remark 2.1. We note that for every $\gamma \in \mathbf{Q}$, we have $V^{\gamma} B_{\mathbf{f}}|_{X \setminus Z} = B_{\mathbf{f}}|_{X \setminus Z}$.

We recall that the V -filtration on $B_{\mathbf{f}}$ induces on $\mathcal{O}_X \simeq \mathcal{O}_X \delta_{\mathbf{f}}$ the filtration by the multiplier ideals of \mathfrak{a} (for the definition and basic properties of multiplier ideals, we refer to [Laz04, Section 9]). More precisely, it follows from [BMS06, Theorem 1] that for every $\gamma \in \mathbf{Q}_{>0}$, we have

$$\{h \in \mathcal{O}_X \mid h \delta_{\mathbf{f}} \in V^{\gamma} B_{\mathbf{f}}\} = \mathcal{J}(\mathfrak{a}^{\gamma-\epsilon}), \quad \text{for } 0 < \epsilon \ll 1.$$

In particular, we have

$$(7) \quad \delta_{\mathbf{f}} \in V^{\gamma} B_{\mathbf{f}} \quad \text{if and only if} \quad \gamma \leq \text{lct}(\mathfrak{a}),$$

where $\text{lct}(\mathfrak{a})$ is the log canonical threshold of \mathfrak{a} , characterized as $\min\{\lambda > 0 \mid \mathcal{J}(\mathfrak{a}^{\lambda}) \neq \mathcal{O}_X\}$ (we also denote this by $\text{lct}(X, Z)$).

The existence of V -filtrations is closely related to the existence of b -functions. Recall that for every $u \in B_{\mathbf{f}}$, the b -function $b_u(s)$ of u is the monic generator of the ideal

$$(8) \quad \{b(s) \in \mathbf{C}[s] \mid b(s)u \in V^1 \mathcal{R} \cdot u\}.$$

We note that the condition in (8) is equivalent to $b(s)V^0 \mathcal{R} \cdot u \subseteq V^1 \mathcal{R} \cdot u$: this follows easily from Lemmas 7.3 and 7.4 in the Appendix and the fact that for all i and j , we have $t_i \partial_{t_j} \cdot V^0 \mathcal{R} \subseteq V^0 \mathcal{R}$ and $t_i \cdot V^0 \mathcal{R} \subseteq V^1 \mathcal{R}$. It follows from the results in [Kas83] (see also

²This means that there is a positive integer ℓ such that $V^{\gamma} B_{\mathbf{f}}$ has constant value for all γ in an interval of the form $(\frac{i-1}{\ell}, \frac{i}{\ell}]$, with $i \in \mathbf{Z}$.

[BMS06]) that for every $u \in B_{\mathbf{f}}$, the ideal (8) is nonzero and thus $b_u(s)$ is well-defined. Moreover, all its roots are rational. The V -filtration can then be described as

$$(9) \quad V^\gamma B_{\mathbf{f}} = \{u \in B_{\mathbf{f}} \mid \text{all roots of } b_u(s) \text{ are } \leq -\gamma\}.$$

In particular, for $u = \delta_{\mathbf{f}} \in B_{\mathbf{f}}$, the b -function $b_u(s)$ is the Bernstein-Sato polynomial of the ideal \mathfrak{a} , introduced and studied in [BMS06]; this only depends on \mathfrak{a} (not on the choice of f_1, \dots, f_d) and we denote it by $b_Z(s)$. In the case $d = 1$ and $f = f_1$, this is the b -function of a hypersurface, introduced independently by Bernstein and Sato; we also denote it by $b_f(s)$. Note that for any d , it follows from (7) and (9) that

$$(10) \quad \max\{\lambda \in \mathbf{Q} \mid b_Z(\lambda) = 0\} = -\text{lct}(X, Z).$$

Suppose now that $d = 1$, so we have only one nonzero regular function, that we denote f . In this case, Saito introduced in [Sai94] the *microlocal V -filtration* associated to f , defined as follows. Instead of B_f , we consider

$$\tilde{B}_f := \bigoplus_{j \in \mathbf{Z}} \mathcal{O}_X \partial_t^j \delta_f,$$

which is a left module over $\tilde{\mathcal{R}} = \mathcal{D}_X \langle t, \partial_t, \partial_t^{-1} \rangle$. Note that the relation $[\partial_t, t] = 1$ implies $[\partial_t^{-1}, t] = -\partial_t^{-2}$. The action of \mathcal{O}_X and of $\partial_t, \partial_t^{-1}$ on \tilde{B}_f are the obvious ones, while the action of derivations and of t are given by the following analogue of (4): for every $D \in \text{Der}_{\mathbf{C}}(\mathcal{O}_X)$, $h \in \mathcal{O}_X$, and $j \in \mathbf{Z}$, we have

$$(11) \quad D \cdot h \partial_t^j \delta_f = D(h) \partial_t^j \delta_f - h D(f) \partial_t^{j+1} \delta_f \quad \text{and} \quad t \cdot h \partial_t^j \delta_f = f h \partial_t^j \delta_f - j h \partial_t^{j-1} \delta_f.$$

The V -filtration on $\tilde{\mathcal{R}}$ is defined as before: for $m \in \mathbf{Z}$, we have

$$V^m \tilde{\mathcal{R}} = \bigoplus_{i-j \geq m} \mathcal{D}_X t^i \partial_t^j,$$

where this time $i \in \mathbf{Z}_{\geq 0}$ and $j \in \mathbf{Z}$. It is easy to see that

$$V^0 \tilde{\mathcal{R}} = \mathcal{D}_X \langle t, t \partial_t, \partial_t^{-1} \rangle \quad \text{and} \quad V^j \tilde{\mathcal{R}} = \partial_t^{-j} \cdot V^0 \tilde{\mathcal{R}} = V^0 \tilde{\mathcal{R}} \cdot \partial_t^{-j} \quad \text{for all } j \in \mathbf{Z}.$$

The Hodge filtration on \tilde{B}_f is given by

$$F_p \tilde{B}_f = \bigoplus_{i \leq p} \mathcal{O}_X \partial_t^i \delta_f.$$

On the other hand, the microlocal V -filtration on \tilde{B}_f is given by

$$V^\gamma \tilde{B}_f = V^\gamma B_f \oplus \bigoplus_{j \geq 1} \mathcal{O}_X \partial_t^{-j} \delta_f \quad \text{for } \gamma \leq 1$$

and

$$V^\gamma \tilde{B}_f = \partial_t^{-j} V^{\gamma-j} \tilde{B}_f \quad \text{for } \gamma > 1,$$

where $j \in \mathbf{Z}$ is such that $0 < \gamma - j \leq 1$. The following properties follow easily from the properties of the V -filtration on B_f :

- i) $V^1 \tilde{\mathcal{R}} \cdot V^\gamma \tilde{B}_f \subseteq V^{\gamma+1} \tilde{B}_f$ for all $\gamma \in \mathbf{Q}$.
- ii) Every $V^\gamma \tilde{B}_f$ is a finitely generated $V^0 \tilde{\mathcal{R}}$ -module and it generates \tilde{B}_f over $\tilde{\mathcal{R}}$.
- iii) $s + \gamma$ is nilpotent on $\text{Gr}_V^\gamma(\tilde{B}_f)$ for every $\gamma \in \mathbf{Q}$.

A useful property of the microlocal V -filtration is that for every $j \in \mathbf{Z}$ and every $\gamma \in \mathbf{Q}$, multiplication by ∂_t^j gives an isomorphism

$$(12) \quad \partial_t^j: V^\gamma \tilde{B}_f \simeq V^{\gamma-j} \tilde{B}_f$$

(see [Sai94, Lemma 2.2]).

Given $u \in \tilde{B}_f$, the microlocal b -function $\tilde{b}_u(s) \in \mathbf{C}[s]$ is the monic generator of the ideal

$$\{b(s) \in \mathbf{C}[s] \mid b(s)u \in V^1 \tilde{\mathcal{R}} \cdot u\}$$

(the fact that this ideal is nonzero and all roots of $\tilde{b}_u(s)$ are rational, follows from the fact that $V^\gamma \tilde{B}_f \subseteq V^1 \tilde{\mathcal{R}} \cdot u$ for $\gamma \gg 0$). We have the following analogue of (9) describing the microlocal V -filtration in terms of microlocal b -functions:

$$(13) \quad V^\gamma \tilde{B}_f = \{u \in \tilde{B}_f \mid \text{all roots of } \tilde{b}_u(s) \text{ are } \leq -\gamma\}.$$

An important example is that when $u = \delta_f \in \tilde{B}_f$, when we write $\tilde{b}_f(s)$ for $\tilde{b}_{\delta_f}(s)$. If f is not invertible, then it is easy to see that $b_f(s) = b_{\delta_f}(s)$ is divisible by $(s+1)$, and in fact we have

$$\tilde{b}_f(s) = b_f(s)/(s+1)$$

(see [Sai94, Proposition 0.3]).

Under the same assumption that f is not invertible, the negative of the largest root of $b_f(s)/(s+1)$ is the *minimal exponent* $\tilde{\alpha}(f)$, that we also write as $\tilde{\alpha}(H)$ if H is the hypersurface defined by f . Here we make the convention that $\tilde{\alpha}(f) = \infty$ if $b_f(s)/(s+1) = 1$. Note that by (10), we have $\min\{\tilde{\alpha}(H), 1\} = \text{lct}(X, H)$. We also recall that by a result of Saito (see [Sai93, Theorem 0.4]), we have $\tilde{\alpha}(H) > 1$ if and only if H has rational singularities. For a discussion of minimal exponents and basic properties, see [MP20, Section 6]. One property that is very relevant for us is its connection with the V -filtration: it follows from (13) and the fact that $b_f(s)/(s+1) = \tilde{b}_f(s)$ that

$$(14) \quad \tilde{\alpha}(H) = \sup\{\gamma \in \mathbf{Q}_{\geq 0} \mid \delta \in V^\gamma \tilde{B}_f\}.$$

In terms of the V -filtration on B_f , this is equivalent to the fact that for every nonnegative integer q and every rational number $\gamma \in (0, 1]$, we have

$$(15) \quad \tilde{\alpha}(H) \geq q + \gamma \quad \text{if and only if} \quad \partial_t^q \delta_f \in V^\gamma B_f$$

(see [Sai16]).

We will also make use of a local version of the minimal exponent of hypersurfaces. If $f \in \mathcal{O}_X(X)$ and H are as above and $x \in H$, then

$$\tilde{\alpha}_x(f) := \max_{U \ni x} \tilde{\alpha}(f|_U),$$

where the maximum is over all open neighborhoods U of x . With this notation, we have

$$\tilde{\alpha}(f) = \min_{x \in H} \tilde{\alpha}_x(f).$$

3. GENERAL V -FILTRATIONS VIA MICROLOCAL V -FILTRATIONS ALONG HYPERSURFACES

Let X be a smooth, irreducible, complex algebraic variety. In this section, we consider nonzero regular functions $f_1, \dots, f_d \in \mathcal{O}_X(X)$ and let $g = \sum_{i=1}^d f_i y_i \in \mathcal{O}_Y(Y)$, where $Y = X \times \mathbf{A}^d$ and we denote by y_1, \dots, y_d the standard coordinates on \mathbf{A}^d . Our goal is to relate the V -filtration on $B_{\mathbf{f}} = \bigoplus_{\alpha \in \mathbf{Z}_{\geq 0}^d} \mathcal{O}_X \partial_t^\alpha \delta_{\mathbf{f}}$ and the microlocal V -filtration on $\tilde{B}_g = \bigoplus_{j \in \mathbf{Z}} \mathcal{O}_Y \partial_z^j \delta_g$ (note that we denote by z the extra variable that acts on \tilde{B}_g in order to avoid confusion with the variables t_1, \dots, t_d that act on $B_{\mathbf{f}}$).

Note that g is homogeneous of degree 1 with respect to the grading on \mathcal{O}_Y such that \mathcal{O}_X lies in degree 0 and $\deg(y_i) = 1$ for all i . We have a corresponding grading on \mathcal{D}_Y such that $\deg(\partial_{y_i}) = -1$ for all i . Furthermore, on $\mathcal{D}_Y \langle z, \partial_z, \partial_z^{-1} \rangle$ we have a grading such that $\deg(z) = 1 = \deg(\partial_z^{-1})$ and $\deg(\partial_z) = -1$.

We can write

$$\tilde{B}_g = \bigoplus_{j \in \mathbf{Z}} \bigoplus_{\alpha \in \mathbf{Z}_{\geq 0}^d} \mathcal{O}_X y^\alpha \partial_z^j \delta_g.$$

If for every $m \in \mathbf{Z}$ we put

$$\tilde{B}_g^{(m)} = \bigoplus_{\alpha \in \mathbf{Z}_{\geq 0}^d} \mathcal{O}_X y^\alpha \partial_z^{|\alpha|-m} \delta_g,$$

then it follows easily from the formulas (11) and the fact that g is homogeneous of degree 1 that the decomposition $\tilde{B}_g = \bigoplus_{m \in \mathbf{Z}} \tilde{B}_g^{(m)}$ makes \tilde{B}_g a graded $\mathcal{D}_Y \langle z, \partial_z, \partial_z^{-1} \rangle$ -module. Let $\theta_y := \sum_{i=1}^d y_i \partial_{y_i} \in \mathcal{D}_Y$.

Lemma 3.1. *For every $m \in \mathbf{Z}$ and every $u \in \tilde{B}_g^{(m)}$, we have $(\theta_y - s)u = mu$.*

Proof. We may and will assume that $u = hy^\alpha \partial_z^{|\alpha|-m} \delta_g$, for some $h \in \mathcal{O}_X$. On one hand, we have

$$\theta_y \cdot u = \sum_{i=1}^d \alpha_i y_i h y^{\alpha - e_i} \partial_z^{|\alpha|-m} \delta_g - \sum_{i=1}^d y_i f_i h y^\alpha \partial_z^{|\alpha|-m+1} \delta_g = |\alpha| h y^\alpha \partial_z^{|\alpha|-m} \delta_g - g h y^\alpha \partial_z^{|\alpha|-m+1} \delta_g.$$

On the other hand, we have

$$su = -\partial_z z u = -g h y^\alpha \partial_z^{|\alpha|-m+1} \delta_g + (|\alpha| - m) h y^\alpha \partial_z^{|\alpha|-m} \delta_g,$$

and the formula in the lemma follows. \square

Note that $V^\gamma \tilde{B}_g$ is preserved by the action of $\theta_y - s$ for every $\gamma \in \mathbf{Q}$. By Lemma 3.1, the decomposition $\tilde{B}_g = \bigoplus_{m \in \mathbf{Z}} \tilde{B}_g^{(m)}$ is an eigenspace decomposition with respect to the endomorphism $\theta_y - s$. We deduce that we get an induced decomposition

$$V^\gamma \tilde{B}_g = \bigoplus_{m \in \mathbf{Z}} V^\gamma \tilde{B}_g^{(m)}, \quad \text{where} \quad V^\gamma \tilde{B}_g^{(m)} = V^\gamma \tilde{B}_g \cap \tilde{B}_g^{(m)}.$$

We get a corresponding decomposition

$$\mathrm{Gr}_V^\gamma(\tilde{B}_g) = \bigoplus_{m \in \mathbf{Z}} \mathrm{Gr}_V^\gamma(\tilde{B}_g^{(m)}), \quad \text{where} \quad \mathrm{Gr}_V^\gamma(\tilde{B}_g^{(m)}) = V^\gamma \tilde{B}_g^{(m)} / V^{>\gamma} \tilde{B}_g^{(m)}.$$

Finally, we note that it follows from (12) that

$$(16) \quad V^\gamma \tilde{B}_g^{(m)} = \partial_z^{-m} V^{\gamma-m} \tilde{B}_g^{(0)} \quad \text{for all} \quad \gamma \in \mathbf{Q}, m \in \mathbf{Z}.$$

We now define the map that will allow us to compare the V -filtration on $B_{\mathbf{f}}$ with the microlocal V -filtration on \tilde{B}_g . Let $\varphi: \tilde{B}_g \rightarrow B_{\mathbf{f}}$ be the unique \mathcal{O}_X -linear map such that

$$(17) \quad \varphi(y^\alpha \partial_z^j \delta_g) = \partial_t^\alpha \delta_{\mathbf{f}} \quad \text{for all } \alpha \in \mathbf{Z}_{\geq 0}^d, j \in \mathbf{Z}.$$

It is clear from the definition that for every $m \in \mathbf{Z}$, φ induces an isomorphism of \mathcal{O}_X -modules $\tilde{B}_g^{(m)} \simeq B_{\mathbf{f}}$. We collect in the following proposition some basic properties of φ .

Proposition 3.2. *With the above notation, the following hold:*

- i) *The map φ is \mathcal{D}_X -linear.*
- ii) *We have $\varphi(\partial_z u) = \varphi(u) = \varphi(\partial_z^{-1} u)$ for every $u \in \tilde{B}_g$.*
- iii) *We have $\varphi(y_i u) = \partial_{t_i} \varphi(u)$ for every $u \in \tilde{B}_g$ and $1 \leq i \leq d$.*
- iv) *We have $\varphi(\partial_{y_i} u) = -t_i \varphi(u)$ for every $u \in \tilde{B}_g$ and $1 \leq i \leq d$.*
- v) *We have $\varphi(su) = (s - m)\varphi(u)$ for every $u \in \tilde{B}_g^{(m)}$, where $m \in \mathbf{Z}$.*

Proof. For i), since φ is \mathcal{O}_X -linear by definition, it is enough to show that $\varphi(Du) = D\varphi(u)$ for every $D \in \text{Der}_{\mathbf{C}}(\mathcal{O}_X)$ and every $u \in \tilde{B}_g$. We may and will assume that $u = hy^\alpha \partial_z^j \delta_g$ for some $h \in \mathcal{O}_X$, $\alpha \in \mathbf{Z}_{\geq 0}^d$ and $j \in \mathbf{Z}$. In this case, we have

$$\begin{aligned} \varphi(Du) &= \varphi(D(h)y^\alpha \partial_z^j \delta_g - hD(g)y^\alpha \partial_z^{j+1} \delta_g) = \varphi(D(h)y^\alpha \partial_z^j \delta_g - \sum_{i=1}^d hD(f_i)y^{\alpha+e_i} \partial_z^{j+1} \delta_g) \\ &= D(h)\partial_t^\alpha \delta_{\mathbf{f}} - \sum_{i=1}^d hD(f_i)\partial_t^{\alpha+e_i} \delta_{\mathbf{f}} = D \cdot h\partial_t^\alpha \delta_{\mathbf{f}} = D\varphi(u). \end{aligned}$$

This completes the proof of i).

The assertions in ii) and iii) follow directly from the definition. In order to prove iv), we may assume that $u = hy^\alpha \partial_z^j \delta_g$ for some $h \in \mathcal{O}_X$, $\alpha \in \mathbf{Z}_{\geq 0}^d$ and $j \in \mathbf{Z}$. We then have

$$\begin{aligned} \varphi(\partial_{y_i} u) &= \varphi(h\alpha_i y^{\alpha-e_i} \partial_z^j \delta_g - f_i h y^\alpha \partial_z^{j+1} \delta_g) \\ &= \alpha_i h \partial_t^{\alpha-e_i} \delta_{\mathbf{f}} - f_i h \partial_t^\alpha \delta_{\mathbf{f}} = -t_i \cdot h \partial_t^\alpha \delta_{\mathbf{f}} = -t_i \varphi(u). \end{aligned}$$

We thus obtain the assertion in iv).

Finally, in order to prove v), we note that by Lemma 3.1, if $u \in \tilde{B}_g^{(m)}$, then $su = (\theta_y - m)u$, hence using iii) and iv), we have

$$\varphi(su) = \varphi((\theta_y - m)u) = \sum_{i=1}^d \varphi(y_i \partial_{y_i} u) - m\varphi(u) = - \sum_{i=1}^d \partial_{t_i} t_i \varphi(u) - m\varphi(u) = (s - m)\varphi(u).$$

□

We now come to the main result of this section. Let us denote by φ_0 the restriction of φ to $\tilde{B}_g^{(0)}$. Note that since φ_0 is bijective, the assertion in the next theorem together with (16) say that the V -filtration on $B_{\mathbf{f}}$ and the microlocal V -filtration on \tilde{B}_g determine each other.

Theorem 3.3. *With the above notation, for every $\gamma \in \mathbf{Q}$, we have*

$$V^\gamma \tilde{B}_g^{(0)} = \varphi_0^{-1}(V^\gamma B_{\mathbf{f}}).$$

Proof. The argument is similar to that proving the uniqueness of V -filtrations (see for example [Sai88, Lemme 3.1.2]). Recall that we write

$$\mathcal{R} = \mathcal{D}_X \langle t_1, \dots, t_d, \partial_{t_1}, \dots, \partial_{t_d} \rangle \quad \text{and} \quad \tilde{\mathcal{R}} = \mathcal{D}_Y \langle z, \partial_z, \partial_z^{-1} \rangle.$$

Let's prove first the inclusion

$$(18) \quad V^\gamma B_{\mathbf{f}} \subseteq W^\gamma B_{\mathbf{f}} := \varphi_0(V^\gamma \tilde{B}_g^{(0)}) \quad \text{for all } \gamma \in \mathbf{Q}.$$

Note that by definition $W^\bullet B_{\mathbf{f}}$ is an exhaustive, decreasing filtration indexed by rational numbers, which is discrete and left continuous (since the microlocal filtration on \tilde{B}_g has these properties) and Proposition 3.2i) implies that each $W^\gamma B_{\mathbf{f}}$ is a \mathcal{D}_X -submodule of $B_{\mathbf{f}}$. This filtration also satisfies

$$(19) \quad t_i \cdot W^\gamma B_{\mathbf{f}} \subseteq W^{\gamma+1} B_{\mathbf{f}} \quad \text{for all } \gamma \in \mathbf{Q}, 1 \leq i \leq d.$$

Indeed, note that if $u \in V^\gamma \tilde{B}_g^{(0)}$, then $-\partial_z^{-1} \partial_{y_i} u \in V^{\gamma+1} \tilde{B}_g^{(0)}$ and it follows from properties ii) and iv) in Proposition 3.2 that

$$t_i \cdot \varphi_0(u) = \varphi_0(-\partial_z^{-1} \partial_{y_i} u) \in W^{\gamma+1} B_{\mathbf{f}}.$$

We also have

$$(20) \quad \partial_{t_i} \cdot W^\gamma B_{\mathbf{f}} \subseteq W^{\gamma-1} B_{\mathbf{f}} \quad \text{for all } \gamma \in \mathbf{Q}, 1 \leq i \leq d.$$

Indeed, if $u \in V^\gamma \tilde{B}_g^{(0)}$, then $\partial_z y_i u \in V^{\gamma-1} \tilde{B}_g^{(0)}$, and it follows from properties ii) and iii) in Proposition 3.2 that

$$\partial_{t_i} \cdot \varphi_0(u) = \varphi_0(\partial_z y_i u) \in W^{\gamma-1} B_{\mathbf{f}}.$$

In particular, we see that each $W^\gamma B_{\mathbf{f}}$ is a $V^0 \mathcal{R}$ -submodule of $B_{\mathbf{f}}$.

Furthermore, for every $\gamma \in \mathbf{Q}$, we have

$$(21) \quad s + \gamma \quad \text{is nilpotent on} \quad \text{Gr}_W^\gamma(B_{\mathbf{f}}).$$

Indeed, assertion v) in Proposition 3.2 gives $\varphi(su) = s\varphi(u)$ for every $u \in \tilde{B}_g^{(0)}$ and we know that $s + \gamma$ is nilpotent on $\text{Gr}_V^\gamma(\tilde{B}_g)$.

We can now prove the inclusion (18). If γ, γ' are distinct rational numbers, then both $s + \gamma$ and $s + \gamma'$ are nilpotent on

$$(22) \quad \frac{V^\gamma B_{\mathbf{f}} \cap W^{\gamma'} B_{\mathbf{f}}}{(V^{>\gamma} B_{\mathbf{f}} \cap W^{\gamma'} B_{\mathbf{f}}) + (V^\gamma B_{\mathbf{f}} \cap W^{>\gamma'} B_{\mathbf{f}})}$$

(this follows from (21) and the fact that $s + \gamma$ is nilpotent on $\text{Gr}_V^\gamma(B_{\mathbf{f}})$ by definition of the V -filtration on $B_{\mathbf{f}}$). This implies that the quotient in (22) is 0. We deduce that

$$(23) \quad V^\gamma B_{\mathbf{f}} \subseteq W^\gamma B_{\mathbf{f}} + V^{>\gamma} B_{\mathbf{f}} \quad \text{for all } \gamma \in \mathbf{Q}.$$

Indeed, if $u \in V^\gamma B_{\mathbf{f}}$, since $W^\bullet B_{\mathbf{f}}$ is exhaustive, there is γ' such that $u \in W^{\gamma'} B_{\mathbf{f}}$. If $\gamma' \geq \gamma$, then we are done. Suppose now that $\gamma' < \gamma$. The fact that the quotient in (22) is 0 implies that we can write $u = u_1 + u_2$, with $u_1 \in V^{>\gamma} B_{\mathbf{f}} \cap W^{\gamma'} B_{\mathbf{f}}$ and $u_2 \in V^\gamma B_{\mathbf{f}} \cap W^{>\gamma'} B_{\mathbf{f}}$. Note that u lies in the right-hand side of (23) if and only if u_2 does. Also, we have $u_2 \in V^\gamma B_{\mathbf{f}} \cap W^{\gamma''} B_{\mathbf{f}}$ for some $\gamma'' > \gamma'$. We can repeat the argument with u replaced by u_2 ; since $W^\bullet B_{\mathbf{f}}$ is discrete, we see that after finitely many steps we conclude that $u \in W^\gamma B_{\mathbf{f}} + V^{>\gamma} B_{\mathbf{f}}$.

Using the fact that the filtration $V^\bullet B_{\mathbf{f}}$ is discrete, we deduce from (23) that for every $\gamma, \gamma' \in \mathbf{Q}$, we have

$$(24) \quad V^\gamma B_{\mathbf{f}} \subseteq W^\gamma B_{\mathbf{f}} + V^{\gamma'} B_{\mathbf{f}}.$$

We next note that given $\gamma \in \mathbf{Q}$, it follows from property iii) in the definition of the V -filtration on $B_{\mathbf{f}}$ that there is an integer q_0 such that for every integer $q \geq q_0$, we have

$$V^{\gamma+q}B_{\mathbf{f}} \subseteq V^{q-q_0}\mathcal{R} \cdot V^{\gamma+q_0}B_{\mathbf{f}}.$$

On the other hand, since $V^{\gamma+q_0}B_{\mathbf{f}}$ is a finitely generated $V^0\mathcal{R}$ -module and $W^\bullet B_{\mathbf{f}}$ is exhaustive, there is β such that $V^{\gamma+q_0}B_{\mathbf{f}} \subseteq W^\beta B_{\mathbf{f}}$. By taking q such that $q - q_0 + \beta \geq \gamma$, we conclude that

$$V^\gamma B_{\mathbf{f}} \subseteq W^\gamma B_{\mathbf{f}} + V^{\gamma+q}B_{\mathbf{f}} \subseteq W^\gamma B_{\mathbf{f}} + V^{q-q_0}\mathcal{R} \cdot W^\beta B_{\mathbf{f}} \subseteq W^\gamma B_{\mathbf{f}} + W^{\beta+q-q_0}B_{\mathbf{f}} \subseteq W^\gamma B_{\mathbf{f}},$$

where the first inclusion follows from (24) and the third one follows from (19) and (20). This completes the proof of (18).

In order to complete the proof of the theorem, it is enough to also show that

$$\varphi_0(V^\gamma \tilde{B}_g^{(0)}) \subseteq V^\gamma B_{\mathbf{f}} \quad \text{for all } \gamma \in \mathbf{Q}.$$

In fact, we will prove the equivalent statement that

$$(25) \quad V^\gamma \tilde{B}_g \subseteq U^\gamma \tilde{B}_g := \bigoplus_{m \in \mathbf{Z}} \partial_z^{-m} \varphi_0^{-1}(V^{\gamma-m} B_{\mathbf{f}}) \quad \text{for all } \gamma \in \mathbf{Q}.$$

It is clear from the definition that $U^\bullet \tilde{B}_g$ is an exhaustive, decreasing filtration indexed by rational numbers (since the V -filtration on $B_{\mathbf{f}}$ has these properties). Moreover, if ℓ is a positive integer such that $V^\gamma B_{\mathbf{f}}$ is constant for γ in each interval of the form $(\frac{i-1}{\ell}, \frac{i}{\ell}]$, with $i \in \mathbf{Z}$, then it follows from the definition that $U^\gamma \tilde{B}_g$ is constant for γ in such an interval. Therefore $U^\bullet \tilde{B}_g$ is discrete and left continuous.

Note next that by Proposition 3.2i), every $U^\gamma \tilde{B}_g$ is a \mathcal{D}_X -submodule of \tilde{B}_g . Moreover, it follows directly from the definition that

$$\partial_z^j \cdot U^\gamma \tilde{B}_g \subseteq U^{\gamma-j} \tilde{B}_g \quad \text{for all } \gamma \in \mathbf{Q}, j \in \mathbf{Z}.$$

We also have

$$(26) \quad z \cdot U^\gamma \tilde{B}_g \subseteq U^{\gamma+1} \tilde{B}_g \quad \text{for all } \gamma \in \mathbf{Q}.$$

Indeed, if $u \in \varphi_0^{-1}(V^{\gamma-m} B_{\mathbf{f}})$ for some $m \in \mathbf{Z}$, then using the fact that $[z, \partial_z^{-m}] = m \partial_z^{-m-1}$ (see Lemma 7.1 in the Appendix), we have

$$z \partial_z^{-m} u = \partial_z^{-m} (zu + m \partial_z^{-1} u) = \partial_z^{-m-1} (\partial_z z u + m u) = \partial_z^{-m-1} (-su + mu).$$

Note that $-su + mu \in \tilde{B}_g^{(0)}$ and using Proposition 3.2 we see that

$$\varphi_0(-su + mu) = -(s-m)\varphi_0(u) \in V^{\gamma-m} B_{\mathbf{f}},$$

hence $z \partial_z^{-m} u \in U^{\gamma+1} \tilde{B}_g$, proving (26). In particular, we see that each $U^\gamma \tilde{B}_g$ is a $V^0 \tilde{\mathcal{R}}$ -module.

Finally, $s + \gamma$ is nilpotent on $\text{Gr}_U^\gamma(\tilde{B}_g)$ for every $\gamma \in \mathbf{Q}$. Indeed, suppose that $u_0 \in \varphi_0^{-1}(V^{\gamma-m} B_{\mathbf{f}})$. In this case

$$\varphi_0((s + \gamma - m)^N u) = (s + \gamma - m)^N \varphi_0(u) \in V^{>\gamma-m} B_{\mathbf{f}}$$

for $N \gg 0$, where the equality follows from Proposition 3.2v). Since $P(s) \partial_z^{-m} = \partial_z^{-m} P(s-m)$ for every $P \in \mathbf{C}[s]$ (see Lemma 7.3 in Appendix), it follows that

$$(s + \gamma)^N \partial_z^{-m} u = \partial_z^{-m} (s + \gamma - m)^N u \in U^{>\gamma} \tilde{B}_g \quad \text{for } N \gg 0.$$

We can now prove the inclusion (25). Since the argument is very similar to that we used in the proof of (18), we omit some of the details. First, we see that

$$(27) \quad \frac{V^\gamma \tilde{B}_g \cap U^{\gamma'} \tilde{B}_g}{(V^{>\gamma} \tilde{B}_g \cap U^{\gamma'} \tilde{B}_g) + (V^\gamma \tilde{B}_g + U^{>\gamma'} \tilde{B}_g)} = 0 \quad \text{for } \gamma \neq \gamma'$$

using the fact that both $s + \gamma$ and $s + \gamma'$ are nilpotent on this quotient. As before, we use the fact that $U^\bullet \tilde{B}_g$ is exhaustive and discrete and $V^\bullet \tilde{B}_g$ is discrete to deduce from (27) that for every $\gamma, \gamma' \in \mathbf{Q}$, we have

$$(28) \quad V^\gamma \tilde{B}_g \subseteq U^\gamma \tilde{B}_g + V^{\gamma'} \tilde{B}_g.$$

Let us fix now $\gamma \in \mathbf{Q}$. Since $V^0 \tilde{B}_g$ is a finitely generated $V^0 \tilde{\mathcal{R}}$ -module and each $U^\beta \tilde{B}_g$ is a $V^0 \tilde{\mathcal{R}}$ -module, it follows that there is $\beta \in \mathbf{Q}$ such that $V^0 \tilde{B}_g \subseteq U^\beta \tilde{B}_g$. If we take $\gamma' \in \mathbf{Z}$ such that $\gamma' + \beta \geq \gamma$, then using (28) and (12) we conclude that

$$\begin{aligned} V^\gamma \tilde{B}_g &\subseteq U^\gamma \tilde{B}_g + V^{\gamma'} \tilde{B}_g = U^\gamma \tilde{B}_g + \partial_z^{-\gamma'} \cdot V^0 \tilde{B}_g \\ &\subseteq U^\gamma \tilde{B}_g + \partial_z^{-\gamma'} \cdot U^\beta \tilde{B}_g \subseteq U^\gamma \tilde{B}_g + U^{\gamma'+\beta} \tilde{B}_g \subseteq U^\gamma \tilde{B}_g. \end{aligned}$$

This completes the proof of the theorem. \square

We end this section with another application of the map φ , relating b -functions (with respect to f_1, \dots, f_d) to microlocal b -functions (with respect to g).

Proposition 3.4. *For every $m \in \mathbf{Z}$ and every $u \in \tilde{B}_g^{(m)}$, we have*

$$\tilde{b}_u(s - m) = b_{\varphi(u)}(s).$$

Proof. By definition of $\tilde{b}_u(s)$, working locally on X , we can find $P \in V^1 \tilde{\mathcal{R}} = \sum_{a-b \geq 1} \mathcal{D}_Y z^a \partial_z^b$ such that

$$(29) \quad \tilde{b}_u(s)u = Pu.$$

We may and will assume that $\deg(P) = 0$. This implies that we can write $P = \sum_{i=1}^d \partial_{y_i} P_i$ for some P_1, \dots, P_d of degree 1. If we put $Q_i = \partial_z P_i$ for all i , then $P = \sum_{i=1}^d \partial_{y_i} \partial_z^{-1} Q_i$. Applying φ to (29), we obtain

$$(30) \quad \varphi(\tilde{b}_u(s)u) = \varphi(Pu).$$

By Proposition 3.2v), the left-hand side of (30) is equal to $\tilde{b}_u(s - m)\varphi(u)$. On the other hand, it follows from properties ii) and iv) in Proposition 3.2 that the right-hand side of (30) is equal to

$$\sum_{i=1}^d \varphi(\partial_{y_i} \partial_z^{-1} Q_i u) = - \sum_{i=1}^d t_i \cdot \varphi(Q_i u).$$

Note that for every i , we have

$$Q_i \in \sum_{\alpha, \beta, a, b} \mathcal{D}_X y^\alpha \partial_y^\beta z^a \partial_z^b,$$

where $a, b \in \mathbf{Z}$ are such that $a \geq b$ and $a \geq 0$, while $\alpha, \beta \in \mathbf{Z}^d$ are such that $|\alpha| \leq |\beta|$ (this follows from the condition on a and b and the fact that $\deg(Q_i) = 0$). We can write

$z^a \partial_z^b = (z^a \partial_z^a) \partial_z^{b-a} \in \mathbf{C}[s] \cdot \partial_z^{b-a}$ (see Lemma 7.3 in Appendix), hence using Proposition 3.2 we conclude that for every i , we have

$$\varphi(Q_i u) \in \sum_{|\alpha| \leq |\beta|} \mathcal{D}_X[s] \cdot \partial_t^\alpha t^\beta \cdot \varphi(u) \subseteq V^0 \mathcal{R} \cdot \varphi(u).$$

We thus conclude that

$$\tilde{b}_u(s-m) \varphi(u) \in \sum_{i=1}^d t_i \cdot V^0 \mathcal{R} \cdot \varphi(u) \subseteq V^1 \mathcal{R} \cdot \varphi(u),$$

hence by the definition of $b_{\varphi(u)}(s)$, we have

$$(31) \quad b_{\varphi(u)}(s) \text{ divides } \tilde{b}_u(s-m).$$

Going in the opposite direction, it follows from the definition of $b_{\varphi(u)}(s)$ that, working locally on X , there is $T \in V^1 \mathcal{R}$ such that

$$b_{\varphi(u)}(s) \varphi(u) = T \varphi(u).$$

We can write

$$T = \sum_{|\alpha| \geq |\beta|+1} T_{\alpha,\beta} t^\alpha \partial_t^\beta, \quad \text{with } T_{\alpha,\beta} \in \mathcal{D}_X.$$

Let us consider

$$\tilde{T} = \sum_{|\alpha| \geq |\beta|+1} T_{\alpha,\beta} (-y)^\alpha \partial_y^\beta \partial_z^{|\beta|-|\alpha|} \in V^1 \tilde{\mathcal{R}},$$

of degree 0, so $\tilde{T}u \in \tilde{B}_g^{(m)}$. Using Proposition 3.2, we see that

$$\varphi(b_{\varphi(u)}(s+m)u) = b_{\varphi(u)}(s) \varphi(u) = T \varphi(u) = \varphi(\tilde{T}u).$$

Since the restriction of φ to $\tilde{B}_g^{(m)}$ is injective, we conclude that

$$b_{\varphi(u)}(s+m)u = \tilde{T}u.$$

We deduce using the definition of \tilde{b}_u that

$$(32) \quad \tilde{b}_u(s) \text{ divides } b_{\varphi(u)}(s+m).$$

By combining (31) and (32), we see that $\tilde{b}_u(s-m)$ and $b_{\varphi(u)}(s)$ are monic polynomials that divide each other, hence they are equal. \square

Remark 3.5. Note that if we apply the above proposition for $u = \delta_g \in \tilde{B}_g^{(0)}$, then we recover the fact that the Bernstein-Sato polynomial of the ideal (f_1, \dots, f_d) coincides with the microlocal b -function of δ_g (which, as we have mentioned in Section 2, is equal to $b_g(s)/(s+1)$). This is the main result in [Mus22].

4. THE MINIMAL EXPONENT OF A LOCAL COMPLETE INTERSECTION SUBSCHEME

Our goal in this section is to define and study the minimal exponent of a local complete intersection subscheme. Let X be a smooth, irreducible complex algebraic variety and Z a (nonempty) proper closed subscheme of X .

Remark 4.1. We note that in general we have $\text{lct}(X, Z) \leq \text{codim}_X(Z)$. Indeed, the inclusion $Z_{\text{red}} \hookrightarrow Z$ implies $\text{lct}(X, Z) \leq \text{lct}(X, Z_{\text{red}})$, hence we may assume that Z is reduced. If U is an open subset of X such that $U \cap Z$ is smooth and irreducible of codimension r in U , then $\text{lct}(X, Z) \leq \text{lct}(U, Z \cap U) = r$.

Remark 4.2. Note also that if Z is Cohen-Macaulay, of pure codimension r , and $\text{lct}(X, Z) = r$, then Z is reduced. Indeed, if this is not the case, then Z is not generically reduced (being Cohen-Macaulay). It follows that we have an irreducible component Z_0 of Z such that the local ring \mathcal{O}_{Z, Z_0} is not a field; therefore the embedding dimension m of \mathcal{O}_{Z, Z_0} is positive. After possibly replacing X by a suitable open subset that intersects Z_0 nontrivially, we may assume that Z is irreducible, Z_0 is smooth, and there is a smooth, irreducible subvariety W of X of dimension $\dim(Z_0) + m = n - r + m$ such that Z is contained in W and, in fact, the ideal defining Z in W is contained in the ideal $I_{Z_0/W}^2$, where $I_{Z_0/W}$ is the ideal defining Z_0 in W . By considering the exceptional divisor on the blow-up of W along Z_0 and the description of $\text{lct}(W, Z)$ in terms of log resolutions (see [Laz04, Example 9.3.16]), it follows easily that $\text{lct}(W, Z) \leq \text{codim}_W(Z_0)/2 = m/2$. On the other hand, we have $\text{lct}(X, Z) = \text{lct}(W, Z) + \text{codim}_X(W)$ (see for example [Mus02, Proposition 2.6]). Therefore we have

$$\text{lct}(X, Z) = r - m + \frac{m}{2} = r - \frac{m}{2} \leq r - \frac{1}{2}.$$

Suppose now that Z is a local complete intersection, of pure codimension $r \geq 1$ in X . We first consider the case when Z is globally a complete intersection, that is, there are $f_1, \dots, f_r \in \mathcal{O}_X(X)$ such that Z is defined by the ideal generated by f_1, \dots, f_r . In this case we consider the V -filtration on $B_{\mathbf{f}}$. Note that by (7) and Remark 4.1, we have $\delta_{\mathbf{f}} \notin V^\gamma B_{\mathbf{f}}$ for $\gamma > r$.

Definition 4.3. We define the minimal exponent $\tilde{\alpha}(Z)$, as explained in the introduction, by the formula

$$(33) \quad \tilde{\alpha}(Z) = \begin{cases} \sup\{\gamma > 0 \mid \delta_{\mathbf{f}} \in V^\gamma B_{\mathbf{f}}\}, & \text{if } \delta_{\mathbf{f}} \notin V^r B_{\mathbf{f}}; \\ \sup\{r - 1 + q + \gamma \mid F_q B_{\mathbf{f}} \subseteq V^{r-1+\gamma} B_{\mathbf{f}}\}, & \text{if } \delta_{\mathbf{f}} \in V^r B_{\mathbf{f}}, \end{cases}$$

where in the latter case, the supremum is over all nonnegative integers q and all rational numbers $\gamma \in (0, 1]$ with the property that $\partial_t^\beta \delta_{\mathbf{f}} \in V^{r-1+\gamma} B_{\mathbf{f}}$ for all $\beta = (\beta_1, \dots, \beta_r) \in \mathbf{Z}_{\geq 0}^r$, with $\beta_1 + \dots + \beta_r \leq q$. We note that the value of $\tilde{\alpha}(Z)$ does not depend just on Z , but also on X (for the precise way in which it depends on X , see Proposition 4.14 below). Because of this, whenever the ambient variety is not clear from the context, we write $\tilde{\alpha}(X, Z)$ instead of $\tilde{\alpha}(Z)$.

Remark 4.4. Since the V -filtration is left continuous, the supremum in the definition is a maximum, unless $\tilde{\alpha}(Z) = \infty$ (which happens if and only if Z is smooth, see Remark 4.15 below).

Remark 4.5. It follows from the definition and (7) that

$$(34) \quad \text{lct}(X, Z) = \min\{\tilde{\alpha}(Z), r\}.$$

Remark 4.6. If q_1 and q_2 are nonnegative integers and $\gamma_1, \gamma_2 \in (0, 1]$ are rational numbers such that $q_1 + \gamma_1 \geq q_2 + \gamma_2$, and if $F_{q_1} B_{\mathbf{f}} \subseteq V^{r-1+\gamma_1} B_{\mathbf{f}}$, then $F_{q_2} B_{\mathbf{f}} \subseteq V^{r-1+\gamma_2} B_{\mathbf{f}}$. Indeed, this is clear if $q_1 = q_2$, and if this is not the case, then our hypothesis implies $q_2 = q_1 - 1$ and it is enough to show that if $u = \partial_t^\beta \delta_{\mathbf{f}}$, with $|\beta| \leq q_1 - 1$, then $u \in V^r B_{\mathbf{f}}$. The assumption

implies $\partial_{t_i} u \in V^{r-1} B_{\mathbf{f}}$ for all i and thus $t_i \partial_{t_i} u \in V^r B_{\mathbf{f}}$. We conclude that

$$(s+r)u = (-\partial_{t_1} t_1 - \dots - \partial_{t_r} t_r + r)u = -\sum_{i=1}^r t_i \partial_{t_i} u \in V^r B_{\mathbf{f}}.$$

For every $\gamma \neq r$, since $s+\gamma$ is nilpotent on $\text{Gr}_V^\gamma(B_{\mathbf{f}})$, it follows that $s+r$ is invertible on this graded piece. Since $(s+r)u \in V^r B_{\mathbf{f}}$, using the discreteness of the V -filtration, we conclude that $u \in V^r B_{\mathbf{f}}$.

The same argument shows that in order to have $F_q B_{\mathbf{f}} \subseteq V^{r-1+\gamma} B_{\mathbf{f}}$, it is enough to require $\partial_t^\beta \delta_{\mathbf{f}} \in V^{r-1+\gamma} B_{\mathbf{f}}$ for all β with $|\beta| = q$.

Remark 4.7. Suppose that U_1, \dots, U_N are open subsets of X such that all $Z \cap U_i$ are nonempty and $Z \subseteq U_1 \cup \dots \cup U_N$. Since $\partial_t^\beta \delta_{\mathbf{f}} \in V^\gamma B_{\mathbf{f}}$ if and only if the same containment holds on each U_i (note that the condition automatically holds over $X \setminus Z$ by Remark 2.1), it follows using also the assertion in Remark 4.6 that

$$\tilde{\alpha}(X, Z) = \min_{1 \leq i \leq N} \tilde{\alpha}(U_i, Z \cap U_i).$$

Remark 4.8. The definition of $\tilde{\alpha}(Z)$ does not depend on the choice of f_1, \dots, f_r . By taking an affine open cover of X and using Remark 4.7, we see that it is enough to prove this assertion when X is affine. Suppose now that we have regular functions f_1, \dots, f_r and g_1, \dots, g_r such that $(f_1, \dots, f_r) = (g_1, \dots, g_r)$. The condition $\delta_{\mathbf{f}} \in V^\gamma B_{\mathbf{f}}$ is equivalent to $\text{lct}(X, Z) \geq \gamma$, hence it is independent of the choice of generators for the ideal. We thus only need to show that if $q \in \mathbf{Z}_{\geq 0}$ and $\gamma \in (0, 1]$ is a rational number, then $F_q B_{\mathbf{f}} \subseteq V^{r-1+\gamma} B_{\mathbf{f}}$ if and only if $F_q B_{\mathbf{g}} \subseteq V^{r-1+\gamma} B_{\mathbf{g}}$.

Let us write $g_i = \sum_j a_{i,j} f_j$ for $1 \leq i \leq r$. Note that $D = \det(a_{i,j})$ does not vanish at any point in Z . After replacing X by the complement of the zero-locus of D , we may assume that D is invertible (see Remark 4.7). In this case

$$u: X \times \mathbf{A}^r \rightarrow X \times \mathbf{A}^r, \quad u(x, t_1, \dots, t_r) = \left(x, \sum_j a_{1,j} t_j, \dots, \sum_j a_{r,j} t_j \right)$$

is an isomorphism such that $u(X \times \{0\}) = X \times \{0\}$ and $B_{\mathbf{g}} = u_+ B_{\mathbf{f}}$. We thus have an isomorphism u^* of $\mathcal{R} = \mathcal{D}_X \langle t_1, \dots, t_r, \partial_{t_1}, \dots, \partial_{t_r} \rangle$ that keeps \mathcal{D}_X fixed and maps each t_i to a linear form in t_1, \dots, t_r and an isomorphism of \mathcal{R} -modules $\tau: B_{\mathbf{g}} \rightarrow B_{\mathbf{f}}$ (where we view $B_{\mathbf{f}}$ as an \mathcal{R} -module via u^*). This clearly has the property that $\tau(F_q B_{\mathbf{g}}) = F_q B_{\mathbf{f}}$ for every q and using the uniqueness of the V -filtration, we see that $\tau(V^\gamma B_{\mathbf{g}}) = V^\gamma B_{\mathbf{f}}$ for all $\gamma \in \mathbf{Q}$. It is then clear that we have $F_q B_{\mathbf{f}} \subseteq V^{r-1+\gamma} B_{\mathbf{f}}$ if and only if $F_q B_{\mathbf{g}} \subseteq V^{r-1+\gamma} B_{\mathbf{g}}$.

Suppose now that Z is an arbitrary local complete intersection closed subscheme of X , of pure codimension $r \geq 1$. We can find open subsets U_1, \dots, U_N of X with $Z \subseteq \bigcup_{i=1}^N U_i$ such that each $Z \cap U_i$ is nonempty and defined in U_i by an ideal generated by r regular functions on U_i . In particular, each $\tilde{\alpha}(U_i, Z \cap U_i)$ is well-defined.

Definition 4.9. With the above notation, the *minimal exponent* of Z is

$$\tilde{\alpha}(Z) = \tilde{\alpha}(X, Z) := \min_{1 \leq i \leq N} \tilde{\alpha}(U_i, Z \cap U_i).$$

Remark 4.10. It is easy to see, using Remark 4.7, that the definition is independent of the choice of open subsets U_1, \dots, U_N . Moreover, given any open subsets V_1, \dots, V_m of X , with

$Z \subseteq \bigcup_{j=1}^m V_j$ such that each $Z \cap V_j$ is nonempty, we have

$$\tilde{\alpha}(X, Z) = \min_{1 \leq j \leq m} \tilde{\alpha}(V_j, Z \cap V_j).$$

Remark 4.11. In the case of hypersurfaces (that is, $r = 1$), we recover the usual definition of the minimal exponent by (15).

Proposition 4.12. *If $\pi: Y \rightarrow X$ is a surjective smooth morphism of smooth, irreducible varieties, and Z is a local complete intersection closed subscheme of X , of pure codimension r , then $\tilde{\alpha}(X, Z) = \tilde{\alpha}(Y, \pi^{-1}(Z))$.*

Proof. We may and will assume that Z is defined in X by the ideal generated by $f_1, \dots, f_r \in \mathcal{O}_X(X)$ and let $g_i = f_i \circ \pi$ for $1 \leq i \leq r$. Using the fact that π is smooth, it is then straightforward to see that we have an isomorphism

$$B_{\mathbf{g}} \simeq \pi^* B_{\mathbf{f}}$$

such that for every $p \in \mathbf{Z}_{\geq 0}$ and every $\alpha \in \mathbf{Q}$, we get

$$F_p B_{\mathbf{g}} \simeq \pi^* F_p B_{\mathbf{f}} \quad \text{and} \quad V^\alpha B_{\mathbf{g}} \simeq \pi^* V^\alpha B_{\mathbf{f}}.$$

The assertion in the proposition then follows directly from the definition of the minimal exponent. \square

Our next goal is to describe the minimal exponent of Z via the minimal exponent of a hypersurface. Suppose that Z is a nonempty closed subscheme of X , of pure codimension $r \geq 1$, whose ideal is generated by $f_1, \dots, f_r \in \mathcal{O}_X(X)$. We put $g = \sum_{i=1}^r f_i y_i \in \mathcal{O}_X(X)[y_1, \dots, y_r]$. Let $U = X \times (\mathbf{A}^r \setminus \{0\}) \subseteq Y = X \times \mathbf{A}^r$. We will freely use the notation in Section 3. The following is the key observation:

Lemma 4.13. *If $\gamma \in \mathbf{Q}$ and $\alpha \in \mathbf{Z}_{\geq 0}^r$ are such that $y^\alpha \partial_z^{|\alpha|} \delta_g \in V^\gamma \tilde{B}_g \setminus V^{>\gamma} \tilde{B}_g$ and $y^\alpha \partial_z^{|\alpha|} \delta_g|_U \in V^{>\gamma} \tilde{B}_g|_U$, then $\gamma \geq r$ and $\gamma \in \mathbf{Z}$.*

Proof. By assumption, if u is the class of $y^\alpha \partial_z^{|\alpha|} \delta_g$ in $\text{Gr}_V^\gamma(\tilde{B}_g)$, then $u \neq 0$, but there is N such that $(y_1, \dots, y_r)^N u = 0$. This implies that there is $\beta \in \mathbf{Z}_{\geq 0}$ such that $v = y^\beta u \neq 0$, but $(y_1, \dots, y_r)v = 0$. Note that $u \in \text{Gr}_V^\gamma(\tilde{B}_g^{(0)})$, hence $v \in \text{Gr}_V^\gamma(\tilde{B}_g^{(m)})$, where $m = |\beta| \geq 0$. By Lemma 3.1, we have $(\theta_y - s)v = mv$.

On the other hand, since $y_i v = 0$ for all i , we have

$$\theta_y v = \sum_{i=1}^r y_i \partial_{y_i} u_i = -rv + \sum_{i=1}^r \partial_{y_i} y_i v = -rv,$$

hence $(s + m + r)v = \theta_y v + rv = 0$. By definition of the V -filtration, $s + \gamma$ is nilpotent on $\text{Gr}_V^\gamma(\tilde{B}_g)$, and thus $s + \lambda$ is invertible on $\text{Gr}_V^\gamma(\tilde{B}_g)$ for every $\lambda \neq \gamma$. Since $v \neq 0$, we conclude that $\gamma = m + r \geq r$, which completes the proof. \square

We can prove now the equality $\tilde{\alpha}(Z) = \tilde{\alpha}(g|_U)$.

Proof of Theorem 1.1. It is enough to show that for every $\beta \in \mathbf{Q}_{>0}$, we have $\tilde{\alpha}(Z) \geq \beta$ if and only if $\tilde{\alpha}(g|_U) \geq \beta$. We treat separately the cases when $\beta \leq r$ and when $\beta > r$.

If $\beta \leq r$, then by definition we have $\tilde{\alpha}(Z) \geq \beta$ if and only if $\delta_{\mathbf{f}} \in V^\beta B_{\mathbf{f}}$. By Theorem 3.3, this is equivalent to $\delta_g \in V^\beta \tilde{B}_g$. On the other hand, it follows from (14) that $\tilde{\alpha}(g|_U) \geq \beta$ if and

only if $\delta_g \in V^\beta \tilde{B}_g$ on U . It is clear that if $\delta_g \in V^\beta \tilde{B}_g$ then this also holds after restricting to U and we need to show that the converse holds. Arguing by contradiction, let us assume that $\delta_g \in V^\beta \tilde{B}_g$ on U , but $\delta_g \notin V^\beta \tilde{B}_g$. In this case, let $\beta' = \max\{\gamma \in \mathbf{Q}_{\geq 0} \mid \delta_g \in V^\gamma \tilde{B}_g\} < \beta \leq r$. By assumption, we have $\delta_g \in V^{>\beta'} \tilde{B}_g$ on U , hence Lemma 4.13 implies $\beta' \geq r$, a contradiction.

If $\beta > r$, let us write $\beta = r - 1 + q + \gamma$, where q is a positive integer and $\gamma \in (0, 1]$ is a rational number. By definition, we have $\tilde{\alpha}(Z) \geq \beta$ if and only if $\partial_t^\alpha \delta_g \in V^{r-1+\gamma} \tilde{B}_g$ for every $\alpha \in \mathbf{Z}_{\geq 0}^r$ with $|\alpha| \leq q$. By Theorem 3.3, this holds if and only if $y^\alpha \partial_z^{|\alpha|} \delta_g \in V^{r-1+\gamma} \tilde{B}_g$ for all such α . On the other hand, it follows from (14) that $\tilde{\alpha}(g|_U) \geq \beta$ if and only if $\delta_g \in V^\beta \tilde{B}_g$ on U .

Note that $U = U_1 \cup \dots \cup U_r$, where U_i is the complement of the zero-locus of y_i . We thus see that if $y_i^q \partial_z^q \delta_g \in V^{r-1+\gamma} \tilde{B}_g$, we have $\partial_z^q \delta_g \in V^{r-1+\gamma} \tilde{B}_g$ on U_i , and thus $\delta_g \in V^\beta \tilde{B}_g$ on U_i by (12). We conclude that if $\tilde{\alpha}(Z) \geq \beta$, then $\delta_g \in V^\beta \tilde{B}_g$ on U and thus $\tilde{\alpha}(g|_U) \geq \beta$.

In order to prove the converse, we argue by contradiction: we assume that $\delta_g \in V^\beta \tilde{B}_g$ on U , but there is $\alpha \in \mathbf{Z}_{\geq 0}^r$ with $|\alpha| \leq q$ such that $y^\alpha \partial_z^{|\alpha|} \delta_g \notin V^{r-1+\gamma} \tilde{B}_g$. Let

$$\beta' = \max\{\eta \in \mathbf{Q}_{\geq 0} \mid y^\alpha \partial_z^{|\alpha|} \delta_g \in V^\eta \tilde{B}_g\}.$$

Note that $\beta' < r - 1 + \gamma \leq r$. On the other hand, since $\delta_g \in V^\beta \tilde{B}_g$ on U , we have $y^\alpha \partial_z^{|\alpha|} \delta_g \in V^{\beta-|\alpha|} \tilde{B}_g \subseteq V^{>\beta'} \tilde{B}_g$ on U . Applying Lemma 4.13, we get $\beta' \geq r$, a contradiction. \square

Proposition 4.14. *If Z is a local complete intersection scheme of pure dimension and $Z \hookrightarrow X$ is a closed embedding, where X is a smooth, irreducible variety, then $\tilde{\alpha}(X, Z) - \dim(X)$ does not depend on X , but only on Z .*

Proof. Let us consider two embeddings $i: Z \hookrightarrow X$ and $i': Z \hookrightarrow X'$, where both X and X' are smooth irreducible varieties. After comparing both these embeddings with the diagonal embedding $(i, i'): Z \hookrightarrow X \times X'$, we see that we may assume that there is a smooth morphism $p: X' \rightarrow X$ such that $p \circ i' = i$. Given any point $x \in Z$, we can choose regular systems of parameters x_1, \dots, x_n in $\mathcal{O}_{X, i(x)}$ and $p^*(x_1), \dots, p^*(x_n), y_1, \dots, y_m$ in $\mathcal{O}_{X', i'(x)}$ such that y_1, \dots, y_m vanish along $i'(Z)$. In a suitable neighborhood of $i'(x)$, we get an étale morphism $X' \rightarrow X \times \mathbf{A}^m$, given by (p, y_1, \dots, y_m) that maps $i'(Z)$ inside $X \times \{0\}$. After taking a suitable open cover of Z and using the invariance of the minimal exponent under étale morphisms (see Proposition 4.12), we see that it is enough to prove that if $X' = X \times \mathbf{A}^m$ and $i' = (i, 0)$, then $\tilde{\alpha}(X', Z) = \tilde{\alpha}(X, Z) + m$. Of course, arguing by induction on m , we see that it is enough to treat the case $m = 1$.

We may and will assume that X is affine, and the ideal defining Z in X is generated by a regular sequence $f_1, \dots, f_r \in \mathcal{O}_X(X)$. Of course, in this case the ideal defining Z in X' is (f_1, \dots, f_r, z) , where z denotes the coordinate on \mathbf{A}^1 . Using the description of the minimal exponent in Theorem 1.1, we see that it is enough to show that if we put $U = X \times (\mathbf{A}^r \setminus \{0\})$ and $U' = X' \times (\mathbf{A}^{r+1} \setminus \{0\})$, and

$$g = \sum_{i=1}^r f_i y_i \quad \text{and} \quad g' = z y_{r+1} + \sum_{i=1}^r f_i y_i,$$

then $\tilde{\alpha}(g'|_{U'}) = \tilde{\alpha}(g|_U)$. Note that we can write $U' = U'_1 \cup U'_2$, where U'_2 is given by $y_{r+1} \neq 0$ and $U'_1 = U \times \text{Spec}(\mathbf{C}[y_{r+1}, z])$. Note that the hypersurface defined by g' in U'_2 is smooth,

while $g'|_{U'_1} = g|_{U_1} + zy_{r+1}$, hence

$$\tilde{\alpha}(g'|_{U'}) = \tilde{\alpha}(g'|_{U'_1}) = \tilde{\alpha}(g|_{U_1}) + 1,$$

where the second equality follows from the Thom-Sebastiani theorem for minimal exponents (see [Sai94, Theorem 0.8]). \square

We next use the description of the minimal exponent in Theorem 1.1 to prove some basic properties of this invariant. Until the end of this section, we assume that X is a smooth, irreducible, n -dimensional variety and Z is a closed subscheme of X that is locally a complete intersection, of pure codimension r . Recall that if $f \in \mathcal{O}_X(X)$ is nonzero and $x \in X$, then the multiplicity $\text{mult}_x(f)$ is the largest d such that $f \in \mathfrak{m}_x^d$, where \mathfrak{m}_x is the ideal defining x . Before introducing a local version of the minimal exponent, we make the following

Remark 4.15. We have $\tilde{\alpha}(Z) < \infty$ if and only if Z is singular (and in this case we have³ $\tilde{\alpha}(Z) \leq \frac{n+r}{2}$). In order to see this, we may and will assume that Z is defined by the ideal generated by $f_1, \dots, f_r \in \mathcal{O}_X(X)$, and let $g = \sum_{i=1}^r f_i y_i$. We use the fact that by Theorem 1.1, we have $\tilde{\alpha}(Z) = \tilde{\alpha}(g|_U)$, where $U = X \times (\mathbf{A}^r \setminus \{0\})$. If Z is smooth, we may assume that f_1, \dots, f_r are part of a system of coordinates f_1, \dots, f_n on X (that is, df_1, \dots, df_n trivialize Ω_X). In this case it is easy to see⁴ that the singular locus of the hypersurface defined by g is contained in $X \times \{0\}$ and thus $\tilde{\alpha}(g|_U) = \infty$. Conversely, if Z is singular, then we may and will assume that $\text{mult}_x(f_1) \geq 2$ for some $x \in Z$. If $p = (1, 0, \dots, 0) \in \mathbf{A}^r$, then $(x, p) \in U$ and $\text{mult}_{(x,p)}(g) \geq 2$, hence $\tilde{\alpha}(g|_U) \leq \frac{n+r}{2}$ by [MP20, Theorem E(3)].

Suppose now that $x \in Z$ is a fixed point. We define the *minimal exponent of Z at x* as follows: it is clear from the definition that if $V' \subseteq V$ are open neighborhoods of x , then $\tilde{\alpha}(V', Z \cap V') \geq \tilde{\alpha}(V, Z \cap V)$. Moreover, there is V such that for all V' as above, the inequality is an equality. Indeed, otherwise we have a decreasing sequence of open neighborhoods V_i of x such that $(\tilde{\alpha}(V_i, Z \cap V_i))_i$ is a strictly increasing sequence. If $\lim_{i \rightarrow \infty} \tilde{\alpha}(V_i, Z \cap V_i) < \infty$, then we easily get a contradiction using the discreteness of the V -filtration. On the other hand, if $\lim_{i \rightarrow \infty} \tilde{\alpha}(V_i, Z \cap V_i) = \infty$, then it follows from Remark 4.15 that $x \in Z$ is a smooth point, hence it is enough to take V to be a neighborhood of x such that $V \cap Z$ is smooth.

Definition 4.16. Given a point $x \in Z$, we put

$$\tilde{\alpha}_x(Z) := \max_{V \ni x} \tilde{\alpha}(V, Z \cap V),$$

where the maximum is over all open neighborhoods V of x in X . Note that this maximum exists by the previous discussion. Moreover, it follows from Remark 4.15 that $\tilde{\alpha}_x(Z) = \infty$ if and only if x is a smooth point of Z . As before, if the ambient space is not clear from the context, we write $\tilde{\alpha}_x(X, Z)$ instead of $\tilde{\alpha}_x(Z)$.

Remark 4.17. It is a consequence of the definition of the minimal exponent, of the discreteness of the V -filtration, and of Remark 4.15, that the set

$$\{\tilde{\alpha}_x(Z) \mid x \in Z\}$$

is a finite set. It is then clear that

$$\tilde{\alpha}(Z) = \min \{\tilde{\alpha}_x(Z) \mid x \in Z\}.$$

Furthermore, for every $\gamma \in \mathbf{Q}_{>0}$, the set $\{x \in Z \mid \tilde{\alpha}_x(Z) \geq \gamma\}$ is open in Z .

³For a sharper estimate, see Remark 4.21 below.

⁴For a more general statement, see Lemma 4.22 below.

Slightly more generally, if X is a smooth, but possibly disconnected variety, and Z is a local complete intersection closed subscheme of X , with both X and Z pure dimensional, then we can define $\tilde{\alpha}_x(Z)$ for every $x \in Z$ by restricting to the connected component of X that contains x . This is useful, for instance, in the setting of Theorem 1.2, when we do not assume that the fibers of μ are connected.

Before we prove the main properties of the minimal exponent in general, we need to handle one such property in the special case of hypersurfaces.

Lemma 4.18. *The assertion in Theorem 1.2ii) holds when $r = 1$.*

We note that in the presence of a section $s: T \rightarrow X$, the openness assertion in the lemma is [MP20, Theorem E(2)]. However, for our purpose it will be important to have the stronger openness assertion, that does not make reference to a section, since this is the one that will allow us to handle arbitrary codimension. The proof follows closely the approach in [MP20] (which in turn was modeled on the approach to prove the semicontinuity of log canonical thresholds via multiplier ideals, see [Laz04, Example 9.5.41]). However, we include a detailed proof for the benefit of the reader.

The proof of assertion ii₁) makes use of the notion of *Hodge ideals* for \mathbf{Q} -divisors, introduced and studied in [MP19]. For every hypersurface Z in a smooth variety X , every nonnegative integer p , and every positive rational number α , the corresponding Hodge ideal is denoted by $I_p(\alpha Z)$. For $p = 0$, this is just the multiplier ideal $\mathcal{J}((\alpha - \epsilon)Z)$, where $0 < \epsilon \ll 1$, see [MP19, Proposition 9.1] (since Z is a hypersurface, we follow the traditional notation to write $\mathcal{J}(\lambda Z)$ for what we denoted before by $\mathcal{J}(\mathfrak{a}^\lambda)$, where \mathfrak{a} is the ideal defining Z). Moreover, it was shown in [MP19] that many basic properties of multiplier ideals admit extensions to Hodge ideals.

Recall that by definition of the log canonical threshold, we have $\text{lct}(X, Z) \geq \alpha$ if and only if $\mathcal{J}((\alpha - \epsilon)Z) = \mathcal{O}_X$ for $0 < \epsilon \ll 1$. Similarly, it was shown in [MP20, Corollary C] that if Z is reduced, p is a nonnegative integer, and $\alpha \in \mathbf{Q} \cap (0, 1]$, then $\tilde{\alpha}(Z) \geq p + \alpha$ if and only if $I_p(\alpha Z) = \mathcal{O}_X$. On the other hand, if Z is not reduced, then we automatically have $\text{lct}(X, Z) < 1$, hence $\tilde{\alpha}(Z) = \text{lct}(X, Z)$ can be characterized using multiplier ideals.

Proof of Lemma 4.18. We first note that for every $t \in T$, the fiber X_t is a smooth subvariety of the smooth variety X that contains no component of the hypersurface Z . Locally around any $x \in X_t$, we can write X_t as a transverse intersection of $\dim(T)$ smooth hypersurfaces in X , so that successively applying [MP20, Theorem E(1)] to restrict to each of these smooth hypersurfaces, we obtain

$$(35) \quad \tilde{\alpha}_x(X_t, Z_t) \leq \tilde{\alpha}_x(X, Z).$$

We next show that there is a nonempty open subset T_0 of T such that for every $t \in T_0$ and every $x \in X_t$, the inequality in (35) is an equality, thus proving the assertion in ii₂) in our setting. One way to see this is by using the characterization of the minimal exponent in terms of Hodge ideals and multiplier ideals and the fact that there is an open subset T_0 such that

$$\mathcal{J}(\lambda Z_t) = \mathcal{J}(\lambda Z) \cdot \mathcal{O}_{Z_t} \quad \text{for all } \lambda > 0, t \in T_0$$

(see [Laz04, Theorem 9.5.35]) and, assuming that Z is reduced and thus Z_t is also reduced for general $t \in T$, a similar formula holds for Hodge ideals

$$I_p(\lambda Z_t) = I_p(\lambda Z) \cdot \mathcal{O}_{Z_t} \quad \text{for all } p \in \mathbf{Z}_{\geq 0}, \lambda \in \mathbf{Q}_{>0}, t \in T_0$$

(see the last assertion in [MP19, Theorem 13.1]). Alternatively, one can use the characterization of the minimal exponent in terms of the V -filtration in (14) and the results concerning the behavior of the V -filtration with respect to non-characteristic restriction in [DMST06].

We next prove the assertion in ii₁). For every α , let

$$W_\alpha = \{x \in Z \mid \tilde{\alpha}_x(X_{\mu(x)}, Z_{\mu(x)}) \geq \alpha\}.$$

We first note that the assertion in ii₁) makes sense also when T is not assumed to be a smooth variety, but just a (reduced, but not necessarily irreducible) algebraic variety. However, in order to have the statement for such varieties of dimension n , it is enough to prove it for smooth, irreducible, n -dimensional varieties. Indeed, using resolution of singularities, we can find a proper surjective morphism $g: T' \rightarrow T$, with T' n -dimensional and smooth (but possibly disconnected). Consider the Cartesian diagram

$$\begin{array}{ccc} X' & \xrightarrow{h} & X \\ f \downarrow & & \downarrow \mu \\ T' & \xrightarrow{g} & T \end{array}$$

and let $Z' = h^*(Z)$. The assertion follows by noting that

$$W'_\alpha := \{x \in Z' \mid \tilde{\alpha}_x(X'_{f(x)}, Z'_{f(x)}) \geq \alpha\} = h^{-1}(W_\alpha),$$

and thus $Z \setminus W_\alpha = h(Z' \setminus W'_\alpha)$ is closed in Z if W'_α is open in Z' .

We now prove that W_α is open in Z by induction on $\dim(T)$. The assertion is clear if $\dim(T) = 0$, hence we may and will assume that $\dim(T) \geq 1$. We first show that the subset $W_\alpha \subseteq Z$ is constructible. Indeed, note first that if $T_0 \subseteq T$ is a nonempty open subset that satisfies condition ii₂), then

$$W_\alpha \cap \mu^{-1}(T_0) = \{x \in \mu^{-1}(T_0) \cap Z \mid \tilde{\alpha}_x(X, Z) \geq \alpha\}$$

is open in $Z \cap \mu^{-1}(T_0)$. On the other hand, we can apply the induction hypothesis to the morphism $\mu^{-1}(T \setminus T_0) \rightarrow T \setminus T_0$ and the hypersurface $Z \cap \mu^{-1}(T \setminus T_0)$ to conclude that $W_\alpha \cap \mu^{-1}(T \setminus T_0)$ is open in $\mu^{-1}(T \setminus T_0)$ (as we have discussed, the fact that $T \setminus T_0$ might not be smooth is not an issue). Therefore W_α is constructible.

Since W_α is constructible, in order to prove that it is open in Z , it is enough to show that if $W \subseteq Z$ is an irreducible locally closed subvariety of Z , of positive dimension, and $x \in W$ is such that $W \setminus \{x\} \subseteq Z \setminus W_\alpha$, then $x \notin W_\alpha$. Of course, we may assume that W dominates T , since otherwise we are done by induction. Arguing by contradiction, let us assume that $x \in W_\alpha$. In this case it follows from (35) that $\tilde{\alpha}_x(X, Z) \geq \alpha$ and thus there is an open neighborhood V of x in Z such that $\tilde{\alpha}_y(X, Z) \geq \alpha$ for all $y \in V$. On the other hand, if $T_0 \subseteq T$ is a nonempty open subset that satisfies property ii₂), then $W \cap \mu^{-1}(T_0) \cap V$ contains some $y \neq x$. In this case we have

$$\tilde{\alpha}_y(X_{\mu(y)}, Z_{\mu(y)}) = \tilde{\alpha}_y(X, Z) \geq \alpha,$$

hence $y \in W_\alpha$, a contradiction. This completes the proof of ii₁).

The finiteness of the set

$$\{\tilde{\alpha}_x(X_{\mu(x)}, Z_{\mu(x)}) \mid x \in Z\}$$

now follows easily by induction on $\dim(T)$, using the fact that if T_0 satisfies the condition in ii₂), then

$$\{\tilde{\alpha}_x(X_{\mu(x)}, Z_{\mu(x)}) \mid x \in \mu^{-1}(T_0) \cap Z\} \subseteq \{\tilde{\alpha}_x(X, Z) \mid x \in \mu^{-1}(T_0) \cap Z\}$$

and the right-hand side is clearly finite. Finally, if $s: T \rightarrow X$ is a section of π such that $s(T) \subseteq Z$, then

$$\{t \in T \mid \tilde{\alpha}_{s(t)}(X_t, Z_t) \geq \alpha\} = s^{-1}(\{x \in Z \mid \tilde{\alpha}_x(X_{\mu(x)}, Z_{\mu(x)}) \geq \alpha\}),$$

and thus it is open in T . This completes the proof of the lemma. \square

We can now prove the properties of the minimal exponent in arbitrary codimension.

Proof of Theorem 1.2. Since all assertions are local with respect to X , we may and will assume that Z is defined by the ideal generated by $f_1, \dots, f_r \in \mathcal{O}_X(X)$. Let $g = \sum_{i=1}^r f_i y_i \in \mathcal{O}_X(X)[y_1, \dots, y_r]$ and let Z' be the hypersurface defined by g in $U = X \times (\mathbf{A}^r \setminus \{0\})$. Note that $Z \times (\mathbf{A}^r \setminus \{0\}) \subseteq Z'$. The plan is to use Theorem 1.1 to reduce to the case of hypersurfaces. The only subtlety is that while the results concern local minimal exponents, the description provided by Theorem 1.1 is not of a local nature. However, we will go around this issue using the homogeneity of g in y_1, \dots, y_r . More precisely, we have the following

Claim 4.19. If $X_0 \subseteq Z$ is a subset such that $\tilde{\alpha}_{(x,\lambda)}(g) \geq \gamma$ for all $(x, \lambda) \in X_0 \times (\mathbf{A}^r \setminus \{0\})$, then after replacing X by an open neighborhood of X_0 , we may assume that $\tilde{\alpha}(g|_U) \geq \gamma$.

Indeed, consider the canonical projection $\pi: U \rightarrow X \times \mathbf{P}^{r-1}$. Since g is homogeneous with respect to y_1, \dots, y_r , it follows that the set

$$W := \{(x, \lambda) \in Z' \mid \tilde{\alpha}_{(x,\lambda)}(g) < \gamma\}$$

is equal to $\pi^{-1}(W')$, for some subset $W' \subseteq Z \times \mathbf{P}^{r-1}$. Since W is closed in U , we see that W' is closed in $Z \times \mathbf{P}^{r-1}$. By assumption, $W' \cap (X_0 \times \mathbf{P}^{r-1}) = \emptyset$. It follows that if $F \subseteq X$ is the projection of W' , then after replacing X by $X \setminus F$, which is an open neighborhood of X_0 , we have $\tilde{\alpha}(g|_U) \geq \gamma$. This proves the above claim.

Let's begin with the proof of i). Note that the hypothesis implies that Z_H is a complete intersection in H , of pure codimension r , defined by the ideal generated by $f_1|_H, \dots, f_r|_H$. Let $U_H = H \times (\mathbf{A}^r \setminus \{0\})$. If $\gamma = \tilde{\alpha}_x(H, Z_H)$, then after replacing X by a suitable open neighborhood of x , we may assume that $\tilde{\alpha}(H, Z_H) = \gamma$, hence $\tilde{\alpha}(g|_{U_H}) = \gamma$ by Theorem 1.1. In this case, it follows from [MP20, Theorem E(1)] that $\tilde{\alpha}_{(z,\lambda)}(g) \geq \gamma$ for every $z \in Z_H$ and every $\lambda \in \mathbf{A}^r \setminus \{0\}$. We deduce using Claim 4.19 that after possibly replacing X by a neighborhood of Z_H , we have $\tilde{\alpha}(g|_U) \geq \gamma$. Another application of Theorem 1.1 gives $\tilde{\alpha}(X, Z) \geq \gamma$, which completes the proof of i).

We next prove ii). Arguing as in the proof of Lemma 4.18, it is straightforward to see that if ii₁) and ii₂) hold, then the other two assertions hold as well. Let us prove first ii₁). We need to show that for every $x \in Z$, there is an open neighborhood U_x of x such that

$$(36) \quad \tilde{\alpha}_z(X_{\mu(z)}, Z_{\mu(z)}) \geq \alpha := \tilde{\alpha}_x(X_{\mu(x)}, Z_{\mu(x)}) \quad \text{for every } z \in U_x \cap Z.$$

Let φ be the composition $U = X \times (\mathbf{A}^r \setminus \{0\}) \rightarrow X \xrightarrow{\mu} T$. For every $t \in T$, we denote by g_t the restriction of g to $X_t \times \mathbf{A}^r$. After possibly replacing X by an open neighborhood of x , we may and will assume that $\tilde{\alpha}(X_{\mu(x)}, Z_{\mu(x)}) \geq \alpha$. By Theorem 1.1, we have

$$(37) \quad \tilde{\alpha}(g_{\mu(x)}|_{U_{\mu(x)}}) \geq \alpha.$$

Applying Lemma 4.18 for the smooth morphism φ and the hypersurface defined by g in U , we see that the set

$$V_\alpha := \{(z, \lambda) \in Z \times (\mathbf{A}^r \setminus \{0\}) \mid \tilde{\alpha}_{(z, \lambda)}(g_{\mu(z)})|_{U_{\mu(z)}} \geq \alpha\}$$

is open in $Z \times (\mathbf{A}^r \setminus \{0\})$. Note that by (37), we have

$$(38) \quad Z_{\mu(x)} \times (\mathbf{A}^r \setminus \{0\}) \subseteq V_\alpha.$$

Arguing as in the proof of Claim 4.19, we see that after possibly replacing X by an open neighborhood of $Z_{\mu(x)}$, we may assume that $\tilde{\alpha}(g_t) \geq \alpha$ for all $t \in T$ (indeed, V_α is the inverse image of an open subset $W \subseteq Z \times \mathbf{P}^{r-1}$ and we may take the open neighborhood of $Z_{\mu(x)}$ to be the complement in X of the projection of $(X \times \mathbf{P}^{r-1}) \setminus W$ onto the first component). In this case, Theorem 1.1 gives $\tilde{\alpha}(X_t, Z_t) \geq \alpha$ for all $t \in T$. Thus ii₁) holds.

Keeping the same notation, we now prove ii₂). Applying Lemma 4.18 for φ and the hypersurface Z' defined by g , we see that there is an open subset T_0 of T such that for every $t \in T_0$ and every $x \in Z'_t$, we have

$$(39) \quad \tilde{\alpha}_{(x, \lambda)}(g_t) = \tilde{\alpha}_{(x, \lambda)}(g) \quad \text{for all } \lambda \in \mathbf{A}^r \setminus \{0\}.$$

It is enough to show that for every $t \in T_0$ and every $x \in Z_t$, we have

$$(40) \quad \tilde{\alpha}_x(X_t, Z_t) = \tilde{\alpha}_x(X, Z).$$

We fix such t and x and note that we may replace X by any open neighborhood of x . The inequality “ \leq ” in (40) always holds by part i) of the theorem, so we only need to prove that $\tilde{\alpha}_x(X_t, Z_t) \geq \alpha_0 := \tilde{\alpha}_x(X, Z)$. After possibly replacing X by a suitable neighborhood of x , we may and will assume that $\tilde{\alpha}(X, Z) = \alpha_0$, hence Theorem 1.1 gives $\tilde{\alpha}(g|_U) = \alpha_0$. By (39), we thus have $\tilde{\alpha}_{(x', \lambda)}(g_t) \geq \alpha_0$ for every $x' \in Z_t$, and $\lambda \in \mathbf{A}^r \setminus \{0\}$, and another application of Theorem 1.1 gives $\tilde{\alpha}_x(X_t, Z_t) \geq \tilde{\alpha}(X_t, Z_t) \geq \alpha_0$. This completes the proof of ii₂).

Let us prove the inequality in iii). For $\lambda = (\lambda_1, \dots, \lambda_r) \in \mathbf{A}^r \setminus \{0\}$, we put $g_\lambda = \sum_{i=1}^r \lambda_i f_i$. It follows from the assertion in ii₂) that if λ is general, then $\tilde{\alpha}(g_\lambda) \geq \tilde{\alpha}(g|_U) = \tilde{\alpha}(Z)$, where the equality follows from Theorem 1.1. Since $\text{mult}_x(g_\lambda) \geq k$, it follows from [MP20, Theorem E(3)] that $\tilde{\alpha}_x(g_\lambda) \leq \frac{n}{k}$, and thus $\tilde{\alpha}(Z) \leq \frac{n}{k}$. Since the same argument applies to any open neighborhood of x , we get $\tilde{\alpha}_x(Z) \leq \frac{n}{k}$. \square

Remark 4.20. Note that the assertion in Theorem 1.2ii₁) makes sense when T is any (reduced, but not necessarily irreducible) variety. Moreover, the assertion in the general case can be easily reduced to the case when T is smooth using resolution of singularities, as explained in the proof of Lemma 4.18. Furthermore, the same argument implies that in this case, too, the set $\{\tilde{\alpha}_x(X_{\mu(x)}, Z_{\mu(x)}) \mid x \in Z\}$ is finite.

Remark 4.21. We can now see that if X is a smooth, irreducible variety and Z is a local complete intersection closed subscheme of X , of pure dimension, then for every singular point $x \in Z$, we have

$$\tilde{\alpha}_x(X, Z) \leq \dim(X) - \frac{1}{2} \text{embdim}_x(Z),$$

where $\text{embdim}_x(Z) = \dim_{\mathbf{C}} T_x Z$. Indeed, note first that if $d = \text{embdim}_x(Z)$, then after possibly replacing X by an open neighborhood of x , we have a closed embedding $Z \hookrightarrow X'$, where X' is smooth, irreducible, of dimension d . Since $x \in Z$ is a singular point, the ideal

defining Z in X' is contained in \mathfrak{m}_x^2 , where \mathfrak{m}_x is the ideal defining x , hence Theorem 1.2iii) gives $\tilde{\alpha}_x(X', Z) \leq \frac{d}{2}$. In this case Proposition 4.14 implies that

$$\tilde{\alpha}_x(X, Z) = \dim(X) - d + \tilde{\alpha}_x(X', Z) \leq \dim(X) - \frac{d}{2}.$$

Our next goal is to give one nontrivial computation of minimal exponent when $r > 1$. Before doing this, we give an easy lemma describing the singular locus of the hypersurface that we associate to a complete intersection subscheme. We assume that we have global coordinates x_1, \dots, x_n on the smooth variety X (that is, dx_1, \dots, dx_n trivialize Ω_X) and let $\partial_{x_1}, \dots, \partial_{x_n}$ be the corresponding derivations. As usual, we suppose that we have $f_1, \dots, f_r \in \mathcal{O}_X(X)$ that define a closed subscheme Z of X , of pure codimension r , and consider $Y = X \times \mathbf{A}^r$, $U = X \times (\mathbf{A}^r \setminus \{0\})$, and $g = \sum_{i=1}^r f_i y_i \in \mathcal{O}_Y(Y)$. We denote by J_f^t the transpose matrix of the Jacobian matrix $(\partial_{x_j}(f_i))_{i,j}$.

Lemma 4.22. *With the above notation, the singular locus of the hypersurface $V(g)$ defined by g in Y is*

$$V(g)_{\text{sing}} = \bigsqcup_{x \in Z} \{x\} \times W_x,$$

where $W_x = \text{Ker } J_f^t(x)$ is a linear subspace of \mathbf{A}^r of dimension $\dim_{\mathbf{C}}(T_x Z) - \dim(Z)$, and thus

$$V(g|_U)_{\text{sing}} = \bigsqcup_{x \in Z_{\text{sing}}} \{x\} \times (W_x \setminus \{0\}).$$

Proof. The singular locus $V(g)_{\text{sing}}$ is defined by the equations

$$\partial_{y_1}(g) = \dots = \partial_{y_r}(g) = \partial_{x_1}(g) = \dots = \partial_{x_n}(g) = 0$$

(note that these imply $g = 0$ since g is homogeneous of degree 1 in y_1, \dots, y_r). The formula for $V(g)_{\text{sing}}$ follows from the fact that

$$\partial_{y_i}(g) = f_i \quad \text{for } 1 \leq i \leq r \quad \text{and} \quad \partial_{x_j}(g) = \sum_{i=1}^r y_i \partial_{x_j}(f_i) \quad \text{for } 1 \leq j \leq n.$$

We deduce the formula for $\dim(W_x)$ from the fact that the rank of J_f at $x \in Z$ is $n - \dim_{\mathbf{C}}(T_x Z)$ and J_f and J_f^t have the same rank. In particular, we see that $W_x \neq \{0\}$ if and only if $x \in Z_{\text{sing}}$, and we obtain the description of $V(g|_U)_{\text{sing}}$. \square

Example 4.23. Let $f_1, \dots, f_r \in \mathbf{C}[x_1, \dots, x_n]$ be homogeneous polynomials of degree $d \geq 2$ that define a smooth, irreducible variety of codimension r in \mathbf{P}^{n-1} . Therefore the subvariety $Z = V(f_1, \dots, f_r) \subseteq \mathbf{A}^n$ is a complete intersection, with a unique singular point at 0. We will show that

$$\tilde{\alpha}(Z) = \frac{n}{d},$$

generalizing the well-known formula for $r = 1$.

Let $g = \sum_{i=1}^r f_i y_i$ and $U = \mathbf{A}^n \times (\mathbf{A}^r \setminus \{0\})$. We denote by B_U , respectively \tilde{B}_U , the \mathcal{D} -modules on which we have the V -filtration (respectively, the microlocal V -filtration) associated to $g|_U$ and we simply write δ_U for $\delta_{g|_U}$. Note that it follows from Lemma 4.22 and our assumption on Z that the singular locus of $V(g|_U)$ is equal to $\{0\} \times (\mathbf{A}^r \setminus \{0\})$. Recall that by (14), if λ is such that $\delta_U \in V^\lambda \tilde{B}_U$ and its class $\overline{\delta_U} \in \text{Gr}_V^\lambda(\tilde{B}_U)$ is nonzero, then $\lambda = \tilde{\alpha}(g|_U)$. We will show that $\lambda = \frac{n}{d}$.

Recall that for every $\alpha \in [0, 1) \cap \mathbf{Q}$, the filtered \mathcal{D}_U -module $(\mathrm{Gr}_V^\alpha(B_U), F)$ is a filtered direct summand of a mixed Hodge module. In particular, it is regular and quasi-unipotent along every hypersurface in the sense of [Sai88, Section 3.2.1]. Moreover, its support is contained in the singular locus of $V(g|_U)$. On the other hand, we have an isomorphism of filtered \mathcal{D}_U -modules

$$(\mathrm{Gr}_V^\alpha(\tilde{B}_U), F) \simeq (\mathrm{Gr}_V^\alpha(B_U), F)$$

by [Sai94, (2.1.4)]. Finally, if we write $\lambda = k + \alpha$, where $k \in \mathbf{Z}$ and $\alpha \in [0, 1)$, it follows from [Sai94, (2.2.3)] that we have a filtered isomorphism

$$\partial_t^k : (\mathrm{Gr}_V^\lambda(\tilde{B}_U), F) \rightarrow (\mathrm{Gr}_V^\alpha(\tilde{B}_U), F[k]),$$

where $F[k]$ is the shifted filtration $F[k]_p = F_{p+k}$. We thus conclude that $(\mathrm{Gr}_V^\lambda(\tilde{B}_U), F)$ is regular and quasi-unipotent along every hypersurface; moreover, its support is contained in the subset defined by (x_1, \dots, x_n) .

Note that by definition of the Hodge filtration on \tilde{B}_U , we have $\delta_U \in F_0 \tilde{B}_U$ and $F_{-1} \tilde{B}_U = \bigoplus_{i \leq -1} \mathcal{O}_U \partial_t^i \delta_U$. Since $\delta_U \in V^\lambda \tilde{B}_U$, it follows that $F_{-1} \tilde{B}_U \subseteq \sum_{i \leq -1} \partial_t^i \tilde{B}_U \subseteq V^{\lambda+1} \tilde{B}_U$. We thus see that $F_{-1} \mathrm{Gr}_V^\lambda(\tilde{B}_U) = 0$.

Since $(\mathrm{Gr}_V^\lambda(\tilde{B}_U), F)$ is regular and quasi-unipotent along every hypersurface, with support contained in the zero-locus of x_1, \dots, x_n , it follows from [Sai88, Lemme 3.2.6] that x_1, \dots, x_n annihilate the first nonzero piece of the Hodge filtration on $\mathrm{Gr}_V^\lambda(\tilde{B}_U)$. In particular, if $\theta_x = \sum_{i=1}^n x_i \partial_{x_i}$, so that $\theta_x + n = \sum_{i=1}^n \partial_{x_i} x_i$, we see that $(\theta_x + n) \overline{\delta_U} = 0$.

Note that by (4), we have

$$\theta_x \delta_U = - \sum_{i=1}^n x_i \partial_{x_i}(g) \partial_t \delta_U = -dg \partial_t \delta_U,$$

where the second equality follows from the fact that g is homogeneous of degree d with respect to x_1, \dots, x_n . On the other hand, using again (4), we have

$$s \delta_U = -\partial_t t \delta_U = -g \partial_t \delta_U.$$

We thus conclude that

$$(\theta_x + n) \overline{\delta_U} = (n + sd) \overline{\delta_U} = 0.$$

Since $s + \lambda$ is nilpotent on $\mathrm{Gr}_V^\lambda(\tilde{B}_U)$, this implies $\lambda = \frac{n}{d}$.

5. THE MINIMAL EXPONENT AND THE HODGE FILTRATION ON LOCAL COHOMOLOGY

In this section we give the proofs of Theorems 1.3 and 1.4 by making use of a key result from [CD23], saying that, under the assumptions of Theorem 1.4, we have an isomorphism of filtered \mathcal{D}_X -modules

$$(41) \quad \sigma : V^r(B_{\mathbf{f}})/(t_1, \dots, t_r) V^{r-1}(B_{\mathbf{f}}) \xrightarrow{\sim} \mathcal{H}_Z^r(\mathcal{O}_X).$$

For us, it is important to have an explicit description of this isomorphism and this is not easy to obtain from the proof in *loc. cit.* Because of this, we proceed in a roundabout way: we first construct a nonzero morphism of \mathcal{D}_X -modules as in (41) and then use the following lemma that describes the endomorphisms of $\mathcal{H}_Z^r(\mathcal{O}_X)$.

Lemma 5.1. *Let X be a smooth, irreducible complex algebraic variety. If Z is a connected, local complete intersection subscheme of X , of pure codimension r , then the canonical map $\mathbf{C} \rightarrow \mathrm{End}_{\mathcal{D}_X}(\mathcal{H}_Z^r(\mathcal{O}_X))$ is bijective.*

Proof. Let X^{an} denote the complex manifold corresponding to X . In this proof we make use of some standard results on holonomic \mathcal{D} -modules. In order to prove that every endomorphism of $\mathcal{H}_Z^r(\mathcal{O}_X)$ is given by multiplication with a scalar, it is enough to prove that the same property holds for $\mathbf{D}_X(\mathcal{H}_Z^r(\mathcal{O}_X))$, where \mathbf{D}_X is the duality functor for holonomic \mathcal{D}_X -modules (see [HTT08, Chapter 2.6]). Moreover, by the Riemann-Hilbert correspondence (see [HTT08, Chapter 7]), it is enough to show that every endomorphism of the perverse sheaf corresponding to $\mathbf{D}_X(\mathcal{H}_Z^r(\mathcal{O}_X))$ is given by multiplication with a scalar. For an arbitrary Z , this perverse sheaf is ${}^p\mathcal{H}^0(\underline{\mathbf{C}}_{Z^{\text{an}}}[n-r])$, where $n = \dim(X)$. However, since Z is locally a complete intersection, the sheaf $\underline{\mathbf{C}}_{Z^{\text{an}}}[n-r]$ is a perverse sheaf (see [Dim04, Theorem 5.1.20]), and thus ${}^p\mathcal{H}^0(\underline{\mathbf{C}}_{Z^{\text{an}}}[n-r]) = \underline{\mathbf{C}}_{Z^{\text{an}}}[n-r]$. The fact that every endomorphism of $\underline{\mathbf{C}}_{Z^{\text{an}}}$ is given by scalar multiplication follows from the fact that Z is connected. \square

Proof of Theorem 1.4. Since the assertion in the theorem can be checked locally on X , we may and will assume that Z is connected. The key point is to construct an explicit filtered isomorphism as in (41). We first consider the morphism

$$\tau: \mathcal{O}_X[1/f_1 \cdots f_r, s_1, \dots, s_r] \mathbf{f}^{\mathbf{s}} \rightarrow \mathcal{O}_X[1/f_1 \cdots f_r]$$

given by specializing to $s_1 = \dots = s_r = -1$, that is,

$$\tau(P(s_1, \dots, s_r) \mathbf{f}^{\mathbf{s}}) = P(-1, \dots, -1) \frac{1}{f_1 \cdots f_r}.$$

It is clear that this is a morphism of left \mathcal{D}_X -modules.

Via the isomorphism (6), we will identify τ with a morphism $B_{\mathbf{f}}^+ \rightarrow \mathcal{O}_X[1/f_1 \cdots f_r]$. Using (5) and the fact that $Q_m(-1) = m!$, we can describe τ by

$$(42) \quad \tau\left(\sum_{\beta} h_{\beta} \partial_t^{\beta} \delta_{\mathbf{f}}\right) = \sum_{\beta} \frac{\beta_1! \cdots \beta_r! h_{\beta}}{f_1^{\beta_1+1} \cdots f_r^{\beta_r+1}},$$

where $h_{\beta} \in \mathcal{O}_X[1/f_1 \cdots f_r]$ for all β .

It is clear from the definition that τ vanishes on $\sum_{i=1}^r (s_i + 1) B_{\mathbf{f}}^+$. Let us consider the image of $t_i B_{\mathbf{f}}$. Note that by Lemma 7.4, if $\beta_i \geq 1$, then

$$t_i Q_{\beta}(s_1, \dots, s_r) \mathbf{f}^{\mathbf{s}} = (s_i + 1) Q_{\beta'}(s_1, \dots, s_r) f_i \cdot \mathbf{f}^{\mathbf{s}} \in \text{Ker}(\tau),$$

where $\beta' = (\beta_1, \dots, \beta_i - 1, \dots, \beta_r)$. On the other hand, if $\sum_{\beta} h_{\beta} \partial_t^{\beta} \delta_{\mathbf{f}} \in B_{\mathbf{f}}$ (so $h_{\beta} \in \mathcal{O}_X$ for all β) and if β is such that $\beta_i = 0$, then

$$\tau(t_i h_{\beta} \partial_t^{\beta} \delta_{\mathbf{f}}) = \tau(f_i h_{\beta} \partial_t^{\beta} \delta_{\mathbf{f}}) = h_{\beta} \prod_{j \neq i} \frac{\beta_j!}{f_j^{\beta_j+1}} \in \mathcal{O}_X[1/f_1 \cdots \widehat{f_i} \cdots f_r].$$

We conclude that τ induces a morphism of \mathcal{D}_X -modules

$$\bar{\tau}: V^r B_{\mathbf{f}} / \sum_{i=1}^r t_i \cdot V^{r-1} B_{\mathbf{f}} \rightarrow \mathcal{O}_X[1/f_1 \cdots f_r] / \sum_{i=1}^r \mathcal{O}_X[1/f_1 \cdots \widehat{f_i} \cdots f_r] \simeq \mathcal{H}_Z^r(\mathcal{O}_X).$$

Let us show that $\bar{\tau}$ is surjective. Given $u = \frac{g}{(f_1 \cdots f_r)^m} \in \mathcal{O}_X[1/f_1 \cdots f_r]$, for some $g \in \mathcal{O}_X$ and some $m \geq 1$, we have $u = \tau(v)$, where $v = \frac{g}{(m-1)!^r} \partial_t^{\beta} \delta_{\mathbf{f}}$, where $\beta = (m-1, \dots, m-1)$. By the properties of the V -filtration, we know that we can find $\alpha_1, \dots, \alpha_N \in \mathbf{Q}$, with $\alpha_i < r$ for all i , such that $w := (s + \alpha_1) \cdots (s + \alpha_N) v \in V^r B_{\mathbf{f}}$ (recall that $s = s_1 + \dots + s_r$). Since we can find $p(s)$ and $q(s)$ such that $p(s)(s+r) + q(s) \prod_{i=1}^N (s + \alpha_i) = 1$ and since $(s+r)v \in \sum_{i=1}^r (s_i + 1) B_{\mathbf{f}} \subseteq \text{Ker}(\tau)$, it follows that $u = \tau(v) = \tau(q(s)w) \in \tau(V^r B_{\mathbf{f}})$. Therefore we see that $\bar{\tau}$ is surjective.

As we have already mentioned, it follows from [CD23, Theorems 1.1 and 1.2(b)] that we have the isomorphism of filtered \mathcal{D}_X -modules (41), where the filtration on the left-hand side is induced by the Hodge filtration on $B_{\mathbf{f}}$ and the filtration on the right-hand side is the Hodge filtration that comes from the mixed Hodge module structure on $\mathcal{H}_Z^r(\mathcal{O}_X)$. We recall that our convention is that the Hodge filtration on $B_{\mathbf{f}}$ is defined to be

$$F_q B_{\mathbf{f}} = \bigoplus_{|\alpha| \leq q} \mathcal{O}_X \cdot \partial_t^\alpha \delta_{\mathbf{f}},$$

which differs by a shift by r from the usual convention followed in [CD23]. In particular, $\bar{\tau} \circ \sigma^{-1}$ is a \mathcal{D}_X -linear endomorphism of $\mathcal{H}_Z^r(\mathcal{O}_X)$; hence, by Lemma 5.1, it is given by multiplication with some $\lambda \in \mathbf{C}$. Since the morphism is nonzero (being surjective), it follows that $\lambda \neq 0$, hence $\bar{\tau}$ is an isomorphism. Moreover, since $\bar{\tau} = \lambda \sigma$, we deduce that $\bar{\tau}$ is a filtered isomorphism, too. Therefore the assertion in the theorem follows from the definition of the Hodge filtration on $B_{\mathbf{f}}$ and the description of τ in (42). \square

Proof of Theorem 1.3. We shall prove the equivalent statement that $F_k B_{\mathbf{f}} \subseteq V^r B_{\mathbf{f}}$ if and only if $F_k \mathcal{H}_Z^r \mathcal{O}_X = O_k \mathcal{H}_Z^r \mathcal{O}_X$. The “only if” part is clear: since the elements

$$\left[\frac{1}{f_1^{\alpha_1+1} f_2^{\alpha_2+1} \dots f_r^{\alpha_r+1}} \right] \in \mathcal{H}_Z^r(\mathcal{O}_X)$$

with $\alpha_1, \dots, \alpha_r \geq 0$ and $\alpha_1 + \alpha_2 + \dots + \alpha_r \leq k$ generate $O_k \mathcal{H}_Z^r \mathcal{O}_X$ (see for example [MP22, Lemma 9.2]), the “only if” part follows from Theorem 1.4.

For the reverse implication, we use induction on k . Suppose first that $k = 0$. Since $\left[\frac{1}{f_1 \dots f_r} \right] \in F_0 \mathcal{H}_Z^r(\mathcal{O}_X)$, it follows from Theorem 1.4 that locally on X , we can find $h \in \mathcal{O}_X$ such that $h \delta_{\mathbf{f}} \in V^r B_{\mathbf{f}}$ and $(h - 1)$ lies in the ideal \mathcal{I}_Z defining Z . Therefore $F_0 B_{\mathbf{f}} \subseteq V^r B_{\mathbf{f}}$ at every point of Z (and outside of Z , this is automatic).

Suppose now we know the assertion for $k \geq 0$ and let us prove it for $k + 1$. Since $F_{k+1} \mathcal{H}_Z^r(\mathcal{O}_X) = O_{k+1} \mathcal{H}_Z^r(\mathcal{O}_X)$, it follows from [MP22, Lemma 9.3] that $F_k \mathcal{H}_Z^r(\mathcal{O}_X) = O_k \mathcal{H}_Z^r(\mathcal{O}_X)$, hence by the induction hypothesis we have $\partial_t^\alpha \delta \in V^r B_{\mathbf{f}}$ for all α with $|\alpha| \leq k$. We need to show that $\partial_t^\alpha \delta_{\mathbf{f}} \in V^r B_{\mathbf{f}}$ also for all α with $|\alpha| = k + 1$.

Since $F_{k+1} \mathcal{H}_Z^r(\mathcal{O}_X) = O_{k+1} \mathcal{H}_Z^r(\mathcal{O}_X)$, it follows from Theorem 1.4 that the map

$$\sigma: F_{k+1} V^r B_{\mathbf{f}} \rightarrow \mathrm{Gr}_{k+1}^O(\mathcal{H}_Z^r(\mathcal{O}_Z)), \quad \sum_{|\beta| \leq k+1} h_\beta \partial_t^\beta \delta_{\mathbf{f}} \rightarrow \sum_{|\beta|=k+1} \left[\frac{\beta_1! \dots \beta_r! h_\beta}{f_1^{\beta_1+1} \dots f_r^{\beta_r+1}} \right]$$

is surjective. This implies that working locally on X , for every α with $|\alpha| = k + 1$ we can find

$$u_\alpha = \sum_{|\beta| \leq k+1} h_{\alpha, \beta} \partial_t^\beta \delta_{\mathbf{f}} \in V^r B_{\mathbf{f}}$$

that is mapped by σ to $v_\alpha = \left[\frac{\alpha_1! \alpha_2! \dots \alpha_r!}{f_1^{\alpha_1+1} \dots f_r^{\alpha_r+1}} \right] \in \mathrm{Gr}_{k+1}^O(\mathcal{H}_Z^r(\mathcal{O}_Z))$. Note that since Z is a complete intersection, $\mathrm{Gr}_{k+1}^O \mathcal{H}_Z^r(\mathcal{O}_Z)$ is a free \mathcal{O}_Z -module, with a basis given by the v_α , with $|\alpha| = k + 1$ (see for example [MP22, Lemmas 9.1, 9.2]). This implies that $h_{\alpha, \alpha} - 1$ and $h_{\alpha, \beta}$, for $\alpha \neq \beta$, lie in \mathcal{I}_Z . It is then clear that for every α with $|\alpha| = k + 1$ we have $\partial_t^\alpha \delta_{\mathbf{f}} \in V^r B_{\mathbf{f}}$ at every point of Z (this holds trivially on the complement of Z). This completes the proof of the induction step and thus the proof of the “if” part. \square

We obtain the following consequence to the characterization of local complete intersection Du Bois singularities. We note that the “if” part is [Sch07, Corollary 5.8] and the full equivalence in the case when Z is normal is [Kov99, Theorem 3.6] (note that if Z is normal and local complete intersection, then $\text{lct}(X, Z) = r$ if and only if Z has log canonical singularities by Inversion of Adjunction, see [EM04, Corollary 3.2]). A version of the “only if” implication also appears in [Doh08, Theorem 4.2].

Corollary 5.2. *If X is a smooth, irreducible variety and $Z \hookrightarrow X$ is a local complete intersection closed subscheme of pure codimension r , then Z has Du Bois singularities⁵ if and only if $\text{lct}(X, Z) = r$.*

Proof. Recall first that $\text{lct}(X, Z) \leq r$ and equality implies that Z is reduced: see Remarks 4.1 and 4.2. We may thus assume that Z is reduced. Since Z is locally a complete intersection, it follows from [MP22, Theorem C] that Z has Du Bois singularities if and only if $F_0\mathcal{H}_Z^r(\mathcal{O}_X) = \mathcal{O}_0\mathcal{H}_Z^r(\mathcal{O}_X)$ and this condition is equivalent to $\text{lct}(X, Z) \geq r$ by Theorem 1.3. \square

6. THE MINIMAL EXPONENT AND THE BERNSTEIN-SATO POLYNOMIAL

Let X be a smooth, irreducible, complex algebraic variety and Z a proper (nonempty) closed subscheme of X , defined by the ideal \mathfrak{a} . In what follows we will make use of the notation and definitions introduced in Section 2. Recall, in particular, that we discussed the Bernstein-Sato polynomial $b_Z(s)$ in the case when \mathfrak{a} is generated by nonzero global regular functions f_1, \dots, f_d . The general case can be easily reduced to this one (in fact, to the case when X is affine) since for open subsets U_1, \dots, U_N of X such that $Z \subseteq U_1 \cup \dots \cup U_N$, we have

$$(43) \quad b_Z(s) = \text{LCM}\{b_{Z \cap U_i}(s) \mid 1 \leq i \leq N\}$$

(with the convention that $b_{Z \cap U_i}(s) = 1$ if $Z \cap U_i = \emptyset$).

Proposition 6.1. *If Z is locally a complete intersection in X , of pure codimension r , then $b_Z(-r) = 0$.*

Proof. If U is an open subset of X , then we deduce from (43) that $b_{Z \cap U}(s)$ divides $b_Z(s)$. It follows that after replacing X by a suitable affine open neighborhood of a point in Z , we may assume that X is affine and the ideal \mathfrak{a} defining Z is generated by a regular sequence $f_1, \dots, f_r \in \mathcal{O}_X(X)$. The condition in the definition of $b_Z(s)$ can be reformulated as saying that $b_Z(s)$ is the monic polynomial of minimal degree such that

$$(44) \quad b_Z(s)f_1^{s_1} \cdots f_r^{s_r} \in \sum_{u \in \mathbf{Z}^r, |u|=1} \mathcal{D}_X[s_1, \dots, s_r] \cdot \prod_{u_i < 0} \binom{s_i}{-u_i} f_1^{s_1+u_1} \cdots f_r^{s_r+u_r}$$

(see [BMS06, Section 2.10]). Here we use the isomorphism (6), that maps $\delta_{\mathbf{f}}$ to $f_1^{s_1} \cdots f_r^{s_r}$, such that the action of $-\partial_{t_i} t_i$ corresponds to the action of s_i (so that $s = s_1 + \dots + s_r$). Note also that $\binom{s_i}{-u_i}$ denotes the polynomial $\frac{1}{(-u_i)!} s_i(s_i - 1) \cdots (s_i + u_i - 1)$.

By making $s_1 = \dots = s_r = -1$ in (44), we obtain the following relation in $\mathcal{O}_X[1/f_1 \cdots f_r]$:

$$(45) \quad b_Z(-r) \frac{1}{f_1 \cdots f_r} \in \sum_{u \in \mathbf{Z}^r, |u|=1} \mathcal{D}_X \cdot f_1^{u_1-1} \cdots f_r^{u_r-1}.$$

⁵We note that the condition of having Du Bois singularities assumes, in particular, that Z is reduced.

Note that if $u = (u_1, \dots, u_r)$ is such that $|u| = 1$, then there is i such that $u_i \geq 1$, in which case

$$\mathcal{D}_X \cdot f_1^{u_1-1} \cdots f_r^{u_r-1} \subseteq \mathcal{D}_X \cdot \prod_{j \neq i} f_j^{u_j-1} \subseteq \mathcal{O}_X[1/f_1 \cdots \widehat{f_i} \cdots f_r].$$

Using (45), we thus conclude that

$$b_Z(-r) \frac{1}{f_1 \cdots f_r} \in \sum_{i=1}^r \mathcal{O}_X[1/f_1 \cdots \widehat{f_i} \cdots f_r],$$

hence the class of $b_Z(-r) \frac{1}{f_1 \cdots f_r}$ in the local cohomology $\mathcal{H}_Z^r(\mathcal{O}_X)$ is 0. Since the class of $\frac{1}{f_1 \cdots f_r}$ in the local cohomology is nonzero, we conclude that $b_Z(-r) = 0$. \square

From now on, for the rest of this section, we assume that Z is a local complete intersection closed subscheme of X , of pure codimension $r \geq 1$. The above result motivates the following

Definition 6.2. We denote by $\tilde{\gamma}(Z)$ the negative of the largest root of $b_Z(s)/(s+r)$ (with the convention that $\tilde{\gamma}(Z) = \infty$ if this polynomial is 1).

Remark 6.3. Note that since all roots of $b_Z(s)$ are rational numbers, the same is true for $\tilde{\gamma}(Z)$. Recall also that by formula (10), the largest root of $b_Z(s)$ is $-\text{lct}(X, Z)$, hence

$$(46) \quad \text{lct}(X, Z) = \min \{ \tilde{\gamma}(Z), r \}.$$

Remark 6.4. Since Z is locally a complete intersection in X , of pure codimension r , it follows from [BMS06, Theorem 4] that $\tilde{\gamma}(Z) > r$ if and only if Z has rational singularities (the result in *loc. cit.* requires Z to also be reduced, but the hypothesis $\tilde{\gamma}(Z) > r$ implies $\text{lct}(X, Z) = r$, and thus Z is reduced by Remark 4.2).

We will need the following easy result:

Lemma 6.5. *If $X = \text{Spec}(R)$ is a smooth affine variety and $f_1, \dots, f_r \in R$ form a regular sequence, then for every $k \geq 0$, the sequence t_1, \dots, t_r is regular on the finitely generated $R[t_1, \dots, t_r]$ -module $F_k B_{\mathbf{f}}$.*

Proof. For every $k \geq 0$, the quotient $F_k B_{\mathbf{f}} / F_{k-1} B_{\mathbf{f}}$ is a free R -module. Moreover, each t_i acts on this quotient as multiplication by f_i . The assertion in the lemma thus follows by induction on k , using the fact that if

$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$

is a short exact sequence of $R[t_1, \dots, t_r]$ -modules such that t_1, \dots, t_r is a regular sequence on both M' and M'' , then it is a regular sequence also on M . \square

We can now prove our result relating $\tilde{\gamma}(Z)$ and $\tilde{\alpha}(Z)$.

Proof of Theorem 1.6. After possibly replacing X by suitable affine open subsets, we may and will assume that X is affine and the ideal defining Z is generated by a regular sequence f_1, \dots, f_r . It follows from (34) and (46) that we have

$$\min \{ \tilde{\gamma}(Z), r \} = \text{lct}(X, Z) = \min \{ \tilde{\alpha}(Z), r \}.$$

Therefore both assertions in the theorem hold if $\text{lct}(X, Z) < r$. Hence from now on we may and will assume that $\text{lct}(X, Z) = r$, which in light of (7), is equivalent to $\delta_{\mathbf{f}} \in V^r B_{\mathbf{f}}$. In order to prove the two assertions in the theorem, it is enough to show the following:

- (i) If q is a nonnegative integer and $\gamma \in (0, 1]$ is a rational number such that $\tilde{\gamma}(Z) \geq r - 1 + q + \gamma$, then $\tilde{\alpha}(Z) \geq r - 1 + q + \gamma$; by definition of the minimal exponent, this is equivalent to $\partial_t^\beta \delta_{\mathbf{f}} \in V^{r-1+\gamma} B_{\mathbf{f}}$ for all $\beta \in \mathbf{Z}_{\geq 0}^r$ with $|\beta| \leq q$.
- (ii) If $\gamma' \in (0, 1]$ is a rational number such that $\tilde{\alpha}(\bar{Z}) \geq r + \gamma'$, then $\tilde{\gamma}(Z) \geq r + \gamma'$.

We first prove (i), arguing by induction on $q \geq 0$. If $q = 0$, then we are done, since we are assuming $\delta_{\mathbf{f}} \in V^r B_{\mathbf{f}}$. Suppose now that $q \geq 1$. By the induction hypothesis, it is enough to show that for every $\beta \in \mathbf{Z}_{\geq 0}^r$, with $|\beta| = q - 1$, if $u = \partial_t^\beta \delta_{\mathbf{f}}$, then $\partial_{t_i} u \in V^{r-1+\gamma} B_{\mathbf{f}}$ for $1 \leq i \leq r$. Furthermore, the induction hypothesis gives $u \in V^r B_{\mathbf{f}}$. Let's write $b(s) := b_Z(s) = (s+r)p(s)$.

By definition of $b_Z(s)$, we have

$$(47) \quad b(s)V^0 \mathcal{R} \cdot \delta_{\mathbf{f}} \subseteq V^1 \mathcal{R} \cdot \delta_{\mathbf{f}}.$$

We first note that for every i with $1 \leq i \leq q - 1$, if $\beta' \in \mathbf{Z}_{\geq 0}^r$ is such that $|\beta'| = q - i$, then there is $\beta'' \in \mathbf{Z}_{\geq 0}^r$ with $|\beta''| = q - i - 1$ such that

$$(48) \quad b(s - q + i) \partial_t^{\beta'} V^0 \mathcal{R} \cdot \delta_{\mathbf{f}} \subseteq \partial_t^{\beta''} V^0 \mathcal{R} \cdot \delta_{\mathbf{f}}.$$

Indeed, it follows from Lemma 7.3 that if we take any $\beta'' \in \mathbf{Z}_{\geq 0}^r$ with $|\beta''| = q - i - 1$ and $\beta'_j \geq \beta''_j \geq 0$ for all j , then

$$b(s - q + i) \partial_t^{\beta'} V^0 \mathcal{R} \cdot \delta_{\mathbf{f}} = \partial_t^{\beta'} b(s) V^0 \mathcal{R} \cdot \delta_{\mathbf{f}} \subseteq \partial_t^{\beta'} V^1 \mathcal{R} \cdot \delta_{\mathbf{f}} \subseteq \partial_t^{\beta''} V^0 \mathcal{R} \cdot \delta_{\mathbf{f}}.$$

Applying (48) for all i with $1 \leq i \leq q - 1$, as well as (47), we obtain

$$b(s)b(s-1) \cdots b(s-q+1) \partial_t^\beta \delta_{\mathbf{f}} \subseteq b(s)V^0 \mathcal{R} \cdot \delta_{\mathbf{f}} \subseteq V^1 \mathcal{R} \cdot \delta_{\mathbf{f}} \subseteq V^{r+1} B_{\mathbf{f}}.$$

Note now that we can write

$$b(s)b(s-1) \cdots b(s-q+1) = (s+r)p_1(s),$$

where

$$p_1(s) = (s+r-1) \cdots (s+r-q+1) \cdot \prod_{i=0}^{q-1} p(s-i).$$

The assumption that $\tilde{\gamma}(Z) \geq r - 1 + q + \gamma$ implies that $p_1(s)$ has no roots in the interval $(-r - \gamma, -r]$. Since $(s+r)u \in V^r B_{\mathbf{f}}$ (recall that $u \in V^r B_{\mathbf{f}}$) and $p_1(s)(s+r)u \in V^{r+1} B_{\mathbf{f}}$, we conclude that $(s+r)u \in V^{r+\gamma} B_{\mathbf{f}}$. By assumption, we have $u \in F_{q-1} B_{\mathbf{f}}$, hence $(s+r)u \in F_q V^{r+\gamma} B_{\mathbf{f}}$ (where for every $p \in \mathbf{Z}_{\geq 0}$ and every $\alpha \in \mathbf{Q}$, we put $F_p V^\alpha B_{\mathbf{f}} = F_p B_{\mathbf{f}} \cap V^\alpha B_{\mathbf{f}}$).

Consider now the map

$$(F_q V^{r-1+\gamma} B_{\mathbf{f}})^{\oplus r} \xrightarrow{(t_1, \dots, t_r)} F_q V^{r+\gamma} B_{\mathbf{f}}.$$

Since $\gamma > 0$, it follows from [CD23, Theorem 1.1] that the map is surjective, hence we may write

$$(s+r)u = \sum_{i=1}^r t_i u_i,$$

with $u_i \in F_q V^{r-1+\gamma} B_{\mathbf{f}}$ for $1 \leq i \leq r$. Since $s+r = -\sum_{i=1}^r t_i \partial_{t_i}$, we obtain

$$\sum_{i=1}^r t_i (u_i + \partial_{t_i} u) = 0.$$

Using the fact that t_1, \dots, t_r form a regular sequence on $F_q B_{\mathbf{f}}$ by Lemma 6.5, we conclude that for every i , we have

$$u_i + \partial_{t_i} u \in \sum_{i=1}^r t_i \cdot F_q B_{\mathbf{f}} \subseteq V^r B_{\mathbf{f}},$$

where the inclusion follows from the fact that we already know, by induction, that $F_{q-1} B_{\mathbf{f}} \subseteq V^r B_{\mathbf{f}}$ and thus $F_q B_{\mathbf{f}} \subseteq \sum_{i=1}^r \partial_{t_i} \cdot V^r B_{\mathbf{f}} \subseteq V^{r-1} B_{\mathbf{f}}$. We thus conclude that

$$\partial_{t_i} u = (u_i + \partial_{t_i} u) - u_i \in V^{r-1+\gamma} B_{\mathbf{f}}.$$

This completes the proof of (i).

We next prove (ii). We will make use of the results in Section 3. By definition of $\tilde{\alpha}(Z)$, we know that $\partial_{t_i} \delta_{\mathbf{f}} \in V^{r-1+\gamma'} B_{\mathbf{f}}$ for $1 \leq i \leq r$. Theorem 3.3 (see also equation (16)) thus gives $y_i \delta_g \in V^{r+\gamma'} \tilde{B}_g$ for $1 \leq i \leq r$. By Lemma 3.1, we have $v := (s+r)\delta_g = \sum_{i=1}^r \partial_{y_i} y_i \delta_g \in V^{r+\gamma'} \tilde{B}_g$. Using the description (13), we deduce that all roots of $\tilde{b}_v(s)$ are $\leq -(r+\gamma')$. On the other hand, the inclusions

$$\tilde{b}_v(s)(s+r)\delta_g \subseteq V^1 \tilde{\mathcal{R}} \cdot (s+r)\delta_g \subseteq V^1 \tilde{\mathcal{R}} \cdot \delta_g,$$

imply that \tilde{b}_{δ_g} divides $\tilde{b}_v(s)(s+r)$. Since $b_Z(s) = \tilde{b}_{\delta_g}$ (see Remark 3.5), we conclude that $\tilde{\gamma}(Z) \geq r + \gamma'$. This completes the proof of the theorem. \square

We can now show that the equality $F_1 = O_1$ on $\mathcal{H}_Z^r(\mathcal{O}_X)$ implies that Z has rational singularities.

Proof of Corollary 1.7. It follows from Theorem 1.6 that $\tilde{\alpha}(Z) > r$ if and only if $\tilde{\gamma}(Z) > r$. By [BMS06, Theorem 4], this holds if and only if Z has rational singularities (see also Remark 6.4). The last assertion in the corollary now follows from Theorem 1.3. \square

7. APPENDIX: SOME FORMULAS INVOLVING DIFFERENTIAL OPERATORS

In this appendix we collect for ease of reference some easy computations involving differential operators. We work in the ring $\mathbf{C}\langle t, \partial_t, \partial_t^{-1} \rangle$ and put $s = -\partial_t t$.

Lemma 7.1. *For every $m \geq 1$ we have $[\partial_t, t^m] = m t^{m-1}$ and for every $m \in \mathbf{Z}$ we have $[t, \partial_t^m] = -m \partial_t^{m-1}$.*

Proof. The first formula follows from the more general fact that for every derivation D and every regular function f , we have $[D, f] = D(f)$. The second formula is clear if $m = 1$ and the general case follows by induction on $|m|$, using the fact that

$$[t, \partial_t^{m+1}] = (t \partial_t^m - \partial_t^m t) \partial_t + \partial_t^m (t \partial_t - \partial_t t) = [t, \partial_t^m] \partial_t - \partial_t^m,$$

which immediately implies that the formula holds for m if and only if it holds for $m+1$. \square

Lemma 7.2. *For every $m \geq 1$, we have*

$$\partial_t^m t^m = (-1)^m s(s-1) \cdots (s-m+1).$$

Proof. The assertion is clear when $m = 1$ and the general case follows by induction on m , writing

$$\partial_t^{m+1} t^{m+1} = \partial_t t \partial_t^m t^m + \partial_t [\partial_t^m, t] t^m = (\partial_t t + m) \partial_t^m t^m,$$

where the last equality follows using Lemma 7.1. \square

Lemma 7.3. *For every $m \in \mathbf{Z}$ and every $P \in \mathbf{C}[s]$, we have $P(s)\partial_t^m = \partial_t^m P(s+m)$.*

Proof. It is easy to see that it is enough to prove the equality when P is a monomial and then that it is enough to prove it when $P = s$. In this case we have

$$s\partial_t^m - \partial_t^m s = -\partial_t t \partial_t^m + \partial_t^{m+1} t = -\partial_t[t, \partial_t^m] = m\partial_t^m,$$

where the last equality follows from Lemma 7.1. \square

The proof of the next lemma is similar and we leave it for the reader.

Lemma 7.4. *For every $m \in \mathbf{Z}_{\geq 0}$ and every $P \in \mathbf{C}[s]$, we have $P(s)t^m = t^m P(s-m)$.*

REFERENCES

- [BMS06] N. Budur, M. Mustařă, and M. Saito, *Bernstein-Sato polynomials of arbitrary varieties*, Compos. Math. **142** (2006), no. 3, 779–797. [↑2, 4, 5, 7, 8, 29, 30, 32](#)
- [CD23] Q. Chen and B. Dirks, *On V -filtration, Hodge filtration and Fourier transform*, Selecta Math. (N.S.) **29** (2023), no. 4, Paper No. 50, 76 pp. [↑4, 26, 28, 31](#)
- [Dim04] A. Dimca, *Sheaves in topology*, Universitext, Springer-Verlag, Berlin, 2004. [↑27](#)
- [DMST06] A. Dimca, P. Maisonobe, M. Saito, and T. Torrelli, *Multiplier ideals, V -filtrations and transversal sections*, Math. Ann. **336** (2006), no. 4, 901–924. [↑22](#)
- [Dir23] B. Dirks, *Some applications of microlocalization for local complete intersection subvarieties*, preprint arXiv:2310.15277 (2023). [↑5](#)
- [Doh08] D. C. Doherty, *Singularities of generic projection hypersurfaces*, Proc. Amer. Math. Soc. **136** (2008), no. 7, 2407–2415. [↑29](#)
- [EM04] L. Ein and M. Mustařă, *Inversion of adjunction for local complete intersection varieties*, Amer. J. Math. **126** (2004), no. 6, 1355–1365. [↑29](#)
- [HTT08] R. Hotta, K. Takeuchi, and T. Tanisaki, *D -modules, perverse sheaves, and representation theory*, Birkhäuser, Boston, 2008. [↑5, 27](#)
- [Kas83] M. Kashiwara, *Vanishing cycle sheaves and holonomic systems of differential equations*, Algebraic geometry (Tokyo/Kyoto, 1982), Lecture Notes in Math., vol. 1016, Springer, Berlin, 1983, pp. 134–142. [↑2, 5, 7](#)
- [Kol97] J. Kollár, *Singularities of pairs*, Algebraic geometry—Santa Cruz 1995, Proc. Sympos. Pure Math., vol. 62, Amer. Math. Soc., Providence, RI, 1997, pp. 221–287. [↑1](#)
- [Kov99] S. J. Kovács, *Rational, log canonical, Du Bois singularities: on the conjectures of Kollár and Steenbrink*, Compositio Math. **118** (1999), no. 2, 123–133. [↑29](#)
- [Laz04] R. Lazarsfeld, *Positivity in algebraic geometry II*, Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 49, Springer-Verlag, Berlin, 2004. [↑7, 16, 21](#)
- [Lic89] B. Lichtin, *Poles of $|f(z, w)|^{2s}$ and roots of the b -function*, Ark. Mat. **27** (1989), no. 2, 283–304. [↑1](#)
- [Loe84] F. Loeser, *Exposant d’Arnold et sections planes*, C. R. Acad. Sci. Paris Sér. I Math. **298** (1984), no. 19, 485–488. [↑1](#)
- [Mal83] B. Malgrange, *Polynomes de Bernstein-Sato et cohomologie évanescence*, Analysis and topology on singular spaces, II, III (Luminy, 1981), 1983, pp. 243–267. [↑2, 7](#)
- [Mus02] M. Mustařă, *The multiplier ideals of a sum of ideals*, Trans. Amer. Math. Soc. **354** (2002), no. 1, 205–217. [↑16](#)
- [Mus22] M. Mustařă, *Bernstein-Sato polynomials for general ideals vs. principal ideals*, Proc. Amer. Math. Soc. **150** (2022), no. 9, 3655–3662. [↑3, 15](#)
- [MP19] M. Mustařă and M. Popa, *Hodge ideals for \mathbf{Q} -divisors: birational approach*, J. de l’École Polytechnique **6** (2019), 283–328. [↑21, 22](#)
- [MP20] M. Mustařă and M. Popa, *Hodge ideals for \mathbf{Q} -divisors, V -filtration, and minimal exponent*, Forum Math. Sigma **8** (2020), Paper No. e19, 41. [↑2, 3, 9, 20, 21, 23, 24](#)
- [MP22] ———, *Hodge filtration on local cohomology, Du Bois complex and local cohomological dimension*, Forum Math. Pi **10** (2022), Paper No. e22, 58 pp. [↑4, 5, 28, 29](#)
- [Sai88] M. Saito, *Modules de Hodge polarisables*, Publ. Res. Inst. Math. Sci. **24** (1988), no. 6, 849–995. [↑5, 12, 26](#)
- [Sai90] ———, *Mixed Hodge modules*, Publ. Res. Inst. Math. Sci. **26** (1990), no. 2, 221–333. [↑1](#)

- [Sai93] ———, *On b -function, spectrum and rational singularity*, Math. Ann. **295** (1993), no. 1, 51–74. [↑9](#)
- [Sai94] ———, *On microlocal b -function*, Bull. Soc. Math. France **122** (1994), no. 2, 163–184. [↑1](#), [2](#), [8](#), [9](#), [20](#), [26](#)
- [Sai16] ———, *Hodge ideals and microlocal V -filtration*, preprint arXiv:1612.08667 (2016). [↑2](#), [9](#)
- [Sch19] C. Schnell, *An overview of Morihiko Saito’s theory of mixed Hodge modules*, Representation theory, automorphic forms & complex geometry, Int. Press, Somerville, MA, 2019, pp. 27–80. [↑](#)
- [Sch07] K. Schwede, *A simple characterization of Du Bois singularities*, Compos. Math. **143** (2007), no. 4, 813–828. [↑29](#)
- [Ste85] J. H. M. Steenbrink, *Semicontinuity of the singularity spectrum*, Invent. Math. **79** (1985), no. 3, 557–565. [↑1](#)
- [Var82] A. N. Varchenko, *The complex singularity index does not change along the stratum $\mu = \text{const}$* , Funktsional. Anal. i Prilozhen. **16** (1982), no. 1, 1–12, 96. [↑1](#)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 530 CHURCH STREET, ANN ARBOR, MI 48109, USA

Email address: `qyc@umich.edu`

DEPARTMENT OF MATHEMATICS, STONY BROOK UNIVERSITY, STONY BROOK, NY 11794-3651, USA

Email address: `bradley.dirks@stonybrook.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 530 CHURCH STREET, ANN ARBOR, MI 48109, USA

Email address: `mmustata@umich.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, 40 ST. GEORGE ST., TORONTO, ONTARIO, CANADA, M5S 2E4

Email address: `seolano@math.toronto.edu`