

## Solutions to Practice Final 2

1. Using induction prove that

$$1^2 + 3^2 + \dots + (2n + 1)^2 = \frac{(n + 1)(2n + 1)(2n + 3)}{3}$$

### Solution

First we verify the base of induction. When  $n = 0$  LHS =  $1^2 = 1$  and RHS =  $\frac{1 \cdot 1 \cdot 3}{3} = 1$ .

Induction step. Assume the formula is true for  $n \geq 0$  and we need to verify it for  $n + 1$ . Then we have

$$\begin{aligned} 1^2 + 3^2 + \dots + (2n + 1)^2 + (2n + 3)^2 &= \frac{(n + 1)(2n + 1)(2n + 3)}{3} + (2n + 3)^2 = \\ &= \frac{(n + 1)(2n + 1)(2n + 3) + 3(2n + 3)^2}{3} = \frac{(2n + 3)(2n^2 + 3n + 1 + 3(2n + 3))}{3} \\ &= \frac{(2n + 3)(2n^2 + 9n + 10)}{3} = \frac{(2n + 3)(2n + 5)(n + 2)}{3} \end{aligned}$$

This completes the induction step and proves the formula for all  $n \geq 0$ .

2. Let  $a, b, c$  be natural numbers.

- (a) Show that the equation  $ax + by = c$  has a solution if and only if  $(a, b) | c$ .  
(b) Find all integer solutions of  $6x + 15y = 9$ .

### Solution

- (a) Suppose  $ax + by = c$  for some integer  $x$  and  $y$ . If  $d | a$  and  $d | b$  then obviously,  $d | ax + by = c$ . In particular, if  $(a, b) | c$ .

Conversely, suppose  $(a, b) | c$  so that  $c = d \cdot (a, b)$ . Then  $ax + by = (a, b)$  has an integer solution by a result from class. Multiplying both sides by  $d$  we get  $a(xd) + b(yd) = (a, b) \cdot d = c$ .

- (b) First, divide both sides by 3. we get  $2x + 5y = 3$ . We have  $(2, 5) = 1$  and we can find integer solution of  $2x + 5y = 1$  using either Euclidean algorithm or just by trying a few small numbers we get

$2 \cdot (-2) + 5 \cdot 1 = 1$ . Multiplying by 3 we get  $2 \cdot (-6) + 5 \cdot (3) = 3$  so  $x_0 = -6, y_0 = 3$  is a solution of  $2x + 5y = 3$ .

It's easy to see that  $x = -6 - 5k, y = 3 + 2k$  is a solution of  $2x + 5y = 3$  for any  $k$ . We claim that any integer solution of  $2x + 5y = 3$  has this form.

Suppose  $2x + 5y = 3$ . we also have  $2 \cdot (-6) + 5 \cdot (3) = 3$ . Subtracting these equations we get  $2(-6 - x) + 5(3 - y) = 0$  or  $2(-6 - x) = 5(y - 3)$ . This implies that  $2 | (y - 3)$  so that  $y - 3 = 2k$  or  $y = 3 + 2k$ . This gives  $2(-6 - x) = 5(y - 3) = 6k, -6 - x = 3k, x = -6 - 3k$ .

Thus the general solution is  $x = -6 - 5k, y = 3 + 2k$  where  $k$  is any integer.

3. Find the last digit of the sum

$$2(1 + 3 + 3^2 + 3^3 + \dots + 3^{309})$$

### Solution

First, we compute

$$2(1 + 3 + 3^2 + 3^3 + \dots + 3^{309}) = 2 \cdot \frac{3^{310} - 1}{3 - 1} = 3^{310} - 1.$$

We have  $\phi(10) = \phi(2 \cdot 5) = 1 \cdot 4 = 4$ . By Euler's theorem this implies that  $3^4 \equiv 1 \pmod{10}$ . Of course, this can also be seen directly as  $3^4 = 81$ .

Therefore  $3^{4k} \equiv 1 \pmod{10}$ . We have  $310 = 308 + 2$  and  $4|308$ . Therefore  $3^{310} \equiv 3^2 \pmod{10}$ . This means that the last digit of  $3^{310}$  is 9 and hence the last digit of  $3^{310} - 1$  is 8.

4. Let  $S$  be infinite and  $A \subset S$  be finite. Prove that  $|S| = |S \setminus A|$ .

### Solution

Let  $A = \{s_1, \dots, s_n\}$ . Since  $S$  is infinite the set  $S \setminus A$  is non empty. Pick any  $s_{n+1} \in S \setminus A = S \setminus \{s_1, \dots, s_n\}$ . Next, since  $S \setminus \{s_1, \dots, s_{n+1}\} \neq \emptyset$  we can choose  $s_{n+2} \in S \setminus \{s_1, \dots, s_{n+1}\}$ . Proceeding by induction we can construct  $s_{m+1} \in S \setminus \{s_1, \dots, s_m\}$  for any  $m \geq n$ .

Now define  $f: S \rightarrow S \setminus A$  by the formula  $f(s_i) = s_{i+n}$  for any  $i$  and  $f(x) = x$  if  $x \in S \setminus \{s_1, s_2, \dots\}$ . By construction,  $f$  is 1-1 and onto.

5. Let  $S = [0, 1]$  and  $T = [0, 2)$ . Let  $f: S \rightarrow T$  be given by  $f(x) = x$  and  $g: T \rightarrow S$  be given by  $g(x) = x/2$ .

(a) Find  $S_S, S_T, S_\infty$ ;

(b) give an explicit formula for a 1-1 and onto map  $h: S \rightarrow T$  coming from  $f$  and  $g$  using the proof of the Schroeder-Berstein theorem.

### Solution

(a) Note that  $1 \notin g(T)$  and therefore  $1 \in S_S$ . Next, we see that  $1/2 \in S_S$  also. Indeed,  $1/2 = g(1)$  and  $1 = f(1)$ . So 1 is the last ancestor of  $1/2$  and hence  $1/2 \in S_S$ . proceeding by induction we see that  $\frac{1}{2^n} \in S_S$  for any  $n \geq 0$ .

Next observe that  $(1/2, 1) \subset S_T$ . Indeed, if  $1/2 < x < 1$  then  $x = g(2x)$  and  $1 < 2x < 2$  so that  $2x \notin f(S)$ .

Proceeding by induction we claim that  $(\frac{1}{2^{n+1}}, \frac{1}{2^n}) \in S_T$  for any  $n \geq 0$ . We just verified the base of induction.

Induction step. Suppose we know the statement of  $n \geq 0$  and we need to prove it for  $n + 1$ . Let  $\frac{1}{2^{n+2}} < x < \frac{1}{2^{n+1}}$  then  $x = g(2x)$  and  $\frac{1}{2^{n+1}} < 2x < \frac{1}{2^n}$ . Also,  $2x = f(2x)$ . By induction assumption,  $2x \in S_T$  and the last ancestor of  $x$  is the last ancestor of  $2x$  so  $x \in S_T$  also.

This concludes the induction step.

It's obvious that  $0 \in S_\infty$ . Therefore  $S_\infty = \{0\}$ ,  $S_S = \{1, \frac{1}{2}, \frac{1}{4}, \dots, \frac{1}{2^n}, \dots\}$  and  $S_T = \{x \in [0, 1] \text{ such that } x \neq 0, 1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots\}$ .

- (b) By the proof of the Schroeder-Bernstein Theorem the following map  $h: S \rightarrow T$  is 1-1 and onto.

$$h(x) = \begin{cases} x & \text{if } x = 0, 1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots \\ 2x & \text{if } x \neq 0, 1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \dots \end{cases}$$

6. Let  $n = 2p$  where  $p$  is an odd prime. Find the remainder when  $\phi(n)!$  is divided by  $n$ . Here  $\phi(n)$  is the Euler function of  $n$ .

### Solution

We have  $\phi(n) = \phi(2p) = (2-1)(p-1) = p-1$ . By Wilson's theorem  $\phi(n)! = (p-1)! \equiv -1 \pmod{p} \equiv p-1 \pmod{p}$ . This means that  $p \mid (p-1)! - (p-1)$ . Since  $p$  is odd  $p-1$  is even and therefore  $2 \mid (p-1)! - (p-1)$  also. Since  $(2, p) = 1$  this implies that  $2p \mid (p-1)! - (p-1)$  or, equivalently  $(p-1)! \equiv p-1 \pmod{2p}$ .

**Answer:**  $p-1$ .

7. Prove that  $q_1\sqrt{3} + q_2\sqrt{5} \neq q'_1\sqrt{3} + q'_2\sqrt{5}$  for any rational  $q_1, q_2, q'_1, q'_2$  unless  $q_1 = q'_1, q_2 = q'_2$ .

### Solution

Suppose  $q_1\sqrt{3} + q_2\sqrt{5} = q'_1\sqrt{3} + q'_2\sqrt{5}$ . Then  $(q_1 - q'_1)\sqrt{3} + (q_2 - q'_2)\sqrt{5} = 0$ . Let  $a = q_1 - q'_1, b = q_2 - q'_2$  are rational and  $a\sqrt{3} + b\sqrt{5} = 0$ . We want to show that  $a = b = 0$ . If  $a \neq 0$  this gives  $\sqrt{\frac{3}{5}} = -\frac{b}{a}$  which is rational. This is a contradiction since  $\sqrt{\frac{3}{5}}$  is irrational. Hence  $a = 0$ . Since  $a\sqrt{3} + b\sqrt{5} = 0$  this implies  $b\sqrt{5} = 0, b = 0$ .

8. Let  $a$  be a root of  $x^5 - 6x^3 + 2x^2 + 5x - 1 = 0$ . Construct a polynomial with integer coefficients which has  $a^2$  as a root.

*Hint:* separate even and odd powers.

### Solution

We can rewrite the equation as  $x^5 - 6x^3 + 5x = 1 - 2x^2$ ,  $x(x^4 - 6x^2 + 5) = 1 - 2x^2$ . Squaring both sides we get  $x^2(x^4 - 6x^2 + 5)^2 = (1 - 2x^2)^2$ . Clearly,  $y = x^2$  satisfies  $y(y^2 - 6y + 5)^2 = (1 - 2y)^2$ .

9. Find all complex roots of  $x^6 + 7x^3 - 8 = 0$ .

*Reminder:* Real numbers are also complex numbers.

### Solution

Let  $z = x^3$ . Then  $z$  satisfies  $z^2 + 7z - 8 = 0$ . Solving this quadratic equation we get  $z = 1, z = -8$ . Thus we need to solve  $x^3 = 1$  and  $x^3 = -8$ . Solving  $x^3 = 1$  gives  $x = 1, x = \cos(2\pi/3) + i \sin(2\pi/3) = \frac{-1+i\sqrt{3}}{2}, x = \cos(4\pi/3) + i \sin(4\pi/3) = \frac{-1-i\sqrt{3}}{2}$

Next we write  $-8$  as  $2^3(\cos \pi + i \sin \pi)$ . Thus solving  $x^3 = -8$  we get  $x = 2(\cos(\pi/3) + i \sin(\pi/3)) = 1 + i\sqrt{3}, x = 2(\cos(\pi/3 + 2\pi/3) + i \sin(\pi/3 + 2\pi/3)) = 2(\cos \pi + i \sin \pi) = -2, x = 2(\cos(\pi/3 + 4\pi/3) + i \sin(\pi/3 + 4\pi/3)) = 2(\cos(5\pi/3) + i \sin(5\pi/3)) = 1 - i\sqrt{3}$

10. Represent  $\sin(5\theta)$  as a polynomial in  $\sin(\theta)$ .

### Solution

We have  $\cos(5\theta) + i \sin(5\theta) = (\cos \theta + i \sin \theta)^5 = (\cos \theta + i \sin \theta)^2(\cos \theta + i \sin \theta)^3$ . We compute separately  $(\cos \theta + i \sin \theta)^2 = (\cos^2 \theta - \sin^2 \theta + 2i \sin \theta \cos \theta)$  and  $(\cos \theta + i \sin \theta)^3 = (\cos \theta + i \sin \theta)^2(\cos \theta + i \sin \theta) = (\cos^2 \theta - \sin^2 \theta + 2i \sin \theta \cos \theta)(\cos \theta + i \sin \theta) = (\cos^2 \theta - \sin^2 \theta) \cos \theta - 2 \sin^2 \theta \cos \theta + i(\cos^2 \theta - \sin^2 \theta) \sin \theta + 2i \sin \theta \cos^2 \theta = \cos^3 \theta - 3 \sin^2 \theta \cos \theta + i(3 \sin \theta \cos^2 \theta - \sin^3 \theta)$ .

Combining these together we get  $\cos(5\theta) + i \sin(5\theta) = (\cos \theta + i \sin \theta)^5 = (\cos \theta + i \sin \theta)^2(\cos \theta + i \sin \theta)^3 = (\cos^2 \theta - \sin^2 \theta + 2i \sin \theta \cos \theta)(\cos^3 \theta - 3 \sin^2 \theta \cos \theta + i(3 \sin \theta \cos^2 \theta - \sin^3 \theta)) = (\cos^2 \theta - \sin^2 \theta)(\cos^3 \theta - 3 \sin^2 \theta \cos \theta) - 2 \sin \theta \cos \theta(3 \sin \theta \cos^2 \theta - \sin^3 \theta) + i(\cos^2 \theta - \sin^2 \theta)(3 \sin \theta \cos^2 \theta - \sin^3 \theta) + 2i \sin \theta \cos \theta(\cos^3 \theta - 3 \sin^2 \theta \cos \theta)$ .

Therefore,  $\sin(5\theta) = (\cos^2 \theta - \sin^2 \theta)(3 \sin \theta \cos^2 \theta - \sin^3 \theta) + 2 \sin \theta \cos \theta(\cos^3 \theta - 3 \sin^2 \theta \cos \theta) = (1 - 2 \sin^2 \theta)(3 \sin \theta(1 - \sin^2 \theta) - \sin^3 \theta) + 2 \sin \theta \cos^4 \theta - 6 \sin^3 \theta \cos^2 \theta = (1 - 2 \sin^2 \theta)(3 \sin \theta(1 - \sin^2 \theta) - \sin^3 \theta) + 2 \sin \theta(1 - \sin^2 \theta)^2 - 6 \sin^3 \theta(1 - \sin^2 \theta)$ .

11. Is  $\frac{\sqrt[6]{5} - \sqrt{5}}{1 + 2\sqrt{7}}$  constructible? Justify your answer.

### Solution

$\frac{\sqrt[6]{5}-\sqrt{5}}{1+2\sqrt{7}}$  is not constructible. We argue by contradiction. Assume  $\frac{\sqrt[6]{5}-\sqrt{5}}{1+2\sqrt{7}}$  is constructible. Since  $\sqrt{5}$  and  $\sqrt{7}$  are constructible this implies that  $\sqrt[6]{5}$  is constructible and hence  $(\sqrt[6]{5})^2 = \sqrt[3]{5}$  is also constructible.  $\sqrt[3]{5}$  is a root of  $x^3 - 5 = 0$  which is a cubic equation with integer coefficients. By a theorem from class if it has a constructible root it must have a rational root as well. Let  $\frac{m}{n}$  be a rational root where  $(m, n) = 1$ . Then  $m|5$  and  $n|1$  which means that  $\frac{m}{n} = \pm 1, \pm 5$ . Plugging these numbers into  $x^3 - 5 = 0$  we see that none of them are roots.


This is a contradiction and therefore  $\frac{\sqrt[6]{5}-\sqrt{5}}{1+2\sqrt{7}}$  is not constructible.

12. For each of the following answer "true" or "false". Justify your answer.

- a) If  $\frac{x}{y}$  is constructible then both  $x$  and  $y$  are constructible.
- b) If  $x$  is constructible then  $\frac{1}{x}$  is constructible.
- c) There is an angle  $\theta$  such that  $\cos \theta$  is constructible but  $\sin \theta$  is not constructible.
- d)  $\sqrt[3]{\frac{10}{27}}$  is constructible.

### Solution

- a) **False.** For example, take  $x = y = \pi$ . Then  $X$  and  $y$  are not constructible but  $x/y = 1$  is constructible.
- b) **True.** See figure below. Draw segments of lengths 1 and  $x$  on one side of an angle and a segment of length 1 on the other side. Connect  $x$  and 1 on opposite sides by a line a draw a parallel line through 1 on the same side as  $x$ . It intersect the second side of the angle at distance  $y$ . Then from similar triangles we get  $\frac{x}{1} = \frac{1}{y}$  or  $y = \frac{1}{x}$



- c) **False.** If  $\cos \theta$  is constructible then so is  $1 - \cos^2 \theta$ . Hence  $\sin \theta = \pm \sqrt{1 - \cos^2 \theta}$  is also constructible since a square root of a constructible number is constructible.
- d) **False.** We argue by contradiction. Suppose  $x = \sqrt[3]{\frac{10}{27}}$  is constructible. It satisfies the equation  $27x^3 - 10 = (3x)^3 - 10 = 0$ . If  $x$  is constructible then so is  $y = 3x$  which satisfies the equation  $y^3 - 10 = 0$ . This is a cubic equation with integer coefficients. If it has a constructible root it must also have a rational one. We can write that rational root as  $\frac{a}{b}$  where  $(a, b) = 1$ . Then  $a|10$  and  $b|1$  which means that  $y = \frac{a}{b} = \pm 1 \pm 2 \pm 5$  or  $\pm 10$ . By plugging these numbers into  $y^3 - 10 = 0$  we see that none of them are roots. This is a contradiction and therefore  $\sqrt[3]{\frac{10}{27}}$  is not constructible.

13. Prove that the equation

$$(1 + x^{19})^3 + (1 + x^{19})^2 - 3 = 0$$

has no constructible solutions.

### Solution

Suppose  $x$  is a constructible root. Then  $y = x^{19} + 1$  is also constructible and it satisfies  $y^3 + y^2 - 3 = 0$ . This is a cubic equation with integer coefficients. If it has

a constructible root it must also have a rational one. We can write that rational root as  $\frac{a}{b}$  where  $(a, b) = 1$ . Then  $a|1$  and  $b|1$  which means that  $y = \frac{a}{b} = \pm 1$ . But neither  $y = 1$  nor  $y = -1$  solve  $y^3 + y^2 - 1 = 0$ . This is a contradiction which means that  $(1 + x^{19})^3 + (1 + x^{19})^2 - 3 = 0$  has no constructible solutions.