



Strand Doubling and Reversal.

$$\begin{matrix} \omega & a & S \\ S & \alpha & \theta \\ a & \phi & \Xi \end{matrix} \xrightarrow[\mu=T_a^{-1}]{\mu=\Delta_{bc}^a} \begin{pmatrix} \omega & b & c & S \\ (\sigma_a - \alpha T_a - \nu T_c)/\mu & (T_b - 1)T_c\nu/\mu & (T_b - 1)T_c\theta/\mu & \\ c & (T_c - 1)\nu/\mu & (\alpha - \sigma_a T_a - \nu T_c)/\mu & (T_c - 1)\theta/\mu \\ S & \phi & \phi & \Xi \end{pmatrix}$$

$$dS^a \downarrow T_a \rightarrow T_a^{-1}$$


$$\begin{pmatrix} \alpha\omega/\sigma_a & a & S \\ a & 1/\alpha & \theta/\alpha \\ S & -\phi/\alpha & (\alpha\Xi - \phi\theta)/\alpha \end{pmatrix}$$

Where σ assigns to every $a \in S$ a Laurent monomial σ_a in $\{t_b\}_{b \in S}$ subject to $\sigma(a \nearrow_b, b \nearrow_a) = (a \rightarrow 1, b \rightarrow t_a^{\pm 1})$, $\sigma(T_1 \sqcup T_2) = \sigma(T_1) \sqcup \sigma(T_2)$, and $\sigma//m_c^{ab} = (\sigma \setminus \{a, b\}) \cup (c \rightarrow \sigma_a \sigma_b)_{t_a, t_b \rightarrow t_c}$.

Algebraic Knot Theory




Abstract. This will be a very “light” talk: I will explain why about 13 years ago, in order to have a say on some problems in knot theory, I’ve set out to find tangle invariants with some nice compositional properties. In other talks in Sydney (ωεβ/talks) I have explained / will explain how such invariants were found - though they are yet to be explored and utilized.

(v-)Tangles.



Genus. Every knot is the boundary of an orientable “Seifert Surface” (ωεβ/SS), and the least of their genera is the “genus” of the knot.

Claim. The knots of genus ≤ 2 are precisely the images of 4-component tangles via




A Bit about Ribbon Knots. A “ribbon knot” is a knot that can be presented as the boundary of a disk that has “ribbon singularities”, but no “clasp singularities”. A “slice knot” is a knot in $S^3 = \partial B^4$ which is the boundary of a non-singular disk in B^4 . Every ribbon knot is clearly slice, yet,

Conjecture. Some slice knots are not ribbon.

Fox-Milnor. The Alexander polynomial of a ribbon knot is always of the form $A(t) = f(t)f(1/t)$. (also for slice)

Theorem. K is ribbon iff it is κT for a tangle T for which τT is the untangle U .



The Gold Standard is set by the “ Γ -calculus” Alexander formulas [BNS, BN]. An S -component tangle T has $\Gamma(T) \in R_S \times M_{S \times S}(R_S) = \left\{ \begin{matrix} \omega & S \\ S & A \end{matrix} \right\}$ with $R_S := \mathbb{Z}\langle\langle T_a : a \in S \rangle\rangle$:

$$(a \nearrow_b, b \nearrow_a) \rightarrow \begin{matrix} 1 & a & b \\ a & 1 & -T_a^{-1} \\ b & 0 & T_a^{-1} \end{matrix} \quad T_1 \sqcup T_2 \rightarrow \begin{matrix} \omega_1 \omega_2 & S_1 & S_2 \\ S_1 & A_1 & 0 \\ S_2 & 0 & A_2 \end{matrix}$$

$$\begin{matrix} \omega & a & b & S \\ a & \alpha & \beta & \theta \\ b & \gamma & \delta & \epsilon \\ S & \phi & \psi & \Xi \end{matrix} \xrightarrow[m_c^{ab}]{T_a, T_b \rightarrow T_c} \begin{pmatrix} (1-\beta)\omega & c & S \\ c & \gamma + \frac{\alpha\delta}{1-\beta} & \epsilon + \frac{\delta\theta}{1-\beta} \\ S & \phi + \frac{\alpha\psi}{1-\beta} & \Xi + \frac{\psi\theta}{1-\beta} \end{pmatrix}$$

For long knots, ω is Alexander, and that’s the fastest Alexander algorithm I know! Dunfield: 1000-crossing fast.



Vo’s Thesis [Vo]. A proof of the Fox-Milnor theorem for ribbon knots using this technology (and more).

Implementation key idea:

ωεβ/AlexDemo

```

(ω, A = (αab)) ↔
(ω, λ = Σ αab t_a h_b)

F /: F[ω_1, λ_1] F[ω_2, λ_2] := F[ω_1 * ω_2, λ_1 * λ_2]
m_h_a -> [Γ[ω_1, λ_1]] := Module[{a, b, γ, δ, θ, e, φ, ψ, ξ, μ},
  {
    {α β θ} = {θ_a, h_a, λ} {α_b, h_b, λ} {α_c, h_c, λ} /. {t | h}_a | b -> 0;
    Γ[(μ-1-β)ω, (t_1, 1)·{γ+αδ/μ, ε+δθ/μ}·{h_c, 1}]
    /. {T_a → T_c, T_b → T_c} // FCollect];
M = Prepend[M, {t_1, t_2} // Transpose;
M = Prepend[M, Prepend[{h_1, h_2} / θ, ω], ω];
M // MatrixForm];
Rm_{a,b} := Γ[1, (t_1, t_2)·{0 1-T_a}·{h_a, h_b}];
Rm_{a,b} := Rm_{a,b} / T_a → 1/T_a;



```

Meta-Associativity

$$\xi = \Gamma[\omega, \{t_1, t_2, t_3, t_4\}] \cdot \begin{pmatrix} \alpha_{11} & \alpha_{12} & \alpha_{13} & \theta_1 \\ \alpha_{21} & \alpha_{22} & \alpha_{23} & \theta_2 \\ \alpha_{31} & \alpha_{32} & \alpha_{33} & \theta_3 \\ \phi_1 & \phi_2 & \phi_3 & \Xi \end{pmatrix} \cdot \{h_1, h_2, h_3, h_4\};$$

$$(\xi // m_{12 \rightarrow 1} // m_{13 \rightarrow 1}) = (\xi // m_{23 \rightarrow 2} // m_{12 \rightarrow 1})$$

True R3 ... divide and conquer!



Fact. Γ is better viewed as an invariant of a certain class of 2D knotted objects in \mathbb{R}^4 [BND, BN].

Fact. Γ is the “0-loop” part of an invariant that generalizes to “ n -loops” (1D tangles only, see further talks and future publications with van der Veen).

Speculation. Stepping stones to categorification?

Ask me about geography vs. identity!



References.

[BN] D. Bar-Natan, *Balloons and Hoops and their Universal Finite Type Invariant, BF Theory, and an Ultimate Alexander Invariant*, ωεβ/KBH, arXiv:1308.1721.
[BND] D. Bar-Natan and Z. Dancso, *Finite Type Invariants of W-Knotted Objects I: w-Knots and the Alexander Polynomial*, Alg. and Geom. Top. **16-2** (2016) 1063–1133, arXiv:1405.1956, ωεβ/WK01.
[BNS] D. Bar-Natan and S. Selmani, *Meta-Monoids, Meta-Bicrossed Products, and the Alexander Polynomial*, J. of Knot Theory and its Ramifications **22-10** (2013), arXiv:1302.5689.
[GST] R. E. Gompf, M. Scharlemann, and A. Thompson, *Fibered Knots and Potential Counterexamples to the Property 2R and Slice-Ribbon Conjectures*, Geom. and Top. **14** (2010) 2305–2347, arXiv:1103.1601.
[Vo] H. Vo, *Alexander Invariants of Tangles via Expansions*, University of Toronto Ph.D. thesis, ωεβ/Vo.

“God created the knots, all else in topology is the work of mortals.”
Leopold Kronecker (modified) www.katlas.org The Knot Atlas