





Knots in Four Dimensions and the Simplest Open Problem About Them

Abstract. I will describe a few 2-dimensional knots in 4 dimensional space in detail, then tell you how to make many more, then tell you that I don't really understand my way of making them, yet I can tell at least some of them apart in a colourful way.

u-Knots.




Satoh




Dalvit
 ω/Dal

The Generators



The Double Inflation Procedure δ .


w-Knots.



"God created the knots, all else in topology is the work of mortals."

Leopold Kronecker (modified)


www.katlas.org



Satoh's Conjecture. (ω/Sat) The "kernel" of the double inflation map δ , mapping w-knot diagrams in the plane to knotted 2D tubes and spheres in 4D, is precisely the moves R2-3, VR1-3, M, CP and OC listed above. In other words, two w-knot diagrams represent via δ the same 2D knot in 4D iff they differ by a sequence of the said moves.

First Isomorphism Thm: $\delta: G \rightarrow H \Rightarrow \text{im } \delta \cong G / \ker(\delta)$
 δ is a map from algebra to topology. So a thing in "hard" topology ($\text{im } \delta$) is the same as a thing in "easy" algebra ($w\mathcal{K}$).

Reidemeister's Theorem.




Proof by a genericity / "shaking" argument



Kurt Reidemeister


3-Colourings. Colour the arcs of a broken arc diagram in RGB so that every crossing is either mono-chromatic or tri-chromatic; $\lambda(K) := |\{3\text{-colourings}\}|$.




Example. $\lambda(\bigcirc) = 3$ while $\lambda(\bigodot) = 9$; so $\bigcirc \neq \bigodot$.

Exercise. Show that the set of colourings of K is a vector space over \mathbb{F}_3 hence $\lambda(K)$ is always a power of 3.


Extend λ to $w\mathcal{K}$ by declaring that arcs "don't see" v-xings, and that caps are always "kosher". Then $\lambda(\bullet\text{---}\bullet) = 3 \neq 9 = \lambda(\text{CS 2-knot})$, so assuming Conjecture, the CS 2-knot is indeed knotted.



The Full 2-Knot Story



Roseman Moves.



Expansions. Given a "ring" K and an ideal $I \subset K$, set $A := I^0/I^1 \oplus I^1/I^2 \oplus I^2/I^3 \oplus \dots$.

A homomorphic expansion is a multiplicative $Z: K \rightarrow A$ such that if $\gamma \in I^m$, then $Z(\gamma) = (0, 0, \dots, 0, \gamma/I^{m+1}, *, *, \dots)$.

Example. Let $K = C^\infty(\mathbb{R}^n)$ be smooth functions on \mathbb{R}^n , and $I := \{f \in K: f(0) = 0\}$. Then $I^m = \{f: f \text{ vanishes as } |x|^m\}$ and I^m/I^{m+1} is {homogeneous polynomials of degree m } and A is the set of power series. So Z is "a Taylor expansion".

Hence Taylor expansions are vastly general; even knots can be Taylor expanded!