

# Meta-Groups, Meta-Bicrossed-Products, and the Alexander Polynomial, 1

Dror Bar-Natan at Sheffield, February 2013.

<http://www.math.toronto.edu/~drorbn/Talks/Sheffield-130206/>




**Abstract.** I will define “meta-groups” and explain how one specific meta-group, which in itself is a “meta-bicrossed-product”, gives rise to an “ultimate Alexander invariant” of tangles, that contains the Alexander polynomial (multivariable, if you wish), has extremely good composition properties, is evaluated in a topologically meaningful way, and is least-wasteful in a computational sense. If you believe in categorification, that’s a wonderful playground.

This work is closely related to work by Le Dimet (Comment. Math. Helv. **67** (1992) 306–315), Kirk, Livingston and Wang (arXiv:math/9806035) and Cimasoni and Turaev (arXiv:math.GT/0406269).

## Alexander Issues.


- Quick to compute, but computation departs from topology.
- Extends to tangles, but at an exponential cost.
- Hard to categorify.

**Idea.** Given a group  $G$  and two “YB” pairs  $R^\pm = (g_o^\pm, g_u^\pm) \in G^2$ , map them to xings and “multiply along”, so that





**This Fails!** R2 implies that  $g_o^\pm g_o^\mp = e = g_u^\pm g_u^\mp$  and then R3 implies that  $g_o^+$  and  $g_u^+$  commute, so the result is a simple counting invariant.

**A Group Computer.** Given  $G$ , can store group elements and perform operations on them:




Also has  $S_x$  for inversion,  $e_x$  for unit insertion,  $d_x$  for register deletion,  $\Delta_{xy}^z$  for element cloning,  $\rho_y^x$  for renamings, and  $(D_1, D_2) \mapsto D_1 \cup D_2$  for merging, and many obvious composition axioms relating those.

$$P = \{x : g_1, y : g_2\} \Rightarrow P = \{d_y P\} \cup \{d_x P\}$$



**A Standard Alexander Formula.** Label the arcs 1 through  $(n+1) = 1$ , make an  $n \times n$  matrix as below, delete one row and one column, and compute the determinant:



$$\begin{pmatrix} 1 & 0 & 0 & 0 & 0 & x-1 & 0 & -x \\ -1 & x & 0 & 0 & 0 & 0 & 1-x & 0 \\ 0 & -1 & x & 0 & 1-x & 0 & 0 & 0 \\ x-1 & 0 & -x & 1 & 0 & 0 & 0 & 0 \\ 0 & 1-x & 0 & -1 & x & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & -x & 1 & 0 & x-1 \\ 0 & 0 & 1-x & 0 & 0 & -1 & x & 0 \\ 0 & 0 & 0 & x-1 & 0 & 0 & -x & 1 \end{pmatrix} \quad [ [1 ; ; 7, 1 ; ; 7] ] // \text{Det}$$

$$-1 + 4x - 8x^2 + 11x^3 - 8x^4 + 4x^5 - x^6$$

**Claim.** From a meta-group  $G$  and YB elements  $R^\pm \in G_2$  we can construct a knot/tangle invariant.

