



From Stonehenge to Witten – Some Further Details

Oporto Meeting on Geometry, Topology and Physics, July 2004


Dror Bar-Natan, University of Toronto



Witten

We the generating function of all stellar coincidences:

$$Z(K) := \lim_{N \rightarrow \infty} \sum_{\substack{D \\ 3\text{-valent}}} \frac{1}{2^c c! \binom{N}{e}} \langle D, K \rangle_{\mathbb{R}} D \cdot \left(\begin{array}{l} \text{framing-} \\ \text{dependent} \\ \text{counter-term} \end{array} \right) \in \mathcal{A}(\odot)$$




Theorem. Modulo Relations, $Z(K)$ is a knot invariant!

Dylan Thurston




$N := \# \text{ of stars}$ $\mathcal{A}(\odot) := \text{Span} \left\langle \left(\begin{array}{c} \square \\ \square \\ \square \end{array} \right) \right\rangle / \text{oriented vertices}$
 $c := \# \text{ of chopsticks}$ AS: $\begin{array}{c} \diagup \\ \diagdown \end{array} + \begin{array}{c} \diagdown \\ \diagup \end{array} = 0$
 $e := \# \text{ of edges of } D$ & more relations

When deforming, catastrophes occur when:



$$\int_{\mathfrak{g}\text{-connections}} \mathcal{D}A \text{hol}_K(A) \exp \left[\frac{ik}{4\pi} \int_{\mathbb{R}^3} \text{tr} \left(A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) \right] \rightarrow \sum_{D: \text{Feynman diagram}} W_{\mathfrak{g}}(D) \int \mathcal{E}(D) \rightarrow \sum_{D: \text{Feynman diagram}} D \int \mathcal{E}(D)$$



Definition. V is finite type (Vassiliev, Goussarov) if it vanishes on sufficiently large alternations as on the right

Theorem. All knot polynomials (Conway, Jones, etc.) are of finite type.

Conjecture. (Taylor's theorem) Finite type invariants separate knots.

Theorem. $Z(K)$ is a universal finite type invariant! (sketch: to dance in many parties, you need many feet).



Goussarov



Vassiliev

Related to Lie algebras

$$\begin{array}{c} x \quad y \\ \diagdown \quad \diagup \\ [x, y] \end{array} = xy - yx$$

$$\begin{array}{c} x \quad y \quad z \\ \diagdown \quad \diagup \quad \diagdown \\ [[x, y], z] = [x, [y, z]] - [y, [x, z]] \end{array}$$



Sophus Lie

More precisely, let $\mathfrak{g} = \langle X_a \rangle$ be a Lie algebra with an orthonormal basis, and let $R = \langle v_\alpha \rangle$ be a representation. Set

$$f_{abc} := \langle [a, b], c \rangle \quad X_a v_\beta = \sum_{\gamma} r_{a\gamma}^{\beta} v_\gamma$$

and then

$$W_{\mathfrak{g}, R} : \begin{array}{c} \gamma \\ \diagdown \quad \diagup \\ \alpha \end{array} \begin{array}{c} a \\ \diagdown \quad \diagup \\ \beta \end{array} \begin{array}{c} b \\ \diagdown \quad \diagup \\ \alpha \end{array} \begin{array}{c} c \\ \diagdown \quad \diagup \\ \alpha \end{array} \rightarrow \sum_{abc\alpha\beta\gamma} f_{abc} r_{a\gamma}^{\beta} r_{b\alpha}^{\gamma} r_{c\beta}^{\alpha}$$

Planar algebra and the Yang-Baxter equation



Yang




Baxter

$W_{\mathfrak{g}, R} \circ Z$ is often interesting:


$\mathfrak{g} = \mathfrak{sl}(2)$ \rightarrow The Jones polynomial

$\mathfrak{g} = \mathfrak{sl}(N)$ \rightarrow The HOMFLYPT polynomial

$\mathfrak{g} = \mathfrak{so}(N)$ \rightarrow The Kauffman polynomial



Parenthesized tangles, the pentagon and hexagon



Reshetikhin



Turaev

Kauffman's bracket and the Jones polynomial

claim $\hat{J}(\mathcal{D}) = \hat{J}(\mathcal{D})$

$\langle X \rangle = \langle Y \rangle - q \langle Z \rangle$ (0-smoothing, 1-smoothing)

$\langle O^k \rangle = (q + q^{-1})^k$

$\hat{J}(L) = (-1)^n q^{n+2n} \langle L \rangle$

(n_+, n_-) count (\nearrow, \searrow)

Indeed, $\langle \mathcal{D} \rangle = \langle \mathcal{D} \rangle - q \langle \mathcal{D} \rangle - 9 \langle \mathcal{D} \rangle + 9^2 \langle \mathcal{D} \rangle = -9 \langle \mathcal{D} \rangle$

"God created the knots, all else in topology is the work of man."

This handout is at <http://www.math.toronto.edu/~drorbn/Talks/Oporto-0407>

More at <http://www.math.toronto.edu/~drorbn/Talks/Oporto-0407/>