
Math 1352 Algebraic Knot Theory – The Knizhnik-Zamolodchikov Connection

Theorem 1. The following is an invariant of braids in $\mathbb{R}_t \times \mathbb{I}_z$ (Fixed endpoints)

$$Z(B) = \oint_{\substack{t_1<\dots<t_m \\ p=(\{z_i,z_i'\})}} \frac{Dp}{(2\pi i)^m} \bigwedge_{i=1}^{m} \frac{dz_i - dz_i'}{z_i - z_i'} \quad \text{in} \quad \mathcal{A}(\mathbb{1}_n) := \left\langle t^{ij} : 1\le i\ne j\le n \right\rangle \Big/ \begin{matrix} t^{ij}=t^{ji}, \\ [t^{ij},t^{kl}]=0 \\ [t^{ij},t^{ik}+t^{jk}]=0 \end{matrix}$$



$$= \left\langle \text{horizontal chords.} \right\rangle / 4T$$

= "Chord diagrams for braids".

Formal Connection & Curvature.

Let $\Omega \in \Omega^1(M,\mathcal{A})$ with $\deg \Omega = 1$. (graded unital.)

$\gamma : [0,1] = I \to M$ induces

$\phi : \Delta^m = \{0 \le t_1 \le \dots \le t_m \le 1\} \to M^m$.

Set $\mathrm{hol}_\gamma(\Omega) = P\exp\int_\gamma \Omega = \oint_{\Delta^m} \phi^* \Omega^m$

where $\Omega^m := \pi_1^*\Omega \wedge \dots \wedge \pi_m^*\Omega$

Theorem 2. If $F_\Omega := d\Omega + \Omega\wedge\Omega = 0$, then $\mathrm{hol}_\gamma(\Omega)$ is invariant under end-point preserving homotopies of γ.

The KZ connection.

$M = \mathbb{C}^n \setminus \{\text{diagonals}\}$, $\mathcal{A} = \mathcal{A}(\mathbb{1}_n)$,

and $\Omega = \sum_{i<j} t^{ij} w_{ij}$ where $w_{ij} := \frac{dz_i - dz_j}{z_i - z_j} = d\log(z_i - z_j)$ locally

Compute $F_\Omega = d\Omega + \Omega\wedge\Omega$: $dw_{ij} = 0$ so $d\Omega = 0$.

$$\Omega\wedge\Omega = \sum_{\substack{i<j \\ k<l}} t^{ij}t^{kl} w_{ij}\wedge w_{kl} = \underbrace{A}_{|\{i,j,k,l\}|=2} + \underbrace{B}_{=3} + \underbrace{C}_{=4} \quad \text{where}$$

$A = C = 0$ as $[t^{ij}, t^{kl}] = 0$ if $|\{i,j,k,l\}| = 2$ or 4 and

$$B = \sum_{\alpha<\beta<\gamma} [t^{\alpha\beta}, t^{\beta\gamma}] w_{\alpha\beta}\wedge w_{\beta\gamma} + \text{cyclic perms}$$

Note: by 4T, $[t^{\alpha\beta}, t^{\beta\gamma}] = r^{\alpha\beta\gamma}$ = 

$$= \sum_{\alpha<\beta<\gamma} r^{\alpha\beta\gamma} \underbrace{\left(w_{\alpha\beta}\wedge w_{\beta\gamma} + \text{cyc perms}\right)}_{=0 \text{ by Arnold's identity}} = 0$$

Proof 2. Let $\Gamma : I_s \times I_t \to M$, $\Phi : I_s \times \Delta^m \to M^m$;

By Stokes',

$$\int_{\Delta^m} \Phi_1^*\Omega^m - \int_{\Delta^m} \Phi_0^*\Omega^m = \int_{I\times\Delta^m} d\Phi^*\Omega^m - \int_{I\times\partial\Delta^m} \Phi^*\Omega^m =: A_m - B_m$$

Now

$$A_m = \sum_{k=1}^{m} (-1)^{k+1} \int_{I\times\Delta^m} \pi_1^*\Omega\wedge\dots\wedge\pi_k^*\,d\Omega\wedge\dots\wedge\pi_m^*\Omega$$

and

$$B_m = \int_{I\times[t_1=0]} \Phi^*\Omega^m \pm \int_{I\times[t_m=1]} \Phi^*\Omega^m + \sum_{k=1}^{m-1} (-1)^k \int_{I\times[t_k=t_{k+1}]} \Phi^*\Omega^m$$

(the first two terms vanish: $\Phi(I\times\{0\})$ and $\Phi(I\times\{1\})$ are fixed endpoints)

$$= \sum_{k=1}^{m-1} (-1)^k \int_{I\times\Delta^{m-1}} \pi_1^*\Omega\wedge\dots\wedge\pi_k^*(\Omega\wedge\Omega)\wedge\dots\wedge\pi_{m-1}^*\Omega$$

and now $\sum A_m = \sum B_m$ by telescopic summation & $F_\Omega = 0$.

Proof of 1

Simply take in theorem 2,

γ = the braid

and

Ω = the KZ connection.

Dror Bar-Natan, Feb 13, 2007