

# ON THE LOCAL CONSTANCY OF CHARACTERS

JONATHAN KORMAN

ABSTRACT. The character of an irreducible admissible representation of a  $p$ -adic reductive group is known to be a constant function in some neighborhood of any regular semisimple element  $\gamma$  in the group. Under certain mild restrictions on  $\gamma$ , we give an explicit description of a neighborhood of  $\gamma$  on which the character is constant.

## INTRODUCTION

Let  $k$  be a  $p$ -adic field of characteristic zero, and let  $\mathbf{G}$  be a connected reductive algebraic group defined over  $k$ . We denote by  $G$  the group of  $k$ -rational points of  $\mathbf{G}$ , and by  $\mathfrak{g}$  the Lie algebra of  $G$ . Let  $\pi$  be an irreducible admissible representation of  $G$ , and let  $\Theta_\pi$  be the (distribution) character of  $\pi$ . In [7] Harish-Chandra showed that  $\Theta_\pi$  can be represented by a function (also denoted by  $\Theta_\pi$ ) which is locally integrable on  $G$  and locally constant on the set  $G^{reg}$  of regular semisimple elements in  $G$ . Thus for any  $\gamma \in G^{reg}$  there exists *some* neighborhood of  $\gamma$  on which the character is constant. In [8, Theorem 2, p. 483], R. Howe gave an elementary proof of Harish-Chandra's result for general linear groups. In this paper we give a precise version of local constancy (near compact regular semisimple tame elements) for all reductive groups. The outline of the approach given here follows the elementary argument of Howe.

Let  $\mathfrak{g}_{x,r}$  (resp.  $G_{x,|r|}$ ) be the Moy-Prasad lattices [10] in  $\mathfrak{g}$  (resp. open compact subgroups of  $G$ ), normalized as in [9, §1.2]. Let  $G_{cpt}$  denote the set of compact elements in  $G$ . For a maximal  $k$ -torus  $T$ , let  $T_r$  denote its filtration subgroups (section 0.3). Let  $\rho(\pi)$  denote the depth of  $\pi$  [10, §5].

Fix a regular semisimple element  $\gamma$  and let  $\mathbf{T} := C_{\mathbf{G}}(\gamma)^\circ$  be the connected component of its centralizer;  $\mathbf{T}$  is a maximal  $k$ -torus in  $\mathbf{G}$ . We assume that it splits over some tamely ramified finite Galois extension  $E$  of  $k$ . Let  $T$  denote the group of  $k$ -rational points of  $\mathbf{T}$ . When  $\gamma \in T \cap G_{cpt}$  we attach to it the nonnegative rational number  $s(\gamma)$ . Using the filtration subgroups  $T_r$  and the parameter  $s(\gamma)$ , we characterize a neighborhood of  $\gamma$  on which the character  $\Theta_\pi$  is constant. Whether or not this neighborhood of constancy is maximal is not addressed here.

The main result of this paper is the following (Theorem 20).

---

*Date:* October 9, 2005.

*1991 Mathematics Subject Classification.* Primary 22E50; Secondary 22E35, 20G25.

*Key words and phrases.* characters, reductive  $p$ -adic groups, Bruhat-Tits building.

**Theorem.** *Let  $r = \max\{s(\gamma), \rho(\pi)\} + s(\gamma)$ . The character  $\Theta_\pi$  is constant on the set  $G(\gamma T_{r+})$ .*

We now give a brief sketch of the proof. Let  $K$  be any open compact subgroup of  $G$ . Decompose  $\Theta_\pi$  into a countable sum of ‘partial trace’ operators  $\Theta_d$ , according to the irreducible representations  $d$  of  $K$  (see section 3). For  $G = GL_n$ , Howe proved [8, p. 499] the following key fact. If  $X$  is a compact subset of  $G^{reg}$ , then  $\Theta_d$  vanishes on  $X$  for all  $d$  not in a certain finite set  $F$  (which depends only on  $X$ ). It follows (see proof of Theorem 20), that  $\Theta_\pi(f) = \int_X (\sum_{d \in F} \Theta_d)(x) f(x) dx$  for all  $f \in C_c^\infty(X)$ . Hence  $\Theta_\pi$  is represented on  $X$  by the locally constant function  $\sum_{d \in F} \Theta_d$ .

The main part of this paper is concerned with formulating an analogue of the above key fact for reductive groups (see Corollary 18).

The rational number  $s(\gamma)$ , defined in section 1, is used (Corollary 18) to make a precise choice of a set  $X$  and a subgroup  $K$ . Corollary 18 characterizes a finite set  $F$  of representations, such that for all  $d$  not in  $F$ ,  $\Theta_d$  vanishes on  $X$  (see Remark 14 for the significance of this fact). Thus the representations  $d \in F$  are those which play a role in understanding the character  $\Theta_\pi$  near  $\gamma$ . The proof of this corollary relies on a special case (Corollary 13), in which we only consider 1-dimensional  $d$ . Such representations have an explicit description in terms of cosets in the lie algebra  $\mathfrak{g}$ . In section 2 we develop the technical tools, using Moy-Prasad lattices, to handle these cosets. Once we have a characterization of the set  $F$ , we can make precise statements about the neighbourhood of constancy of the character near  $\gamma$  (Theorem 20).

**Acknowledgments.** I would like to thank Fiona Murnaghan, Jeff Adler, Stephen DeBacker, Ju-Lee Kim and Joe Repka for helpful conversations and comments. I would also like to thank the referees for their careful reading and detailed comments.

## NOTATION AND CONVENTIONS

Let  $k$  be a  $p$ -adic field (a finite extension of some  $\mathbb{Q}_p$ ) with residue field  $\mathbb{F}_{p^n}$ . Let  $\nu$  be a valuation on  $k$  normalized such that  $\nu(k^\times) = \mathbb{Z}$ .

For any algebraic extension field  $E$  of  $k$ ,  $\nu$  extends uniquely to a valuation (also denoted  $\nu$ ) of  $E$ .

We denote the ring of integers in  $E$  by  $R_E$  (write  $R$  for  $R_k$ ), and the prime ideal in  $R_E$  by  $\wp_E$  (write  $\wp$  for  $\wp_k$ ).

Let  $\mathbf{G}$  be a connected reductive group defined over  $k$ , and  $\mathbf{G}(E)$  the group of  $E$ -rational points of  $\mathbf{G}$ . We denote by  $G$  the group of  $k$ -rational points of  $\mathbf{G}$ . Denote the Lie algebras of  $\mathbf{G}$  and  $\mathbf{G}(E)$  by  $\mathfrak{g}$  and  $\mathfrak{g}(E)$ , respectively. Write  $\mathfrak{g}$  for the Lie algebra of  $k$ -rational points of  $\mathfrak{g}$ .

Let  $\mathcal{N}$  be the set of nilpotent elements in  $\mathfrak{g}$ . There are different notions of nilpotency, but since we assume that  $\text{char}(k) = 0$ , these notions all coincide.

Let  $\text{Ad}$  (resp.  $\text{ad}$ ) denote the adjoint representation of  $\mathbf{G}$  (resp.  $\mathfrak{g}$ ) on its Lie algebra  $\mathfrak{g}$ . For elements  $g \in G$  and  $X \in \mathfrak{g}$  (resp.  $x \in G$ ) we will

sometimes write  ${}^gX$  (resp.  ${}^gx$ ) instead of  $\text{Ad}(g)X$  (resp.  $gxg^{-1}$ ). For a subset  $S$  of  $\mathfrak{g}$  (resp.  $G$ ) let  ${}^G S$  denote the set  $\{{}^g s \mid g \in G \text{ and } s \in S\}$ .

Let  $n$  denote the (absolute) rank of  $\mathbf{G}$ . We say that an element  $g \in G$  is *regular semisimple* if the coefficient of  $t^n$  in  $\det(t - 1 + \text{Ad}(g))$  is nonzero. We denote the set of regular semisimple elements in  $G$  by  $G^{\text{reg}}$ . Similarly we say that an element  $X \in \mathfrak{g}$  is *regular semisimple* if the coefficient of  $t^n$  in  $\det(t - \text{ad}(X))$  is nonzero. We denote the set of regular semisimple elements in  $\mathfrak{g}$  by  $\mathfrak{g}^{\text{reg}}$ . Let  $G_{\text{cpt}}$  denotes the set of compact elements in  $G$ . For a subset  $S$  of  $G$  we will sometimes write  $S_{\text{cpt}}$  for  $S \cap G_{\text{cpt}}$ .

For a subset  $S$  of  $\mathfrak{g}$  (resp.  $G$ ) let  $[S]$  denote the characteristic function of  $S$  on  $\mathfrak{g}$  (resp.  $G$ ).

For any compact group  $K$ , let  $K^\wedge$  denote the set of equivalence classes of irreducible, continuous representations of  $K$ .

Let  $\pi$  be an irreducible admissible representation of  $G$ . We denote by  $\Theta_\pi$  the character of  $\pi$  thought of as a locally constant function on the set  $G^{\text{reg}}$ . Let  $\rho(\pi)$  denote the depth of  $\pi$  [10, §5].

## 0. PRELIMINARIES

**0.1. Apartments and buildings.** For a finite extension  $E$  of  $k$ , let  $\mathcal{B}(\mathbf{G}, E)$  denote the extended Bruhat-Tits building of  $\mathbf{G}$  over  $E$ ; write  $\mathcal{B}(G)$  for  $\mathcal{B}(\mathbf{G}, k)$ . It is known (e.g. [13]) that if  $E$  is a tamely ramified finite Galois extension of  $k$  then  $\mathcal{B}(\mathbf{G}, k)$  can be embedded into  $\mathcal{B}(\mathbf{G}, E)$  and its image is equal to the set of Galois fixed points in  $\mathcal{B}(\mathbf{G}, E)$ . If  $\mathbf{T}$  is a maximal  $k$ -torus in  $\mathbf{G}$  that splits over  $E$ , let  $\mathcal{A}(\mathbf{T}, E)$  be the corresponding apartment over  $E$ . Let  $\mathbf{X}^*(\mathbf{T}, E)$  (resp.  $\mathbf{X}_*(\mathbf{T}, E)$ ) denote the group of  $E$ -rational characters (resp. cocharacters) of  $\mathbf{T}$ .

It is known in the tame case [1, §1.9] that there is a Galois equivariant embedding of  $\mathcal{B}(\mathbf{T}, E)$  into  $\mathcal{B}(\mathbf{G}, E)$ , which in turn induces an embedding of  $\mathcal{B}(\mathbf{T}, k)$  into  $\mathcal{B}(\mathbf{G}, k)$ . Such embeddings are only unique modulo translations by elements of  $\mathbf{X}_*(\mathbf{T}, k) \otimes \mathbb{R}$ , however their images are all the same and are equal to the set  $\mathcal{A}(\mathbf{T}, E) \cap \mathcal{B}(\mathbf{G}, k)$ . From now on we fix a  $T$ -equivariant embedding  $i : \mathcal{B}(\mathbf{T}, k) \rightarrow \mathcal{B}(\mathbf{G}, k)$ , and use it to regard  $\mathcal{B}(\mathbf{T}, k)$  as a subset of  $\mathcal{B}(\mathbf{G}, k)$ ; write  $x$  for  $i(x)$ .

**Notation.** We write  $\mathcal{A}(\mathbf{T}, k)$  for  $\mathcal{A}(\mathbf{T}, E) \cap \mathcal{B}(\mathbf{G}, k)$ . This is well defined independent of the choice of  $E$  [15]. Moreover,  $\mathcal{A}(\mathbf{T}, k)$  is the set of Galois fixed points in  $\mathcal{A}(\mathbf{T}, E)$ .

We remark that the image of  $\mathcal{B}(\mathbf{T}, E)$  in  $\mathcal{B}(\mathbf{G}, E)$  is the apartment  $\mathcal{A}(\mathbf{T}, E)$ , while the image of  $\mathcal{B}(\mathbf{T}, k)$  in  $\mathcal{B}(\mathbf{G}, k)$  is the set  $\mathcal{A}(\mathbf{T}, k)$ .

**0.2. Moy-Prasad filtrations.** Regarding  $\mathbf{G}$  as a group over  $E$ , Moy and Prasad (see [10] and [11]) define lattices in  $\mathfrak{g}(E)$  and subgroups of  $\mathbf{G}(E)$ .

We can and will normalize (with respect to the normalized valuation  $\nu$ ) the indexing  $(x, r) \in \mathcal{B}(\mathbf{G}, E) \times \mathbb{R}$  of these lattices and subgroups as in [9, §1.2]. We will denote the (normalized) lattices by  $\mathfrak{g}(E)_{x,r}$ , and the (normalized) subgroups by  $\mathbf{G}(E)_{x,|r|}$ .

If  $\varpi_E$  is a uniformizing element of  $E$ , and  $e = e(E/k)$  is the ramification index of  $E$  over  $k$ , then these normalized lattices (resp. subgroups) satisfy  $\varpi_E \mathfrak{g}(E)_{x,r} = \mathfrak{g}(E)_{x,r+\frac{1}{e}}$ . Write  $\mathfrak{g}_{x,r}$  (resp.  $G_{x,|r|}$ ) for  $\mathfrak{g}(E)_{x,r}$  (resp.  $\mathbf{G}(E)_{x,|r|}$ ).

The above normalization was chosen to have the following property [1, 1.4.1]: when  $E$  is a tamely ramified Galois extension of  $k$  and  $x \in \mathcal{B}(\mathbf{G}, k) \subset \mathcal{B}(\mathbf{G}, E)$ , we have

$$(1) \quad \mathfrak{g}_{x,r} = \mathfrak{g}(E)_{x,r} \cap \mathfrak{g}, \quad \text{and (for } r > 0) \quad G_{x,r} = \mathbf{G}(E)_{x,r} \cap G.$$

We will also use the following notation. Let  $r \in \mathbf{R}$  and  $x \in \mathcal{B}(G)$ .

- $\mathfrak{g}_{x,r+} = \bigcup_{s>r} \mathfrak{g}_{x,s}$  and  $G_{x,|r|+} = \bigcup_{s>|r|} G_{x,s}$ .
- $G_r = \bigcup_{x \in \mathcal{B}(G)} G_{x,r}$  and  $G_{r+} = \bigcup_{s>r} G_s$  for  $r \geq 0$ .

The lattices  $\mathfrak{g}_{x,r+}$  (resp. groups  $G_{x,|r|+}$ ) have analogous properties to those of  $\mathfrak{g}_{x,r}$  (resp.  $G_{x,|r|}$ ). The set  $G_0$  is the set of compact elements  $G_{cpt}$ . We remark that  $G_{cpt} \subset \mathbf{G}(E)_{cpt} \cap G$ , and in general they need not be equal [3, §2.2.3].

**Lemma 1.** *Let  $\gamma$  be a compact regular semisimple element, and consider the maximal  $k$ -torus  $\mathbf{T} := C_{\mathbf{G}}(\gamma)^\circ$ . Suppose that  $\mathbf{T}$  splits over a tamely ramified finite Galois extension  $E$  of  $k$ . Then  $\gamma$  fixes  $\mathcal{B}(\mathbf{T}, k)$  pointwise.*

*Proof.* Recall that  $\gamma$  acts on  $\mathcal{A}(\mathbf{T}, E)$  by translations [14, §1]. Since  $\gamma$  belongs to a compact subgroup, it has a fixed point  $x \in \mathcal{B}(\mathbf{G}, E)$ .

If  $\gamma$  acts on  $\mathcal{A}(\mathbf{T}, E)$  by a nontrivial translation, then for any  $y \in \mathcal{A}(\mathbf{T}, E)$  there is an  $n \in \mathbb{N}$  such that  $d(x, y) \neq d(x, \gamma^n \cdot y)$ . This contradicts the fact that the action preserves distances. So  $\gamma$  must act trivially on  $\mathcal{A}(\mathbf{T}, E)$ . In particular,  $\gamma$  fixes  $\mathcal{A}(\mathbf{T}, k)$ , and hence  $\mathcal{B}(\mathbf{T}, k)$ , pointwise.  $\square$

**0.3. Root decomposition.** Let  $\mathbf{T}$  be a maximal  $k$ -torus in  $\mathbf{G}$  that splits over a tamely ramified finite Galois extension  $E$  of  $k$ . Let  $\Phi(\mathbf{T}, E)$  denote the set of roots of  $\mathbf{G}$  with respect to  $E$  and  $\mathbf{T}$ , and let  $\Psi(\mathbf{T}, E)$  denote the corresponding set of affine roots of  $\mathbf{G}$  with respect to  $E$ ,  $\mathbf{T}$  and  $\nu$ . When  $\mathbf{T}$  is  $k$ -split, we also write  $\Phi(\mathbf{T})$  for  $\Phi(\mathbf{T}, k)$  (resp.  $\Psi(\mathbf{T})$  for  $\Psi(\mathbf{T}, k)$ ). If  $\psi \in \Psi(\mathbf{T}, E)$ , let  $\dot{\psi} \in \Phi(\mathbf{T}, E)$  be the gradient of  $\psi$ , and let  $\mathfrak{g}(E)_{\dot{\psi}} \subset \mathfrak{g}(E)$  be the root space corresponding to  $\dot{\psi}$ . We denote the root lattice in  $\mathfrak{g}(E)_{\dot{\psi}}$  corresponding to  $\psi$  by  $\mathfrak{g}(E)_{\psi}$  [10, 3.2].

For  $x \in \mathcal{A}(\mathbf{T}, E)$  and  $r \in \mathbb{R}$ , let  $\mathfrak{t}(E)_r := \mathfrak{t}(E) \cap \mathfrak{g}(E)_{x,r}$  and  $\mathfrak{t}(E)_{r+} := \mathfrak{t}(E) \cap \mathfrak{g}(E)_{x,r+}$ . Note that  $\mathfrak{t}(E)_r$  and  $\mathfrak{t}(E)_{r+}$  are defined independent of the choice of  $x \in \mathcal{A}(\mathbf{T}, E)$ . Similarly one defines the subgroups  $\mathbf{T}(E)_r$  and  $\mathbf{T}(E)_{r+}$  for  $r \geq 0$ ; they have analogous properties. Note that using our conventions we will sometimes denote  $\mathbf{T}(E)_0$  by  $\mathbf{T}(E)_{cpt}$ .

An alternative description is [9, §2.1]: for  $r \in \mathbb{R}$ ,

$$\mathfrak{t}(E)_r = \{\Gamma \in \mathfrak{t}(E) \mid \nu(d\chi(\Gamma)) \geq r \text{ for all } \chi \in \mathbf{X}^*(\mathbf{T}, E)\}$$

and for  $r > 0$ ,

$$\mathbf{T}(E)_r = \{t \in \mathbf{T}(E) \mid \nu(\chi(t) - 1) \geq r \text{ for all } \chi \in \mathbf{X}^*(\mathbf{T}, E)\}.$$

Since  $\mathbf{G}$  splits over  $E$ , we have

$$\begin{aligned}\mathfrak{g}(E)_{x,r} &= \mathfrak{t}(E)_r \oplus \sum_{\psi \in \Psi(\mathbf{T}, E), \psi(x) \geq r} \mathfrak{g}(E)_\psi, \\ \mathfrak{g}(E)_{x,r+} &= \mathfrak{t}(E)_{r+} \oplus \sum_{\psi \in \Psi(\mathbf{T}, E), \psi(x) > r} \mathfrak{g}(E)_\psi.\end{aligned}$$

Let  $\mathfrak{t} := \text{Lie}(T)$ , and define  $\mathfrak{t}^\perp := (\text{Ad}(\gamma) - 1)\mathfrak{g}$ . We have the following decomposition [7, §18]

$$(2) \quad \mathfrak{g} = \mathfrak{t} \oplus \mathfrak{t}^\perp.$$

We write  $X = Y + Z$  with respect to this decomposition; when convenient, we also write  $X_t$  for  $Y$ .

Fix  $x \in \mathcal{B}(\mathbf{T}, k) \subset \mathcal{B}(\mathbf{G}, k)$  and  $r \in \mathbb{R}$ . Write  $\mathfrak{t}_r$  for  $\mathfrak{t} \cap \mathfrak{g}_{x,r}$  (resp.  $\mathfrak{t}_{r+}$  for  $\mathfrak{t} \cap \mathfrak{g}_{x,r+}$ ); as mentioned earlier, these definitions are independent of  $x$ . Define  $\mathfrak{t}_{x,r}^\perp := \mathfrak{t}^\perp \cap \mathfrak{g}_{x,r}$  (resp.  $\mathfrak{t}_{x,r+}^\perp := \mathfrak{t}^\perp \cap \mathfrak{g}_{x,r+}$ ). We have [1, 1.9.3],

$$(3) \quad \begin{aligned}\mathfrak{g}_{x,r} &= \mathfrak{t}_r \oplus \mathfrak{t}_{x,r}^\perp, \\ \mathfrak{g}_{x,r+} &= \mathfrak{t}_{r+} \oplus \mathfrak{t}_{x,r+}^\perp.\end{aligned}$$

**0.4. Hypotheses.** (HB) There is a nondegenerate  $G$ -invariant symmetric bilinear form  $B$  on  $\mathfrak{g}$  such that we can identify  $\mathfrak{g}_{x,r}^*$  with  $\mathfrak{g}_{x,r}$  via the map  $\Omega : \mathfrak{g} \rightarrow \mathfrak{g}^*$  defined by  $\Omega(X)(Y) = B(X, Y)$ .

Groups satisfying the above hypothesis are discussed in [4, §4].

Fix  $r \in \mathbb{R}_{>0}$  and  $x \in \mathcal{B}(\mathbf{G}, k)$ . For any  $r \leq t \leq 2r$  the group  $(G_{x,r}/G_{x,t})$  is abelian. By hypothesis (HB), there exists a ( $G_{x,0}$ -equivariant) isomorphism (see [1, §1.7] or [12, p.16])

$$(4) \quad (G_{x,r}/G_{x,t})^\wedge \cong \mathfrak{g}_{x,(-t)+}/\mathfrak{g}_{x,(-r)+}.$$

## 1. REGULAR DEPTH

From now on let  $\gamma \in G^{reg}$ , and assume that the  $k$ -torus  $\mathbf{T} := C_{\mathbf{G}}(\gamma)^\circ$  splits over a tamely ramified finite Galois extension  $E$  of  $k$ . We attach to  $\gamma$  the following rational number  $s(\gamma)$ .

**Definition 2.** For each  $\alpha \in \Phi(\mathbf{T}, E)$  let  $s_\alpha(\gamma) := \nu(\alpha(\gamma) - 1)$  and define  $s(\gamma) := \max\{s_\alpha(\gamma) \mid \alpha \in \Phi(\mathbf{T}, E)\}$ .

**Remark 3.** Note that  $s(\gamma)$  is not the same as the depth of  $\gamma$  (as defined in [2]). But for good elements [1, §2.2], these two notions agree.

**Remark 4.** A priori  $s(\gamma) \in \mathbb{Q} \cup \{+\infty\}$ , but since  $\gamma$  is regular,  $\alpha(\gamma) \neq 1$  for all  $\alpha \in \Phi(\mathbf{T}, E)$  and so  $s(\gamma) \in \mathbb{Q}$ . If  $\gamma$  is compact then  $s(\gamma) \geq 0$ . Also note that  $s(\gamma z) = s(\gamma)$  for all  $z$  in the center  $Z(G)$  of  $G$  and that  $s(g\gamma g^{-1}) = s(\gamma)$  for all  $g \in G$ .

We will need the following basic properties of  $s(\gamma)$ .

**Lemma 5.** *Suppose  $\gamma \in T_{cpt}$  and  $\gamma' \in T_{s(\gamma)+}$ .*

- (1)  $s(\gamma\gamma') = s(\gamma)$  and for  $\alpha \in \Phi(\mathbf{T}, E)$ , we have  $|\alpha(\gamma\gamma') - 1| = |\alpha(\gamma) - 1|$ .
- (2)  $\gamma\gamma' \in T_{cpt}$ .

*Proof.* (1) Fix  $r > s(\gamma) \geq 0$  such that  $T_r = T_{s(\gamma)+}$ . With this notation  $\gamma' \in T_r$ . By the alternative description of  $T_r$ , for any  $\chi \in \mathbf{X}^*(\mathbf{T}, E)$ ,  $\chi(\gamma') = 1 + \mu'$  where  $\nu(\mu') \geq r$ . Thus for any  $\alpha \in \Phi(\mathbf{T}, E)$ ,  $\alpha(\gamma') = 1 + \lambda'$  where  $\nu(\lambda') \geq r$ .

Note that since each  $\alpha \in \Phi(\mathbf{T}, E)$  is continuous,  $\alpha(T(E)_{cpt}) \subset R_E^\times$ . Since  $\gamma \in T_{cpt} \subset T(E)_{cpt}$  we get that  $\alpha(\gamma)$  is a unit.

Now  $\alpha(\gamma\gamma') - 1 = \alpha(\gamma)\alpha(\gamma') - 1 = \alpha(\gamma)(1 + \lambda') - 1 = (\alpha(\gamma) - 1) + \alpha(\gamma)\lambda'$ . Using  $\nu(\alpha(\gamma) - 1) =: s_\alpha(\gamma)$ ,  $\alpha(\gamma)$  is a unit, and  $\nu(\lambda') \geq r > s(\gamma) \geq s_\alpha(\gamma)$ , we have  $\nu(\alpha(\gamma\gamma') - 1) = \nu(\alpha(\gamma) - 1)$  (or equivalently  $|\alpha(\gamma\gamma') - 1| = |\alpha(\gamma) - 1|$ ) for all  $\alpha \in \Phi(\mathbf{T}, E)$ . Thus  $s(\gamma\gamma') := \max_\alpha \{\nu(\alpha(\gamma\gamma') - 1)\} = \max_\alpha \{\nu(\alpha(\gamma) - 1)\} =: s(\gamma)$ .

- (2) Since  $\gamma$  and  $\gamma'$  are in  $T_{cpt}$ , so is their product. □

**Corollary 6.** *Let  $\gamma \in T$  be a compact regular semisimple element. Then  $\gamma T_{s(\gamma)+} \subset G^{reg}$ .*

*Proof.* For  $t \in T \cap G^{reg}$ , following [7, §18], define

$$D_{G/T}(t) := \det(\text{Ad}(t) - 1)|_{\mathfrak{g}/\mathfrak{t}} = \prod_{\alpha \in \Phi(\mathbf{T}, E)} (\alpha(t) - 1).$$

Then  $t \in T \cap G^{reg} \Leftrightarrow D_{G/T}(t) \neq 0 \Leftrightarrow |D_{G/T}(t)| \neq 0$ . Using Lemma 5 with  $\gamma \in T \cap G_{cpt}$  and  $\gamma' \in T_{s(\gamma)+}$ , we get  $|D_{G/T}(\gamma\gamma')| = \prod_\alpha |\alpha(\gamma\gamma') - 1| = \prod_\alpha |\alpha(\gamma) - 1| = |D_{G/T}(\gamma)| \neq 0$ . □

## 2. SOME TECHNICAL LEMMAS

The next lemma will generalize the following example.

**Example 7.**  $\mathbf{G} = \mathbf{GL}_2$ ,  $\mathbf{T}$  a  $k$ -split maximal torus. Choose  $x_0 \in \mathcal{B}(\mathbf{G}, k)$  so that  $G_{x_0,0} = GL_2(R)$ . Any  $X \in \mathcal{N} \cap (\mathfrak{g}_{x_0,r} \setminus \mathfrak{g}_{x_0,r+})$  is of the form  ${}^k \begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix}$ ,

for some  $k \in G_{x_0,0}$  (see [5, 9.2.1]). Thus

$$\begin{aligned} X &= \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 0 & x \\ 0 & 0 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} \\ &= \frac{x}{ad-bc} \begin{pmatrix} -ac & a^2 \\ -c^2 & ac \end{pmatrix}. \end{aligned}$$

Write  $X = Y + Z$  as in (2) and note that the depth of  $X$  with respect to the filtration  $\{\mathfrak{g}_{x_0,r}\}_{r \in \mathbb{R}}$  of  $\mathfrak{g}$  is controlled by  $Z$ . This is the case since  $\max\{\nu(a^2), \nu(-c^2)\} \geq \nu(ac)$  and  $ad - bc \in R^\times$ .

**Lemma 8.** *Fix  $x \in \mathcal{B}(\mathbf{T}, k)$  and  $r \in \mathbb{R}$ . For  $X \in \mathcal{N} \cap (\mathfrak{g}_{x,r} \setminus \mathfrak{g}_{x,r+})$ , write  $X = Y + Z$  as in (2). Then  $Z \in \mathfrak{g}_{x,r} \setminus \mathfrak{g}_{x,r+}$ .*

*Proof.* We first prove the case when the maximal  $k$ -torus  $\mathbf{T}$  is  $k$ -split and then reduce the general case to this case.

**Split case.** Assume  $\mathbf{T}$  is  $k$ -split. Note that  $\mathfrak{t}^\perp = \bigoplus_{\alpha \in \Phi(\mathbf{T})} \mathfrak{g}_\alpha$ . Fix a system of simple roots  $\Delta$  in  $\Phi(\mathbf{T})$  and choose a Chevalley basis for  $\mathfrak{g}$  as in [1, §1.2]. Such a basis contains elements  $H_b$  and  $E_b$  in  $\mathfrak{g}$  for each  $b \in \Phi(\mathbf{T})$ . If  $\mathbf{G}$  is semisimple, then the set  $\{H_b \mid b \in \Delta\} \cup \{E_b \mid b \in \Phi(\mathbf{T})\}$  is a basis for  $\mathfrak{g}$ . These elements also satisfy the commutation relations listed in [1, 1.2.1]. With respect to this choice of Chevalley basis, the adjoint representation is determined by the following formulas [1, 1.2.5]:

$$(5) \quad \begin{cases} \text{Ad}(e_b(\lambda))E_c &= \begin{cases} E_b & \text{if } c = b \\ E_c + \lambda H_b - \lambda^2 E_b & \text{if } c = -b \\ \sum_{i \geq 0} M_{b,c;i} \lambda^i E_{ib+c} & \text{if } c \neq \pm b \end{cases} \\ \text{Ad}(t)E_c &= c(t)E_c \\ \text{Ad}(e_b(\lambda))H &= H - db(H)\lambda E_b \\ \text{Ad}(t)H &= H \end{cases}$$

for all  $H \in \text{Lie}(T)$ , all  $t \in T$  and all  $\lambda \in k$ . Here  $e_b$  is the unique map  $e_b : \mathbf{Add} \rightarrow \mathbf{G}$  such that  $de_b(1) = E_b$  ( $de_b$  is the derivative of  $e_b$ ); and  $M_{b,c;i}$  are constants with  $M_{b,c;0} = 1$ .

Let  $B$  be the Borel subgroup associated to  $\Delta$  (with Levi decomposition  $B = TN$  and opposite Borel  $\bar{B} = T\bar{N}$ ). We have  $\mathfrak{g} = \mathfrak{t} \oplus \mathfrak{n} \oplus \bar{\mathfrak{n}}$ , where  $\mathfrak{n} := \text{Lie}(N)$  and  $\bar{\mathfrak{n}} := \text{Lie}(\bar{N})$ . Note that  $\mathfrak{n} \oplus \bar{\mathfrak{n}} = \bigoplus_{\alpha \in \Phi(\mathbf{T})} \mathfrak{g}_\alpha = \mathfrak{t}^\perp$ . Recall that  $G_{x,0}$  acts on  $\mathfrak{g}_{x,r}$  (and on  $\mathfrak{g}_{x,r+}$ ).

Given  $X \in \mathcal{N} \cap (\mathfrak{g}_{x,r} \setminus \mathfrak{g}_{x,r+})$ , we can use [2, Proposition 3.5.1] (with  $T$  playing the role of  $M$ ) to conclude that there exists a group element  $n \in N \cap G_{x,0}$  such that  $({}^n X)_\mathfrak{t} \in \mathfrak{t}_{r+}$  (where  ${}^n X$  denotes  $\text{Ad}(n)X$ ).

Write  $X = Y + Z$  as in (2) and assume for a contradiction that  $Z \in \mathfrak{g}_{x,r+}$ . Since  $X \in \mathfrak{g}_{x,r} \setminus \mathfrak{g}_{x,r+}$ , the assumption implies that  $Y \in \mathfrak{t} \cap (\mathfrak{g}_{x,r} \setminus \mathfrak{g}_{x,r+}) = \mathfrak{t}_r \setminus \mathfrak{t}_{r+}$ .

Using the properties (5) of the Chevalley basis, one can easily check that the set  $(\mathfrak{t}_r \setminus \mathfrak{t}_{r+}) \oplus \mathfrak{n}$  is preserved under the action of  $\text{Ad}(e_b(\lambda))$  for all  $b \in \Phi^+(\mathbf{T})$ , where  $\Phi^+(\mathbf{T})$  are the positive roots with respect to  $\Delta$ . Since

$\{e_b(\lambda) \mid b \in \Phi^+(\mathbf{T})\}$  generates  $N$ , we conclude that  ${}^n Y \in (\mathfrak{t}_r \setminus \mathfrak{t}_{r+}) \oplus \mathfrak{n}$ , and hence that  $({}^n Y)_\mathfrak{t} \in \mathfrak{t}_r \setminus \mathfrak{t}_{r+}$ .

On the other hand we have  ${}^n X = {}^n Y + {}^n Z$ , where  ${}^n Z \in \mathfrak{g}_{x,r+}$ . Taking the  $\mathfrak{t}$  components, we get,  $({}^n X)_\mathfrak{t} = ({}^n Y)_\mathfrak{t} + ({}^n Z)_\mathfrak{t}$ , with  $({}^n Z)_\mathfrak{t} \in \mathfrak{t}_{r+}$ . Since  $({}^n X)_\mathfrak{t} \in \mathfrak{t}_{r+}$ , we conclude that  $({}^n Y)_\mathfrak{t} \in \mathfrak{t}_{r+}$ . This contradicts  $({}^n Y)_\mathfrak{t} \in \mathfrak{t}_r \setminus \mathfrak{t}_{r+}$ .

Hence  $Z \in \mathfrak{g}_{x,r} \setminus \mathfrak{g}_{x,r+}$  (note that from the decomposition (3) it is clear that  $Z \in \mathfrak{g}_{x,r}$ ).

**General case.** We now assume  $\mathbf{T}$  is an  $E$ -split maximal  $k$ -torus. Define  $\mathfrak{t}(E)^\perp := (\text{Ad}(\gamma) - 1)\mathfrak{g}(E)$ . We have the following analogue of (2)

$$(6) \quad \mathfrak{g}(E) = \mathfrak{t}(E) \oplus \mathfrak{t}(E)^\perp.$$

Note that  $\mathfrak{t} \subset \mathfrak{t}(E)$  and  $\mathfrak{t}^\perp \subset \mathfrak{t}(E)^\perp$ . So the decomposition  $X = Y + Z$  (as in (2)) for  $X \in \mathfrak{g}$  is the same whether viewed in  $\mathfrak{g}$  or in  $\mathfrak{g}(E)$ .

Since  $X \in \mathfrak{g}_{x,r} \setminus \mathfrak{g}_{x,r+}$ , equations (1) imply that  $X \in (\mathfrak{g}(E)_{x,r} \setminus \mathfrak{g}(E)_{x,r+}) \cap \mathfrak{g}$ . Since  $X \in \mathcal{N} \subset \mathcal{N}(E)$  (where  $\mathcal{N}(E)$  is the set of nilpotent elements in  $\mathfrak{g}(E)$ ), we have that  $X \in \mathcal{N}(E) \cap (\mathfrak{g}(E)_{x,r} \setminus \mathfrak{g}(E)_{x,r+})$ . Now since  $\mathbf{T}$  splits over  $E$  we can regard  $\mathbf{G}$  over  $E$  as a split group and hence apply all the constructions of the split case above. So by the considerations of the split case above we conclude that  $Z \in \mathfrak{g}(E)_{x,r} \setminus \mathfrak{g}(E)_{x,r+}$ . Intersecting with  $\mathfrak{g}$  gives  $Z \in \mathfrak{g}_{x,r} \setminus \mathfrak{g}_{x,r+}$ .  $\square$

From now on we assume that  $\gamma$  is also compact. Recall that this implies that  $s(\gamma) \geq 0$  (see Remark 4).

**Lemma 9.** *Let  $t \in \mathbb{R}$  and  $x \in \mathcal{B}(\mathbf{T}, k)$ . If  $Z \in \mathfrak{t}^\perp \cap (\mathfrak{g}_{x,-t} \setminus \mathfrak{g}_{x,(-t)+})$  then  ${}^\gamma Z - Z \notin \mathfrak{g}_{x,(-t+s(\gamma))+}$ .*

*Proof.* Using the root decomposition  $\mathfrak{t}(E)^\perp = \bigoplus_{\alpha \in \Phi(\mathbf{T}, E)} \mathfrak{g}(E)_\alpha$ , for  $Z \in \mathfrak{t}^\perp \subset \mathfrak{t}(E)^\perp$  we write  $Z = \sum Z_\alpha$ . Then  ${}^\gamma Z - Z = \sum ({}^\gamma Z_\alpha - Z_\alpha) = \sum (\alpha(\gamma) - 1)Z_\alpha$ .

By assumption  $Z \notin \mathfrak{g}_{x,(-t)+}$ , hence (see equations (1))  $Z \notin \mathfrak{g}(E)_{x,(-t)+}$ . Thus for some  $\alpha \in \Phi(\mathbf{T}, E)$ ,  $Z_\alpha \notin \mathfrak{g}(E)_{x,(-t)+}$ , and so by definition of  $s_\alpha(\gamma)$ ,  $(\alpha(\gamma) - 1)Z_\alpha \notin \mathfrak{g}(E)_{x,(-t+s_\alpha(\gamma))+}$ . It follows by definition of  $s(\gamma)$ , that  $(\alpha(\gamma) - 1)Z_\alpha \notin \mathfrak{g}(E)_{x,(-t+s(\gamma))+}$ . Hence  ${}^\gamma Z - Z = \sum (\alpha(\gamma) - 1)Z_\alpha \notin \mathfrak{g}(E)_{x,(-t+s(\gamma))+}$ . Intersecting with  $\mathfrak{g}$  we get that  ${}^\gamma Z - Z \notin \mathfrak{g}_{x,(-t+s(\gamma))+}$ .  $\square$

**Proposition 10.** *Let  $r \in \mathbb{R}$  and  $x \in \mathcal{B}(\mathbf{T}, k)$ . If  $X \in \mathcal{N} \cap \mathfrak{g}_{x,(-2r)+}$  satisfies  ${}^\gamma X - X \in \mathfrak{g}_{x,(-r)+}$ , then  $X \in \mathfrak{g}_{x,(-r-s(\gamma))+}$ .*

*Proof.* Fix  $t < 2r$  such that  $X \in \mathcal{N} \cap (\mathfrak{g}_{x,-t} \setminus \mathfrak{g}_{x,(-t)+})$ .

Write  $X = Y + Z$  as in (2). By Lemma 8,  $Z \in \mathfrak{t}^\perp \cap (\mathfrak{g}_{x,-t} \setminus \mathfrak{g}_{x,(-t)+})$ , and so by Lemma 9,  ${}^\gamma Z - Z \notin \mathfrak{g}_{x,(-t+s(\gamma))+}$ .

On the other hand, since  $\gamma$  acts trivially on  $Y$  (because  $Y \in \mathfrak{t} = C_{\mathfrak{g}}(\gamma)$ ),  ${}^\gamma Z - Z = {}^\gamma X - X \in \mathfrak{g}_{x,(-r)+}$ .

Thus  $-t + s(\gamma) > -r$ , or equivalently  $-t > -r - s(\gamma)$ , which implies that  $X \in \mathfrak{g}_{x,-t} \subseteq \mathfrak{g}_{x,(-r-s(\gamma))+}$ .  $\square$



**Definition 11.** A character  $d \in (G_{x,r}/G_{x,2r})^\wedge$  is called degenerate if under the isomorphism (4), the corresponding coset  $X + \mathfrak{g}_{x,(-r)_+}$  contains nilpotent elements.

**Definition 12.** Let  $K$  be a compact subgroup of  $G$  and  $d \in K^\wedge$ . For  $g \in G$ , let  ${}^g d$  denote the representation of  $gKg^{-1}$  defined as  ${}^g d(gkg^{-1}) := d(k)$ . We say that  $g$  intertwines  $d$  with itself if upon restriction to  $gKg^{-1} \cap K$ ,  $d$  and  ${}^g d$  contain a common representation (up to isomorphism) of  $gKg^{-1} \cap K$ .

**Corollary 13.** Let  $x \in \mathcal{B}(\mathbf{T}, k)$ ,  $r \in \mathbb{R}_{>0}$ , and assume  $d \in (G_{x,r}/G_{x,2r})^\wedge$  is degenerate. If  $\gamma$  intertwines  $d$  with itself then  $d \in (G_{x,r}/G_{x,r+s(\gamma)})^\wedge$ .

*Proof.* Let  $X + \mathfrak{g}_{x,(-r)_+}$  be the coset in  $\mathfrak{g}_{x,(-2r)_+}/\mathfrak{g}_{x,(-r)_+}$  corresponding to  $d$  under the isomorphism (4). Since this coset is degenerate, we can assume that  $X \in \mathcal{N}$ .

Since  $\gamma$  fixes  $x$  (Lemma 1),  $\gamma$  stabilizes  $G_{x,r}$ . Thus having  $\gamma$  intertwine  $d$  with itself amounts to having  $d \cong \gamma d$ ; or furthermore, since  $d$  is one-dimensional,  $d = \gamma d$ . Under the isomorphism (4), we get  $X + \mathfrak{g}_{x,(-r)_+} = \gamma(X + \mathfrak{g}_{x,(-r)_+})$ , or equivalently that  $\gamma X - X \in \mathfrak{g}_{x,(-r)_+}$ . Now apply Proposition 10 to conclude that  $X \in \mathfrak{g}_{x,(-r-s(\gamma))_+}$ , which under the isomorphism (4) gives that  $d \in (G_{x,r}/G_{x,r+s(\gamma)})^\wedge$ .  $\square$

### 3. PARTIAL TRACES

Let  $(\pi, V)$  be an irreducible admissible representation of  $G$ . Let  $K$  be an open compact subgroup of  $G$ . Let  $V = \bigoplus_{d \in K^\wedge} V_d$  be the decomposition of  $V$  into  $K$ -isotypic components. Let  $E_d$  denote the  $K$ -equivariant projection from  $V$  to  $V_d$ . For  $f \in C_c^\infty(G)$  define the distribution  $\Theta_d(f) := \text{trace}(E_d \pi(f) E_d)$ , the ‘partial trace of  $\pi$  with respect to  $d$ ’. The distribution  $\Theta_d$  is represented by the locally constant function  $\Theta_d(x) := \text{trace}(E_d \pi(x) E_d)$  on  $G$ . Recall that it is known that the distribution  $\Theta_\pi(f) := \text{trace} \pi(f)$  is also represented by a locally constant function,  $\Theta_\pi$ , on  $G^{reg}$ ; we will not use this fact here. It follows from the definitions that as distributions

$$\Theta_\pi(f) = \sum_{d \in K^\wedge} \Theta_d(f) \text{ for all } f \in C_c^\infty(G).$$

**Remark 14.** For (some)  $\omega \subset G^{reg}$  compact, Corollary 19 and the proof of Theorem 20 will imply that, for all  $f \in C_c^\infty(\omega)$ , this sum is *finite*.

**Lemma 15.**  $\Theta_d(kxk^{-1}) = \Theta_d(x)$  for all  $x \in G$  and all  $k \in K$ .

*Proof.* Since  $E_d$  is  $K$ -equivariant, it commutes with  $\pi(k)$  for all  $k \in K$ .

$$\begin{aligned} \Theta_d(kxk^{-1}) &= \text{trace}(E_d \pi(kxk^{-1}) E_d) \\ &= \text{trace}(E_d \pi(k) \pi(x) \pi(k^{-1}) E_d) \\ &= \text{trace}(\pi(k) E_d \pi(x) E_d \pi(k^{-1})) \\ &= \text{trace}(E_d \pi(x) E_d) = \Theta_d(x). \end{aligned}$$

$\square$

Let  $N$  be an open compact subgroup of  $G$  which is normal in  $K$ . Suppose  $g \in G$  normalizes  $K$  and  $N$ . Let  $d \in K^\wedge$ . Considered as a representation of  $N$ ,  $d$  decomposes into a finite sum of irreducible representations

$$d_1 \oplus \cdots \oplus d_n.$$

**Proposition 16.** *Suppose  $\Theta_d(g) \neq 0$ . Then  $d \cong {}^g d$  as representations of  $K$  and also for some  $i \in \{1, \dots, n\}$ ,  $d_i \cong {}^g d_i$  as representations of  $N$ .*

*Proof.* We refer to the appendix. Since  $g$  permutes the  $V_{d'}$ 's (Theorem 22.1),  $0 \neq \Theta_d(g) = \text{trace}(E_d \pi(g) E_d)$  implies that  $g$  must stabilize  $V_d$ . Fix a decomposition (‡) as in Theorem 22.2, and let  $E_{W_i}$  denote the  $K$ -equivariant projections onto  $W_i$ . Since  $E_d = \sum E_{W_i}$ ,  $\text{trace}(E_d \pi(g) E_d) \neq 0$  implies that for some  $i$ ,  $g$  must stabilize  $W_i$ , and that  $\text{trace}(E_{W_i} \pi(g) E_{W_i}) \neq 0$ . By Theorem 22.3,  $d \cong {}^g d$ , which proves the first part of the theorem.

Now as a representation of  $N$ ,

$$W_i \cong_N \bigoplus_j W_{i,d_j},$$

where  $W_{i,d_j}$  are the  $d_j$ -isotypic components of  $W_i$ . Since  $g$  stabilizes  $N$ , it must permute the  $W_{i,d_j}$ 's (Theorem 22.1). Since  $E_{W_i} = \sum E_{W_{i,d_j}}$ , having  $\text{trace}(E_{W_i} \pi(g) E_{W_i}) \neq 0$  implies that for some  $j$ ,  $g$  must stabilize  $W_{i,d_j}$ , and that  $\text{trace}(E_{W_{i,d_j}} \pi(g) E_{W_{i,d_j}}) \neq 0$ . Fix a decomposition (‡) as in Theorem 22.2 for  $W_{i,d_j}$ :

$$W_{i,d_j} \cong_N \bigoplus d_j.$$

Since  $E_{W_{i,d_j}} = \sum E_{d_j}$ ,  $\text{trace}(E_{W_{i,d_j}} \pi(g) E_{W_{i,d_j}}) \neq 0$  implies that  $g$  must stabilize one of the  $d_j$ 's. By Theorem 22.3,  $d_j \cong {}^g d_j$ , which proves the second part of the theorem.  $\square$

The following theorem and corollaries are used in the proof of Theorem 20 to show that for  $f$  with compact support, the sum  $\sum_{d \in K^\wedge} \Theta_d(f)$  is finite (see also Remark 14).

**Theorem 17.** *Fix  $x \in \mathcal{B}(\mathbf{T}, k)$  and let  $r > \max\{s(\gamma), \rho(\pi)\}$ . If  $d \in (G_{x,r})^\wedge$  satisfies  $\Theta_d(\gamma) \neq 0$ , then  $d \in (G_{x,r}/G_{x,r+s(\gamma)})^\wedge$ .*

*Proof.* If  $d$  is trivial we are done, so assume it is not. Let  $t$  be the smallest number such that  $d|_{G_{x,t+}}$  is trivial (so in particular  $d|_{G_{x,t}}$  is nontrivial).

**Case  $t < 2r$ :** Pick  $s \leq 2r$  such that  $G_{x,s} = G_{x,t+}$ . Consider  $d$  as an element of  $(G_{x,r}/G_{x,2r})^\wedge$ . By Proposition 16,  $\Theta_d(\gamma) \neq 0$  implies that  $d \cong {}^\gamma d$ . Also,  $\Theta_d(\gamma) \neq 0$  implies that  $d \subset \pi|_{G_{x,r}}$ ; since  $r > \rho(\pi)$  this means that  $d$  is degenerate (see [5, §7.6]). Now apply Corollary 13.

**Case  $t \geq 2r$ :** Note that  $\frac{t}{2} \geq r > s(\gamma)$ . For  $\epsilon > 0$  such that  $\frac{t}{2} > \frac{\epsilon}{2} + s(\gamma)$ , let  $s = t + \epsilon$ . By making  $\epsilon$  smaller if necessary, we can make sure that  $G_{x,s} = G_{x,t+}$ . Note that  $t > \frac{t}{2} + \frac{\epsilon}{2} + s(\gamma) = \frac{s}{2} + s(\gamma)$ .

Since  $\frac{s}{2} > \frac{t}{2} \geq r$  it makes sense to restrict  $d$  to  $G_{x, \frac{s}{2}}$  and think of it as an element of  $(G_{x, \frac{s}{2}}/G_{x, s})^\wedge$ . As a representation of  $G_{x, \frac{s}{2}}/G_{x, s}$ ,  $d$  decomposes into a finite sum of irreducible (one-dimensional) representations

$$d_1 \oplus \cdots \oplus d_n.$$

Let  $X_i + \mathfrak{g}_{x, (-\frac{s}{2})_+}$  be the coset in  $\mathfrak{g}_{x, (-s)_+}/\mathfrak{g}_{x, (-\frac{s}{2})_+}$  corresponding to  $d_i$  under the isomorphism (4).

By Proposition 16,  $0 \neq \Theta_d(\gamma)$  implies that for some  $j$ ,  $d_j \cong \gamma d_j$ .

Now  $d \subset \pi|_{G_{x, r}}$ , implies that  $d_j \subset \pi|_{G_{x, \frac{s}{2}}}$  and since  $\frac{s}{2} > r > \rho(\pi)$  we have that  $d_j$  is degenerate. Apply Corollary 13 to  $d_j$  to conclude that  $d_j \in (G_{x, \frac{s}{2}}/G_{x, \frac{s}{2}+s(\gamma)})^\wedge$ . In particular  $d_j$  is trivial on  $G_{x, \frac{s}{2}+s(\gamma)}$ , and hence on  $G_{x, t}$ .

Since  $G_{x, r}$  normalizes  $G_{x, \frac{s}{2}}$ , it acts by permutations on the  $d_i$ 's. Since  $d$  is irreducible, this action is transitive. Hence all the  $d_i$ 's are conjugate by elements of  $G_{x, r}$ . By the conjugation of the  $d_i$ 's and the fact that  $d_j|_{G_{x, t}} = 1$  it follows that  $d_i|_{G_{x, t}} = 1$  for all  $i$ , and so  $d$  itself is trivial on  $G_{x, t}$ . This contradicts the definition of  $t$ . Hence this case is not possible and  $t < 2r$ .  $\square$

**Corollary 18.** *Fix  $x \in \mathcal{B}(\mathbf{T}, k)$  and let  $r > \max\{s(\gamma), \rho(\pi)\}$ . Let  $X$  denote  $\gamma T_{r+s(\gamma)}$ , a compact subset of  $T \cap G^{reg}$ . If  $d \in (G_{x, r})^\wedge$  satisfies  $\Theta_d(\gamma') \neq 0$  for some  $\gamma' \in X$ , then  $d \in (G_{x, r}/G_{x, r+s(\gamma)})^\wedge$ .*

*Proof.* Lemma 5 implies that  $\gamma'$  fixes  $x$  and that  $s(\gamma')=s(\gamma)$ . Now apply Theorem 17 to  $\gamma'$ .  $\square$

**Corollary 19.** *Fix  $x \in \mathcal{B}(\mathbf{T}, k)$  and let  $r > \max\{s(\gamma), \rho(\pi)\}$ . Let  $\omega$  denote  $G_{x, r}(\gamma T_{r+s(\gamma)})$ , an open compact subset of  $G^{reg}$ . Then  $\Theta_d$  vanishes on  $\omega$  for all  $d \notin (G_{x, r}/G_{x, r+s(\gamma)})^\wedge$ . Furthermore,  $\Theta_d(x) = \Theta_d(\gamma)$  for all  $x \in \omega$  and all  $d \in (G_{x, r}/G_{x, r+s(\gamma)})^\wedge$ .*

*Proof.* Follows immediately from Lemma 15 and Corollary 18.  $\square$

#### 4. PROOF OF THE MAIN THEOREM

Let  $r > \max\{s(\gamma), \rho(\pi)\}$ . Denote the finite set  $(G_{x, r}/G_{x, r+s(\gamma)})^\wedge$  by  $F$ .

**Theorem 20.** *The distribution  $\Theta_\pi$  is represented on the set  $G(\gamma T_{r+s(\gamma)})$  by a constant function.*

*Proof.* Using Corollary 19, we have for all  $f \in C_c^\infty(G)$  whose support is contained in  $\omega$ ,

$$\begin{aligned} \Theta_\pi(f) &= \sum_{d \in (G_{x, r})^\wedge} \Theta_d(f) = \sum_{d \in F} \int_\omega \Theta_d(x) f(x) dx = \sum_{d \in F} \int_\omega \Theta_d(\gamma) f(x) dx \\ &= \int_\omega \left( \sum_{d \in F} \Theta_d(\gamma) \right) f(x) dx. \end{aligned}$$

Thus  $\Theta_\pi$  is represented by the constant function  $\sum_{d \in F} \Theta_d(\gamma)$  on  $\omega$ , i.e.  $\Theta_\pi(x) = \sum_{d \in F} \Theta_d(\gamma)$  for all  $x \in \omega$ . Since  $\Theta_\pi$  is conjugation invariant, we get  $\Theta_\pi(gxg^{-1}) = \Theta_\pi(x) = \sum_{d \in F} \Theta_d(\gamma)$  for all  $x \in \omega$  and all  $g \in G$ .  $\square$

**Remark 21.** This gives a new proof of the local constancy (near compact regular semisimple tame elements  $\gamma$ ) of the character of an irreducible admissible representation for an arbitrary reductive  $p$ -adic group  $G$ .

## 5. APPENDIX

We prove some variations of Clifford theory [6, §14]. Let  $K$  and  $N$  be open compact subgroups of  $G$ , such that  $N$  is a normal subgroup of  $K$ . Let  $(\pi, V)$  be an irreducible admissible representation of  $G$  and let

$$(\dagger) \quad V \underset{K}{=} \bigoplus_{d \in K^\wedge} V_d$$

be the (canonical) decomposition of  $V$  into  $K$ -isotypic components. Here  $V_d$  denotes the  $d$ -isotypic component of  $V$ , i.e. the sum of all the  $K$ -submodules of  $V$  isomorphic to  $d = (d, W)$ . Each isotypic component  $V_d$  decomposes (non-canonically) into a finite sum of isomorphic copies  $W_i \underset{K}{\cong} W$  of  $(d, W)$

$$(\ddagger) \quad V_d \underset{K}{\cong} \bigoplus_i W_i.$$

**Theorem 22.** *Suppose  $g \in G$  normalizes  $K$  and  $N$ . Then*

- (1) *The action of  $g$  permutes the  $V_d$ 's.*
- (2) *Suppose  $g$  stabilizes  $V_d$ . Then there exists a decomposition  $(\ddagger)$  such that the action of  $g$  permutes the  $W_i$ .*
- (3) *Suppose  $W'$  is a  $K$ -submodule of  $V$ , isomorphic to  $W$ , and stable under the action  $g$ . Then  ${}^g d \cong d$ .*

*Proof.* (1) This follows from the fact that for any two  $K$ -submodules  $W'$  and  $W''$  of  $V$ , if  $W' \underset{K}{\cong} W''$  then  $gW' \underset{K}{\cong} gW''$ .

- (2) Let  $W'$  be an irreducible  $K$ -submodule of  $V_d$ , isomorphic to  $W$ . Since  $g$  normalizes  $K$  and stabilizes  $V_d$ ,  $gW'$  is a  $K$ -submodule of  $V_d$ . Since  $W'$  is irreducible, so is  $gW'$ . As an irreducible submodule of  $V_d$ ,  $gW'$  must be isomorphic to  $W$ . By irreducibility either  $W' \cap gW' = \{0\}$  or  $W' = gW'$ . Thus the orbit of  $W'$  under  $g$  is a collection of subspaces with trivial pairwise intersection, and so  $g$  acts on their sum as a desired. By complete reducibility of  $V_d$  (being a finite-dimensional representation of the compact group  $K$ ) we can now use induction on the dimension of  $V_d$ .
- (3) This follows from the following commutative diagram (in which all the arrows are isomorphisms of vector spaces and  $k \in K$ ).

$$\begin{array}{ccccccccc}
W & \longrightarrow & W' & \xrightarrow{\pi(g)} & gW' & \xlongequal{\quad} & W' & \longrightarrow & W \\
d(k) \downarrow & & \pi(k) \downarrow & & \pi(gkg^{-1}) \downarrow & & \pi(k^g) \downarrow & & d(k^g) \downarrow \quad {}^g d(k) \\
W & \longrightarrow & W' & \xrightarrow{\pi(g)} & gW' & \xlongequal{\quad} & W' & \longrightarrow & W
\end{array}$$

□

## REFERENCES

- [1] J. Adler, *Refined anisotropic  $K$ -types and supercuspidal representations*, Pacific J. Math **185** (1998), pp. 1–32.
- [2] J. Adler and S. DeBacker, *Some applications of Bruhat-Tits theory to harmonic analysis on the Lie algebra of a reductive  $p$ -adic group*, Michigan Math. J. **50** (2002), no. 2, pp. 263–286.
- [3] ———, *Murnaghan-Kirillov theory for supercuspidal representations of tame general linear groups*, J. Reine Angew. Math. **575** (2004).
- [4] J. Adler and A. Roche, *An intertwining result for  $p$ -adic groups*, Canad. J. Math., vol. **52** (2000), no. 3, pp. 449–467.
- [5] S. DeBacker, *Lectures on harmonic analysis for reductive  $p$ -adic groups*, Representation Theory of Lie Groups, Institute of Mathematical Sciences, National University of Singapore, 2002.
- [6] L. Dornhoff *Group representation theory, Part A*, Pure and applied mathematics, vol. **7**, Marcel Dekker Inc., 1971.
- [7] Harish-Chandra, *Admissible invariant distributions on reductive  $p$ -adic groups*, Notes by Stephen DeBacker and Paul J. Sally, Jr., University Lecture Series, **16**, Amer. Math. Soc., 1999.
- [8] R. Howe, *Some qualitative results on the representation theory of  $GL_n$  over a  $p$ -adic field*, Pacific Journal of Math., **73** (1977), pp. 479–538.
- [9] J.-L. Kim and F. Murnaghan, *Character expansions and unrefined minimal  $K$ -types*, Amer. J. Math., **125** (2003), no. 6, pp. 1199–1234.
- [10] A. Moy and G. Prasad, *Unrefined minimal  $K$ -types for  $p$ -adic groups*, Inv. Math. **116** (1994), pp. 393–408.
- [11] ———, *Jacquet functors and unrefined minimal  $K$ -types*, Comment. Math. Helvetici **71** (1996), pp. 98–121.
- [12] F. Murnaghan, *Characters of admissible representations of reductive  $p$ -adic groups*, Lecture notes for the thematic year (1999) on number theory at the CRM, Montreal.
- [13] G. Prasad, *Galois fixed points in the Bruhat-Tits buildings of a reductive group*, Bull. Soc. Math. France, **129** (2001), pp. 169–174.
- [14] J. Tits, *Reductive groups over local fields*, Automorphic Forms, Representation and  $L$ -functions. Proc. Symp. Pure Math. 33 Vol. 1 (1979), 29–69.
- [15] J.-K. Yu, *Construction of tame supercuspidal representations*, J. Amer. Math. Soc., **14** (2001), no. 3, pp. 579–622.

*E-mail address:* jkorman@math.toronto.edu

THE UNIVERSITY OF TORONTO, TORONTO, ONTARIO M5S 3G3