

Shannon's theorem and Poisson boundaries for locally compact groups

Giulio Tiozzo
University of Toronto

June 17, 2020

Summary

1. The Poisson representation formula - classical case

Summary

1. The Poisson representation formula - classical case
2. Poisson representation for groups

Summary

1. The Poisson representation formula - classical case
2. Poisson representation for groups
3. The ray (and strip) criterion

Summary

1. The Poisson representation formula - classical case
2. Poisson representation for groups
3. The ray (and strip) criterion
4. The Shannon-McMillan-Breiman theorem

Summary

1. The Poisson representation formula - classical case
2. Poisson representation for groups
3. The ray (and strip) criterion
4. The Shannon-McMillan-Breiman theorem
5. Proofs

Summary

1. The Poisson representation formula - classical case
2. Poisson representation for groups
3. The ray (and strip) criterion
4. The Shannon-McMillan-Breiman theorem
5. Proofs


joint with Behrang Forghani

The Poisson representation formula

$$h^\infty(\mathbb{D}) := \{u : \mathbb{D} \rightarrow \mathbb{R} \text{ bounded} : \Delta u = 0\}$$


The Poisson representation formula

$$h^\infty(\mathbb{D}) := \{u : \mathbb{D} \rightarrow \mathbb{R} \text{ bounded} : \Delta u = 0\}$$



The Poisson representation formula

$$h^\infty(\mathbb{D}) := \{u : \mathbb{D} \rightarrow \mathbb{R} \text{ bounded} : \Delta u = 0\}$$



Theorem (Poisson representation)

There is a bijection

$$h^\infty(\mathbb{D}) \quad \leftrightarrow \quad L^\infty(S^1, \lambda)$$

The Poisson representation formula

Theorem

Given an integrable function $f : S^1 \rightarrow \mathbb{R}$,

The Poisson representation formula

Theorem

Given an integrable function $f : S^1 \rightarrow \mathbb{R}$, the function $u : \mathbb{D} \rightarrow \mathbb{R}$

The Poisson representation formula

Theorem

Given an integrable function $f : S^1 \rightarrow \mathbb{R}$, the function $u : \mathbb{D} \rightarrow \mathbb{R}$

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(\theta - t) f(e^{it}) dt$$

is *harmonic* on \mathbb{D} , i.e. $\Delta u = 0$.

The Poisson representation formula

Theorem

Given an integrable function $f : S^1 \rightarrow \mathbb{R}$, the function $u : \mathbb{D} \rightarrow \mathbb{R}$

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(\theta - t) f(e^{it}) dt$$

is *harmonic* on \mathbb{D} , i.e. $\Delta u = 0$. Here,

$$P_r(\theta) := \frac{1 - r^2}{1 + r^2 - 2r \cos \theta}$$

is the *Poisson kernel*.

The Poisson representation formula - II

$$h^\infty(\mathbb{D}) := \{u : \mathbb{D} \rightarrow \mathbb{R} \text{ bounded} : \Delta u = 0\}$$

The Poisson representation formula - II

$$h^\infty(\mathbb{D}) := \{u : \mathbb{D} \rightarrow \mathbb{R} \text{ bounded} : \Delta u = 0\}$$

Theorem (Poisson)

There is a bijection

$$h^\infty(\mathbb{D}) \quad \leftrightarrow \quad L^\infty(\partial\mathbb{D}, \lambda)$$

The Poisson representation formula - II

$$h^\infty(\mathbb{D}) := \{u : \mathbb{D} \rightarrow \mathbb{R} \text{ bounded} : \Delta u = 0\}$$

Theorem (Poisson)

There is a bijection

$$h^\infty(\mathbb{D}) \quad \leftrightarrow \quad L^\infty(\partial\mathbb{D}, \lambda)$$

► \rightarrow boundary values

$$f(\xi) := \lim_{z \rightarrow \xi} u(z)$$

The Poisson representation formula - II

$$h^\infty(\mathbb{D}) := \{u : \mathbb{D} \rightarrow \mathbb{R} \text{ bounded} : \Delta u = 0\}$$

Theorem (Poisson)

There is a bijection

$$h^\infty(\mathbb{D}) \quad \leftrightarrow \quad L^\infty(\partial\mathbb{D}, \lambda)$$

► \rightarrow boundary values

$$f(\xi) := \lim_{z \rightarrow \xi} u(z)$$

► \leftarrow Poisson integral

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(\theta - t) f(e^{it}) dt$$

The Poisson representation formula - II

$$h^\infty(\mathbb{D}) := \{u : \mathbb{D} \rightarrow \mathbb{R} \text{ bounded} : \Delta u = 0\}$$

Theorem (Poisson)

There is a bijection

$$h^\infty(\mathbb{D}) \quad \leftrightarrow \quad L^\infty(\partial\mathbb{D}, \lambda)$$

- \rightarrow boundary values (probabilistic interpretation)

$$f(\xi) := \mathbb{E}\left(\lim_{z_t \rightarrow \xi} u(z_t)\right)$$

where z_t is Brownian motion

- \leftarrow Poisson integral

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(\theta - t) f(e^{it}) dt$$

The Poisson representation formula - II

Theorem (Poisson)


There is a bijection

$$h^\infty(\mathbb{D}) \quad \leftrightarrow \quad L^\infty(\partial\mathbb{D}, \lambda)$$

- \rightarrow boundary values (probabilistic interpretation)

$$f(\xi) := \mathbb{E}\left(\lim_{z_t \rightarrow \xi} u(z_t)\right)$$

where z_t is Brownian motion



The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$.

The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$. Given $a = re^{i\theta} \in \mathbb{D}$,

The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$. Given $a = re^{i\theta} \in \mathbb{D}$, consider

$$g_a(z) := \frac{z - a}{1 - \bar{a}z}$$

so that $g_a(a) = 0$,

The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$. Given $a = re^{i\theta} \in \mathbb{D}$, consider

$$g_a(z) := \frac{z - a}{1 - \bar{a}z}$$

so that $g_a(a) = 0$, hence

$$g'_a(z) := \frac{1 - |a|^2}{(1 - \bar{a}z)^2}$$

The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$. Given $a = re^{i\theta} \in \mathbb{D}$, consider

$$g_a(z) := \frac{z - a}{1 - \bar{a}z}$$

so that $g_a(a) = 0$, hence

$$g'_a(z) := \frac{1 - |a|^2}{(1 - \bar{a}z)^2}$$

and setting $z = e^{it}$

The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$. Given $a = re^{i\theta} \in \mathbb{D}$, consider

$$g_a(z) := \frac{z - a}{1 - \bar{a}z}$$

so that $g_a(a) = 0$, hence

$$g'_a(z) := \frac{1 - |a|^2}{(1 - \bar{a}z)^2}$$

and setting $z = e^{it}$ we obtain

$$\frac{dg_{a\lambda}}{d\lambda}(t)$$

The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$. Given $a = re^{i\theta} \in \mathbb{D}$, consider

$$g_a(z) := \frac{z - a}{1 - \bar{a}z}$$

so that $g_a(a) = 0$, hence

$$g'_a(z) := \frac{1 - |a|^2}{(1 - \bar{a}z)^2}$$

and setting $z = e^{it}$ we obtain

$$\frac{dg_{a\lambda}}{d\lambda}(t) = \left| g'_a(e^{it}) \right|$$

The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$. Given $a = re^{i\theta} \in \mathbb{D}$, consider

$$g_a(z) := \frac{z - a}{1 - \bar{a}z}$$

so that $g_a(a) = 0$, hence

$$g'_a(z) := \frac{1 - |a|^2}{(1 - \bar{a}z)^2}$$

and setting $z = e^{it}$ we obtain

$$\frac{dg_{a\lambda}}{d\lambda}(t) = \left| g'_a(e^{it}) \right| = \frac{1 - r^2}{|1 - re^{i(t-\theta)}|^2}$$

The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$. Given $a = re^{i\theta} \in \mathbb{D}$, consider

$$g_a(z) := \frac{z - a}{1 - \bar{a}z}$$

so that $g_a(a) = 0$, hence

$$g'_a(z) := \frac{1 - |a|^2}{(1 - \bar{a}z)^2}$$

and setting $z = e^{it}$ we obtain

$$\frac{dg_{a\lambda}}{d\lambda}(t) = \left| g'_a(e^{it}) \right| = \frac{1 - r^2}{|1 - re^{i(t-\theta)}|^2} = P_r(\theta - t)$$

The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$. Given $a = re^{i\theta} \in \mathbb{D}$, consider

$$g_a(z) := \frac{z - a}{1 - \bar{a}z}$$

so that $g_a(a) = 0$, hence

$$g'_a(z) := \frac{1 - |a|^2}{(1 - \bar{a}z)^2}$$

and setting $z = e^{it}$ we obtain

$$\frac{dg_{a\lambda}}{d\lambda}(t) = \left| g'_a(e^{it}) \right| = \frac{1 - r^2}{|1 - re^{i(t-\theta)}|^2} = P_r(\theta - t)$$

RN derivative of boundary action

The Poisson representation formula - III

Recall $PSL(2, \mathbb{R}) = Isom^+(\mathbb{H})$. Given $a = re^{i\theta} \in \mathbb{D}$, consider

$$g_a(z) := \frac{z - a}{1 - \bar{a}z}$$

so that $g_a(a) = 0$, hence

$$g'_a(z) := \frac{1 - |a|^2}{(1 - \bar{a}z)^2}$$

and setting $z = e^{it}$ we obtain

$$\frac{dg_{a\lambda}}{d\lambda}(t) = \left| g'_a(e^{it}) \right| = \frac{1 - r^2}{|1 - re^{i(t-\theta)}|^2} = P_r(\theta - t)$$

RN derivative of boundary action = Poisson kernel

The Poisson representation formula - IV

Recall the Poisson formula

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt$$

The Poisson representation formula - IV

Recall the Poisson formula

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt$$

Setting $a = re^{i\theta}$, $d\lambda = \frac{dt}{2\pi}$

The Poisson representation formula - IV

Recall the Poisson formula

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt$$

Setting $a = re^{i\theta}$, $d\lambda = \frac{dt}{2\pi}$

$$u(a) = \int_{\partial\mathbb{D}} f(e^{it}) \frac{dg_{a\lambda}}{d\lambda}(t) d\lambda(t)$$

The Poisson representation formula - IV

Recall the Poisson formula

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt$$

Setting $a = re^{i\theta}$, $d\lambda = \frac{dt}{2\pi}$

$$u(a) = \int_{\partial\mathbb{D}} f(e^{it}) \frac{dg_{a\lambda}}{d\lambda}(t) d\lambda(t)$$

Finally if $\xi = e^{it}$

The Poisson representation formula - IV

Recall the Poisson formula

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{it}) P_r(\theta - t) dt$$

Setting $a = re^{i\theta}$, $d\lambda = \frac{dt}{2\pi}$

$$u(a) = \int_{\partial\mathbb{D}} f(e^{it}) \frac{dg_{a\lambda}}{d\lambda}(t) d\lambda(t)$$

Finally if $\xi = e^{it}$

$$u(a) = \int_{\partial\mathbb{D}} f(\xi) dg_{a\lambda}(\xi)$$

The Poisson representation formula - IV

Theorem (Poisson representation)

If $f : L^\infty(\partial\mathbb{D}, \lambda) \rightarrow \mathbb{R}$, then

$$u(a) = \int_{\partial\mathbb{D}} f(\xi) \, dg_a\lambda(\xi)$$

is harmonic on \mathbb{D} .

The Poisson representation formula - IV

Theorem (Poisson representation)

If $f : L^\infty(\partial\mathbb{D}, \lambda) \rightarrow \mathbb{R}$, then

$$u(a) = \int_{\partial\mathbb{D}} f(\xi) dg_a\lambda(\xi)$$

is harmonic on \mathbb{D} .

Question. Can we generalize this to other groups
 $G \neq \mathrm{PSL}_2(\mathbb{R})$?

General Poisson representation (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G .

General Poisson representation (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G .

Definition

A function $u : G \rightarrow \mathbb{R}$ is μ -harmonic if

General Poisson representation (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G .

Definition

A function $u : G \rightarrow \mathbb{R}$ is μ -harmonic if

$$u(g) = \int_G u(gh) \, d\mu(h)$$

for all $g \in G$.

General Poisson representation (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G .

Definition

A function $u : G \rightarrow \mathbb{R}$ is μ -harmonic if

$$u(g) = \int_G u(gh) \, d\mu(h)$$

for all $g \in G$.

Let B be a space on which G acts (measurably).

General Poisson representation (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G .

Definition

A function $u : G \rightarrow \mathbb{R}$ is μ -harmonic if

$$u(g) = \int_G u(gh) \, d\mu(h)$$

for all $g \in G$.

Let B be a space on which G acts (measurably). A measure ν on B is μ -stationary if

$$\nu = \int_G g\nu \, d\mu(g).$$

Random walks and μ -boundaries

Let μ be a prob. measure on G .

Random walks and μ -boundaries

Let μ be a prob. measure on G .

Consider the random walk

Random walks and μ -boundaries

Let μ be a prob. measure on G .

Consider the random walk

$$w_n := g_1 g_2 \dots g_n$$

Random walks and μ -boundaries

Let μ be a prob. measure on G .

Consider the random walk

$$w_n := g_1 g_2 \dots g_n$$

where (g_i) are i.i.d. with distribution μ .

Random walks and μ -boundaries

Let μ be a prob. measure on G .

Consider the random walk

$$w_n := g_1 g_2 \dots g_n$$

where (g_i) are i.i.d. with distribution μ .

We denote $\Omega := (w_n)$ the space of sample paths, and

$T((w_n)) := (w_{n+1})$ is the shift on Ω .

Random walks and μ -boundaries

Let μ be a prob. measure on G .

Consider the random walk

$$w_n := g_1 g_2 \dots g_n$$

where (g_i) are i.i.d. with distribution μ .

We denote $\Omega := (w_n)$ the space of sample paths, and

$T((w_n)) := (w_{n+1})$ is the shift on Ω .

Definition

A space (B, ν) is a μ -boundary if there exists a measurable map

$$\text{bnd} : \Omega \rightarrow B$$

such that $\text{bnd} = \text{bnd} \circ T$.

Boundary convergence


Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$.

Boundary convergence

Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$. Let $o \in X$ a base point.

Boundary convergence

Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$. Let $o \in X$ a base point.




In most situations,

$$\lim_{n \rightarrow \infty} w_n o \in \partial X$$

exists a.s.

Boundary convergence

Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$. Let $o \in X$ a base point.




In most situations,

$$\lim_{n \rightarrow \infty} w_n o \in \partial X$$

exists a.s.

Boundary convergence

Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$. Let $o \in X$ a base point.




In most situations,

$$\lim_{n \rightarrow \infty} w_n o \in \partial X$$

exists a.s.

Boundary convergence

Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$. Let $o \in X$ a base point.




In most situations,

$$\lim_{n \rightarrow \infty} w_n o \in \partial X$$

exists a.s.

Boundary convergence

Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$. Let $o \in X$ a base point.




In most situations,

$$\lim_{n \rightarrow \infty} w_n o \in \partial X$$

exists a.s.

Boundary convergence

Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$. Let $o \in X$ a base point.




In most situations,

$$\lim_{n \rightarrow \infty} w_n o \in \partial X$$

exists a.s.

Boundary convergence

Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$. Let $o \in X$ a base point.




In most situations,

$$\lim_{n \rightarrow \infty} w_n o \in \partial X$$

exists a.s.

Boundary convergence

Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$. Let $o \in X$ a base point.




In most situations,

$$\lim_{n \rightarrow \infty} w_n o \in \partial X$$

exists a.s.

Boundary convergence

Suppose that $G < Isom(X, d)$ a metric space, and that X has a “bordification” $\overline{X} = X \cup \partial X$. Let $o \in X$ a base point.



In most situations,

$$\lim_{n \rightarrow \infty} w_n o \in \partial X$$

exists a.s.

Boundary convergence

Then we define the hitting measure

$$\nu(A) := \mathbb{P}(\lim_{n \rightarrow \infty} w_n o \in A)$$

Boundary convergence

Then we define the hitting measure

$$\nu(A) := \mathbb{P}(\lim_{n \rightarrow \infty} w_n o \in A)$$


which is μ -stationary.

Boundary convergence

Then we define the **hitting measure**

$$\nu(A) := \mathbb{P}\left(\lim_{n \rightarrow \infty} w_n o \in A\right)$$

which is μ -stationary.




Boundary convergence

Then we define the **hitting measure**

$$\nu(A) := \mathbb{P}(\lim_{n \rightarrow \infty} w_n o \in A)$$

which is μ -stationary.



Moreover, $(\partial X, \nu)$ is a **μ -boundary**, given by the map

$$\Omega \ni \text{bnd}(\omega) := \lim_{n \rightarrow \infty} w_n o \in \partial X$$

General Poisson representation - II (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G , let (B, ν) be a μ -boundary.

General Poisson representation - II (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G , let (B, ν) be a μ -boundary.

The **Poisson transform** $\Phi : L^\infty(B, \nu) \rightarrow H^\infty(G, \mu)$ is

General Poisson representation - II (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G , let (B, ν) be a μ -boundary.

The **Poisson transform** $\Phi : L^\infty(B, \nu) \rightarrow H^\infty(G, \mu)$ is

$$\Phi_f(g) := \int_B f \, dg\nu$$

General Poisson representation - II (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G , let (B, ν) be a μ -boundary.

The **Poisson transform** $\Phi : L^\infty(B, \nu) \rightarrow H^\infty(G, \mu)$ is

$$\Phi_f(g) := \int_B f \, dg\nu$$

Definition

A μ -boundary (B, ν) is the **Poisson boundary** if Φ is an isomorphism

$$L^\infty(B, \nu) \rightarrow H^\infty(G, \mu)$$

General Poisson representation - II (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G , let (B, ν) be a μ -boundary.

The **Poisson transform** $\Phi : L^\infty(B, \nu) \rightarrow H^\infty(G, \mu)$ is

$$\Phi_f(g) := \int_B f \, dg\nu$$

Definition

A μ -boundary (B, ν) is the **Poisson boundary** if Φ is an isomorphism

$$L^\infty(B, \nu) \rightarrow H^\infty(G, \mu)$$

Corollary

Poisson boundary is trivial (= 1 point) \Leftrightarrow bounded harmonic functions are constant

General Poisson representation - II (Furstenberg)

Let G be a (lcsc) group, μ a probability measure on G , let (B, ν) be a μ -boundary.

The **Poisson transform** $\Phi : L^\infty(B, \nu) \rightarrow H^\infty(G, \mu)$ is

$$\Phi_f(g) := \int_B f \, dg\nu$$

Definition

A μ -boundary (B, ν) is the **Poisson boundary** if Φ is an isomorphism

$$L^\infty(B, \nu) \rightarrow H^\infty(G, \mu)$$

Corollary

Poisson boundary is trivial (= 1 point) \Leftrightarrow bounded harmonic functions are constant

Examples. Abelian groups; nilpotent groups

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Question. Is $(\partial X, \nu)$ the Poisson boundary for (G, μ) ?

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Question. Is $(\partial X, \nu)$ the Poisson boundary for (G, μ) ?

(Some) History.

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Question. Is $(\partial X, \nu)$ the Poisson boundary for (G, μ) ?

(Some) History.

- ▶ Furstenberg '63: semisimple Lie groups

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Question. Is $(\partial X, \nu)$ the Poisson boundary for (G, μ) ?

(Some) History.

- ▶ Furstenberg '63: semisimple Lie groups
- ▶ Kaimanovich '94: hyperbolic groups

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Question. Is $(\partial X, \nu)$ the Poisson boundary for (G, μ) ?

(Some) History.

- ▶ Furstenberg '63: semisimple Lie groups
- ▶ Kaimanovich '94: hyperbolic groups
- ▶ Karlsson-Margulis '99: isometries of CAT(0) spaces

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Question. Is $(\partial X, \nu)$ the Poisson boundary for (G, μ) ?

(Some) History.

- ▶ Furstenberg '63: semisimple Lie groups
- ▶ Kaimanovich '94: hyperbolic groups
- ▶ Karlsson-Margulis '99: isometries of CAT(0) spaces
- ▶ Kaimanovich-Masur '99: mapping class group

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Question. Is $(\partial X, \nu)$ the Poisson boundary for (G, μ) ?

(Some) History.

- ▶ Furstenberg '63: semisimple Lie groups
- ▶ Kaimanovich '94: hyperbolic groups
- ▶ Karlsson-Margulis '99: isometries of CAT(0) spaces
- ▶ Kaimanovich-Masur '99: mapping class group
- ▶ Bader-Shalom '06: isometries of affine buildings

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Question. Is $(\partial X, \nu)$ the Poisson boundary for (G, μ) ?

(Some) History.

- ▶ Furstenberg '63: semisimple Lie groups
- ▶ Kaimanovich '94: hyperbolic groups
- ▶ Karlsson-Margulis '99: isometries of CAT(0) spaces
- ▶ Kaimanovich-Masur '99: mapping class group
- ▶ Bader-Shalom '06: isometries of affine buildings
- ▶ Gautero-Math  us '12: relatively hyperbolic groups

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Question. Is $(\partial X, \nu)$ the Poisson boundary for (G, μ) ?

(Some) History.

- ▶ Furstenberg '63: semisimple Lie groups
- ▶ Kaimanovich '94: hyperbolic groups
- ▶ Karlsson-Margulis '99: isometries of CAT(0) spaces
- ▶ Kaimanovich-Masur '99: mapping class group
- ▶ Bader-Shalom '06: isometries of affine buildings
- ▶ Gautero-Math  us '12: relatively hyperbolic groups
- ▶ Horbez '16 : $Out(F_n)$

Identification of the Poisson boundary

Let $G < Isom(X, d)$ be a group, μ a measure on G , ν the hitting measure on ∂X .

Question. Is $(\partial X, \nu)$ the Poisson boundary for (G, μ) ?

(Some) History.

- ▶ Furstenberg '63: semisimple Lie groups
- ▶ Kaimanovich '94: hyperbolic groups
- ▶ Karlsson-Margulis '99: isometries of CAT(0) spaces
- ▶ Kaimanovich-Masur '99: mapping class group
- ▶ Bader-Shalom '06: isometries of affine buildings
- ▶ Gautero-Math  us '12: relatively hyperbolic groups
- ▶ Horbez '16 : $Out(F_n)$
- ▶ Maher-T. '18: Cremona group

Ray approximation

Let G be a countable group and μ a measure on G .

Ray approximation

Let G be a countable group and μ a measure on G . A metric d on G is temperate if $\exists C$ s.t.

$$\#\{g \in G : d(o, go) \leq R\} \leq Ce^{CR}$$

Ray approximation

Let G be a countable group and μ a measure on G . A metric d on G is temperate if $\exists C$ s.t.

$$\#\{g \in G : d(o, go) \leq R\} \leq Ce^{CR}$$

The measure has finite first moment if $\int_G d(o, go) d\mu(g) < +\infty$.

Ray approximation

Let G be a countable group and μ a measure on G . A metric d on G is temperate if $\exists C$ s.t.

$$\#\{g \in G : d(o, go) \leq R\} \leq Ce^{CR}$$

The measure has finite first moment if $\int_G d(o, go) d\mu(g) < +\infty$.

Theorem (Kaimanovich \sim '94)

Let d be a temperate metric on G ,

Ray approximation

Let G be a countable group and μ a measure on G . A metric d on G is temperate if $\exists C$ s.t.

$$\#\{g \in G : d(o, go) \leq R\} \leq Ce^{CR}$$

The measure has finite first moment if $\int_G d(o, go) d\mu(g) < +\infty$.

Theorem (Kaimanovich \sim '94)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d ,

Ray approximation

Let G be a countable group and μ a measure on G . A metric d on G is temperate if $\exists C$ s.t.

$$\#\{g \in G : d(o, go) \leq R\} \leq Ce^{CR}$$

The measure has finite first moment if $\int_G d(o, go) d\mu(g) < +\infty$.

Theorem (Kaimanovich \sim '94)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary.

Ray approximation

Let G be a countable group and μ a measure on G . A metric d on G is temperate if $\exists C$ s.t.

$$\#\{g \in G : d(o, go) \leq R\} \leq Ce^{CR}$$

The measure has finite first moment if $\int_G d(o, go) d\mu(g) < +\infty$.

Theorem (Kaimanovich \sim '94)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary. If there are measurable functions $\pi_n : B \rightarrow G$

Ray approximation

Let G be a countable group and μ a measure on G . A metric d on G is temperate if $\exists C$ s.t.

$$\#\{g \in G : d(o, go) \leq R\} \leq Ce^{CR}$$

The measure has finite first moment if $\int_G d(o, go) d\mu(g) < +\infty$.

Theorem (Kaimanovich \sim '94)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary. If there are measurable functions $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability,

Ray approximation

Let G be a countable group and μ a measure on G . A metric d on G is temperate if $\exists C$ s.t.

$$\#\{g \in G : d(o, go) \leq R\} \leq Ce^{CR}$$

The measure has finite first moment if $\int_G d(o, go) d\mu(g) < +\infty$.

Theorem (Kaimanovich \sim '94)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary. If there are measurable functions $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability, then (B, λ) is the Poisson boundary.

Ray approximation

Theorem (Kaimanovich \sim '94)

If there exists a sequence $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability,

Ray approximation

Theorem (Kaimanovich \sim '94)

If there exists a sequence $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability, then (B, λ) is the Poisson boundary.

Ray approximation

Theorem (Kaimanovich \sim '94)

If there exists a sequence $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability, then (B, λ) is the Poisson boundary.

Corollary

If random walk tracks geodesics sublinearly

Ray approximation

Theorem (Kaimanovich \sim '94)

If there exists a sequence $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability, then (B, λ) is the Poisson boundary.

Corollary

*If **random walk tracks geodesics sublinearly** \rightarrow geometric boundary is Poisson boundary.*

Ray approximation

Theorem (Kaimanovich \sim '94)

If there exists a sequence $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability, then (B, λ) is the Poisson boundary.

Corollary

If random walk tracks geodesics sublinearly \rightarrow geometric boundary is Poisson boundary.


Pick $\pi_n(\omega) := \gamma_\omega(\ell n)$.

Ray approximation

If

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

then (B, λ) is the Poisson boundary.




Ray approximation

If

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

then (B, λ) is the Poisson boundary.




Ray approximation

If

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

then (B, λ) is the Poisson boundary.




Ray approximation

If

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

then (B, λ) is the Poisson boundary.




Ray approximation

If

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

then (B, λ) is the Poisson boundary.




Ray approximation

If

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

then (B, λ) is the Poisson boundary. Pick $\pi_n(\omega) := \gamma_\omega(\ell n)$.




Ray approximation

If

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

then (B, λ) is the Poisson boundary. Pick $\pi_n(\omega) := \gamma_\omega(\ell n)$.




Ray approximation


If

$$\frac{d(w_n, \gamma_\omega(\ell n))}{n} \rightarrow 0$$


then (B, λ) is the Poisson boundary. Pick $\pi_n(\omega) := \gamma_\omega(\ell n)$.




Sublinear tracking and entropy




Sublinear tracking and entropy



Sublinear tracking and entropy




Sublinear tracking and entropy



$$H_n(\mathbb{P}^\xi) \lesssim \log \#B(\gamma(\ell n), r_n)$$


Sublinear tracking and entropy



$$H_n(\mathbb{P}^\xi) \lesssim \log \#B(\gamma(\ell n), r_n)$$

since $r_n = o(n)$

Sublinear tracking and entropy



$$H_n(\mathbb{P}^\xi) \lesssim \log \#B(\gamma(\ell n), r_n)$$

since $r_n = o(n)$

$$h(\mathbb{P}^\xi) = \lim_{n \rightarrow \infty} \frac{H_n(\mathbb{P}^\xi)}{n} \lesssim \lim_{n \rightarrow \infty} \frac{1}{n} \log \#B(\gamma(\ell n), r_n) \rightarrow 0$$

Ray approximation

Theorem (Kaimanovich \sim '94)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary. If there are measurable functions $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability, then (B, λ) is the Poisson boundary.

Ray approximation

Theorem (Kaimanovich \sim '94)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary. If there are measurable functions $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability, then (B, λ) is the Poisson boundary.

Question. (Kaimanovich \sim '96) Does the same criterion work for **locally compact** groups?

Ray approximation

Theorem (Kaimanovich \sim '94)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary. If there are measurable functions $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability, then (B, λ) is the Poisson boundary.

Question. (Kaimanovich \sim '96) Does the same criterion work for **locally compact** groups?

(Kaimanovich-Woess '02) Ray criterion for invariant Markov operators (intermediate between discrete and lcsc groups)

Ray approximation for locally compact groups

Let G be a locally compact, second countable group,

Ray approximation for locally compact groups

Let G be a locally compact, second countable group, and μ a measure on G .

Ray approximation for locally compact groups

Let G be a locally compact, second countable group, and μ a measure on G . Let m be Haar measure, and $\rho := \frac{d\mu}{dm}$.

Ray approximation for locally compact groups

Let G be a locally compact, second countable group, and μ a measure on G . Let m be Haar measure, and $\rho := \frac{d\mu}{dm}$.

A metric d on G is temperate if $\exists C$ s.t.

$$m(\{g \in G : d(o, go) \leq R\}) \leq Ce^{CR}$$

Ray approximation for locally compact groups

Let G be a locally compact, second countable group, and μ a measure on G . Let m be Haar measure, and $\rho := \frac{d\mu}{dm}$.

A metric d on G is temperate if $\exists C$ s.t.

$$m(\{g \in G : d(o, go) \leq R\}) \leq Ce^{CR}$$

Theorem (Forghani-T. '19)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d ,

Ray approximation for locally compact groups

Let G be a locally compact, second countable group, and μ a measure on G . Let m be Haar measure, and $\rho := \frac{d\mu}{dm}$.

A metric d on G is temperate if $\exists C$ s.t.

$$m(\{g \in G : d(o, go) \leq R\}) \leq Ce^{CR}$$

Theorem (Forghani-T. '19)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary.

Ray approximation for locally compact groups

Let G be a locally compact, second countable group, and μ a measure on G . Let m be Haar measure, and $\rho := \frac{d\mu}{dm}$.

A metric d on G is temperate if $\exists C$ s.t.

$$m(\{g \in G : d(o, go) \leq R\}) \leq Ce^{CR}$$

Theorem (Forghani-T. '19)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary. If there are measurable functions $\pi_n : B \rightarrow G$ such that

Ray approximation for locally compact groups

Let G be a locally compact, second countable group, and μ a measure on G . Let m be Haar measure, and $\rho := \frac{d\mu}{dm}$.

A metric d on G is temperate if $\exists C$ s.t.

$$m(\{g \in G : d(o, go) \leq R\}) \leq Ce^{CR}$$

Theorem (Forghani-T. '19)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary. If there are measurable functions $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability,

Ray approximation for locally compact groups

Let G be a locally compact, second countable group, and μ a measure on G . Let m be Haar measure, and $\rho := \frac{d\mu}{dm}$.

A metric d on G is temperate if $\exists C$ s.t.

$$m(\{g \in G : d(o, go) \leq R\}) \leq Ce^{CR}$$

Theorem (Forghani-T. '19)

Let d be a temperate metric on G , suppose that μ has finite first moment w.r.t. d , and let (B, λ) be a μ -boundary. If there are measurable functions $\pi_n : B \rightarrow G$ such that

$$\frac{d(w_n, \pi_n(\text{bnd}(\omega)))}{n} \rightarrow 0$$

in probability, then (B, λ) is the Poisson boundary.

Applications - Isometries of hyperbolic spaces

Recall the drift

$$\ell := \lim_{n \rightarrow \infty} \frac{d(o, w_n o)}{n}$$

Applications - Isometries of hyperbolic spaces

Recall the drift

$$\ell := \lim_{n \rightarrow \infty} \frac{d(o, w_n o)}{n}$$

Theorem (Forghani-T. '19)

Let X be a proper δ -hyperbolic space

Applications - Isometries of hyperbolic spaces

Recall the drift

$$\ell := \lim_{n \rightarrow \infty} \frac{d(o, w_n o)}{n}$$

Theorem (Forghani-T. '19)

Let X be a proper δ -hyperbolic space and $G < \text{Isom}(X)$ be a closed subgroup with bounded exponential growth.

Applications - Isometries of hyperbolic spaces

Recall the drift

$$\ell := \lim_{n \rightarrow \infty} \frac{d(o, w_n o)}{n}$$

Theorem (Forghani-T. '19)

Let X be a proper δ -hyperbolic space and $G < \text{Isom}(X)$ be a closed subgroup with bounded exponential growth. Let μ be a (spread-out) measure on G with finite first moment.

Applications - Isometries of hyperbolic spaces

Recall the drift

$$\ell := \lim_{n \rightarrow \infty} \frac{d(o, w_n o)}{n}$$

Theorem (Forghani-T. '19)

Let X be a proper δ -hyperbolic space and $G < \text{Isom}(X)$ be a closed subgroup with bounded exponential growth. Let μ be a (spread-out) measure on G with finite first moment. Then:

- ▶ *if $\ell = 0$, then the Poisson boundary of (G, μ) is trivial;*

Applications - Isometries of hyperbolic spaces

Recall the drift

$$\ell := \lim_{n \rightarrow \infty} \frac{d(o, w_n o)}{n}$$

Theorem (Forghani-T. '19)

Let X be a proper δ -hyperbolic space and $G < \text{Isom}(X)$ be a closed subgroup with bounded exponential growth. Let μ be a (spread-out) measure on G with finite first moment. Then:

- ▶ *if $\ell = 0$, then the Poisson boundary of (G, μ) is trivial;*
- ▶ *if $\ell > 0$, then the Poisson boundary of (G, μ) is $(\partial X, \nu)$, where ∂X is the Gromov boundary.*

Applications - Isometries of hyperbolic spaces

Recall the drift

$$\ell := \lim_{n \rightarrow \infty} \frac{d(o, w_n o)}{n}$$

Theorem (Forghani-T. '19)

Let X be a proper δ -hyperbolic space and $G < \text{Isom}(X)$ be a closed subgroup with bounded exponential growth. Let μ be a (spread-out) measure on G with finite first moment. Then:

- ▶ *if $\ell = 0$, then the Poisson boundary of (G, μ) is trivial;*
- ▶ *if $\ell > 0$, then the Poisson boundary of (G, μ) is $(\partial X, \nu)$, where ∂X is the Gromov boundary.*
- ▶ For countable G : [Kaimanovich '94] .

Applications - Isometries of CAT(0) spaces

Theorem (Forghani-T. '19)

Let X be a proper CAT(0) space

Applications - Isometries of CAT(0) spaces

Theorem (Forghani-T. '19)

Let X be a proper CAT(0) space and $G < \text{Isom}(X)$ be a closed subgroup with bounded exponential growth.

Applications - Isometries of CAT(0) spaces

Theorem (Forghani-T. '19)

Let X be a proper CAT(0) space and $G < \text{Isom}(X)$ be a closed subgroup with bounded exponential growth. Let μ be a (spread-out) measure on G with finite first moment.

Applications - Isometries of CAT(0) spaces

Theorem (Forghani-T. '19)

Let X be a proper CAT(0) space and $G < \text{Isom}(X)$ be a closed subgroup with bounded exponential growth. Let μ be a (spread-out) measure on G with finite first moment. Then:

- ▶ *if $\ell = 0$, then the Poisson boundary of (G, μ) is trivial;*

Applications - Isometries of CAT(0) spaces

Theorem (Forghani-T. '19)

Let X be a proper CAT(0) space and $G < \text{Isom}(X)$ be a closed subgroup with bounded exponential growth. Let μ be a (spread-out) measure on G with finite first moment. Then:

- ▶ *if $\ell = 0$, then the Poisson boundary of (G, μ) is trivial;*
- ▶ *if $\ell > 0$, then the Poisson boundary of (G, μ) is $(\partial X, \nu)$, where ∂X is the visual boundary.*

Applications - Isometries of CAT(0) spaces

Theorem (Forghani-T. '19)

Let X be a proper CAT(0) space and $G < \text{Isom}(X)$ be a closed subgroup with bounded exponential growth. Let μ be a (spread-out) measure on G with finite first moment. Then:

- ▶ *if $\ell = 0$, then the Poisson boundary of (G, μ) is trivial;*
 - ▶ *if $\ell > 0$, then the Poisson boundary of (G, μ) is $(\partial X, \nu)$, where ∂X is the visual boundary.*
- ▶ For countable G : [Karlsson-Margulis '99].

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$


and edges $(z_1, z_2) \sim (y_1, y_2)$ if $z_1 \sim y_1$ and $z_2 \sim y_2$.

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$

and edges $(z_1, z_2) \sim (y_1, y_2)$ if $z_1 \sim y_1$ and $z_2 \sim y_2$.



Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$

and edges $(z_1, z_2) \sim (y_1, y_2)$ if $z_1 \sim y_1$ and $z_2 \sim y_2$.

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$

and edges $(z_1, z_2) \sim (y_1, y_2)$ if $z_1 \sim y_1$ and $z_2 \sim y_2$.

Theorem (Forghani-T. '19)

Let $G < \text{Aut}(DL_{q,r})$ be a closed subgroup,

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$

and edges $(z_1, z_2) \sim (y_1, y_2)$ if $z_1 \sim y_1$ and $z_2 \sim y_2$.

Theorem (Forghani-T. '19)

Let $G < \text{Aut}(DL_{q,r})$ be a closed subgroup, let μ be a measure on G

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$

and edges $(z_1, z_2) \sim (y_1, y_2)$ if $z_1 \sim y_1$ and $z_2 \sim y_2$.

Theorem (Forghani-T. '19)

Let $G < \text{Aut}(DL_{q,r})$ be a closed subgroup, let μ be a measure on G with finite first moment,

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$

and edges $(z_1, z_2) \sim (y_1, y_2)$ if $z_1 \sim y_1$ and $z_2 \sim y_2$.

Theorem (Forghani-T. '19)

Let $G < \text{Aut}(DL_{q,r})$ be a closed subgroup, let μ be a measure on G with finite first moment, and let V be its vertical drift.

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$

and edges $(z_1, z_2) \sim (y_1, y_2)$ if $z_1 \sim y_1$ and $z_2 \sim y_2$.

Theorem (Forghani-T. '19)

Let $G < \text{Aut}(DL_{q,r})$ be a closed subgroup, let μ be a measure on G with finite first moment, and let V be its vertical drift.

Then:

- 1. if $V = 0$, then the Poisson boundary of (G, μ) is trivial;*

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$

and edges $(z_1, z_2) \sim (y_1, y_2)$ if $z_1 \sim y_1$ and $z_2 \sim y_2$.

Theorem (Forghani-T. '19)

Let $G < \text{Aut}(DL_{q,r})$ be a closed subgroup, let μ be a measure on G with finite first moment, and let V be its vertical drift.

Then:

- 1. if $V = 0$, then the Poisson boundary of (G, μ) is trivial;*
- 2. if $V < 0$, then the Poisson boundary of (G, μ) is $\partial\mathbb{T}_q \times \{\omega_2\}$ with the hitting measure;*

Applications - Diestel-Leader graphs

Consider the Diestel-Leader graph

$$DL_{q,r} := \{(o_1, o_2) \in \mathbb{T}_q \times \mathbb{T}_r : h_1(o_1) + h_2(o_2) = 0\}.$$

and edges $(z_1, z_2) \sim (y_1, y_2)$ if $z_1 \sim y_1$ and $z_2 \sim y_2$.

Theorem (Forghani-T. '19)

Let $G < \text{Aut}(DL_{q,r})$ be a closed subgroup, let μ be a measure on G with finite first moment, and let V be its vertical drift.

Then:

- 1. if $V = 0$, then the Poisson boundary of (G, μ) is trivial;*
- 2. if $V < 0$, then the Poisson boundary of (G, μ) is $\partial\mathbb{T}_q \times \{\omega_2\}$ with the hitting measure;*
- 3. if $V > 0$, then the Poisson boundary of (G, μ) is $\{\omega_1\} \times \partial\mathbb{T}_r$ with the hitting measure.*

The Shannon-McMillan-Breiman theorem

The (Avez) entropy is

$$H(\mu) := - \sum_g \mu(g) \log \mu(g)$$

The Shannon-McMillan-Breiman theorem

The (Avez) entropy is

$$H(\mu) := - \sum_g \mu(g) \log \mu(g)$$

$$h := \lim_{n \rightarrow \infty} \frac{H(\mu_n)}{n}$$

The Shannon-McMillan-Breiman theorem

The (Avez) entropy is

$$H(\mu) := - \sum_g \mu(g) \log \mu(g)$$

$$h := \lim_{n \rightarrow \infty} \frac{H(\mu_n)}{n}$$

where $\mu_n(g) := \mathbb{P}(w_n = g)$.

The Shannon-McMillan-Breiman theorem

The (Avez) entropy is

$$H(\mu) := - \sum_g \mu(g) \log \mu(g)$$

$$h := \lim_{n \rightarrow \infty} \frac{H(\mu_n)}{n}$$

where $\mu_n(g) := \mathbb{P}(w_n = g)$.

Theorem (Kaimanovich-Vershik \sim '83, Derriennic \sim '85)

Let G be a countable group and μ a measure on G ,

The Shannon-McMillan-Breiman theorem

The (Avez) entropy is

$$H(\mu) := - \sum_g \mu(g) \log \mu(g)$$

$$h := \lim_{n \rightarrow \infty} \frac{H(\mu_n)}{n}$$

where $\mu_n(g) := \mathbb{P}(w_n = g)$.

Theorem (Kaimanovich-Vershik \sim '83, Derriennic \sim '85)

Let G be a countable group and μ a measure on G , with finite entropy h .

The Shannon-McMillan-Breiman theorem

The (Avez) entropy is

$$H(\mu) := - \sum_g \mu(g) \log \mu(g)$$

$$h := \lim_{n \rightarrow \infty} \frac{H(\mu_n)}{n}$$

where $\mu_n(g) := \mathbb{P}(w_n = g)$.

Theorem (Kaimanovich-Vershik \sim '83, Derriennic \sim '85)

Let G be a countable group and μ a measure on G , with finite entropy h . Then almost surely

$$\lim_{n \rightarrow \infty} \left(-\frac{1}{n} \log \mu_n(w_n) \right) = h.$$

The Shannon-McMillan-Breiman theorem

The (Avez) entropy is

$$H(\mu) := - \sum_g \mu(g) \log \mu(g)$$

$$h := \lim_{n \rightarrow \infty} \frac{H(\mu_n)}{n}$$

where $\mu_n(g) := \mathbb{P}(w_n = g)$.

Theorem (Kaimanovich-Vershik \sim '83, Derriennic \sim '85)

Let G be a countable group and μ a measure on G , with finite entropy h . Then almost surely

$$\lim_{n \rightarrow \infty} \left(-\frac{1}{n} \log \mu_n(w_n) \right) = h.$$

Proof. Using subadditivity:

$$\mu_{n+m}(w_{n+m}) \geq \mu_n(w_n) \mu_m(w_n^{-1} w_{n+m})$$

The Shannon-McMillan-Breiman theorem

Theorem (Kaimanovich-Vershik '83, Derriennic '85)

Let G be a countable group and μ a measure on G , with finite entropy h . Then almost surely

$$\lim_{n \rightarrow \infty} \left(-\frac{1}{n} \log \mu_n(w_n) \right) = h.$$

The Shannon-McMillan-Breiman theorem

Theorem (Kaimanovich-Vershik '83, Derriennic '85)

Let G be a countable group and μ a measure on G , with finite entropy h . Then almost surely

$$\lim_{n \rightarrow \infty} \left(-\frac{1}{n} \log \mu_n(w_n) \right) = h.$$

Proof. Using subadditivity:

$$\mu_{n+m}(w_{n+m}) \geq \mu_n(w_n) \mu_m(w_n^{-1} w_{n+m})$$

The Shannon-McMillan-Breiman theorem

Theorem (Kaimanovich-Vershik '83, Derriennic '81)

Let G be a countable group and μ a measure on G , with finite entropy h . Then almost surely

$$\lim_{n \rightarrow \infty} \left(-\frac{1}{n} \log \mu_n(w_n) \right) = h.$$

Proof. Using subadditivity:

$$-\log \mu_{n+m}(w_{n+m}) \leq -\log \mu_n(w_n) - \log \mu_m(w_n^{-1} w_{n+m})$$

The Shannon-McMillan-Breiman theorem

Theorem (Kaimanovich-Vershik '83, Derriennic '81)

Let G be a countable group and μ a measure on G , with finite entropy h . Then almost surely

$$\lim_{n \rightarrow \infty} \left(-\frac{1}{n} \log \mu_n(w_n) \right) = h.$$

Proof. Using subadditivity:

$$-\log \mu_{n+m}(w_{n+m}) \leq -\log \mu_n(w_n) - \log \mu_m(w_n^{-1} w_{n+m})$$

and by Kingman's theorem.

The Shannon-McMillan-Breiman theorem

Theorem (Kaimanovich-Vershik '83, Derriennic '81)

Let G be a countable group and μ a measure on G , with finite entropy h . Then almost surely

$$\lim_{n \rightarrow \infty} \left(-\frac{1}{n} \log \mu_n(w_n) \right) = h.$$

Proof. Using subadditivity:

$$-\log \mu_{n+m}(w_{n+m}) \leq -\log \mu_n(w_n) - \log \mu_m(w_n^{-1} w_{n+m})$$

and by Kingman's theorem.

Question. (Derriennic) Can we generalize to locally compact groups?

Entropies

Let μ be a measure on G , abs. cont. w.r.t. the Haar measure m

Entropies

Let μ be a measure on G , abs. cont. w.r.t. the Haar measure m and let $\rho := \frac{d\mu}{dm}$.

Entropies

Let μ be a measure on G , abs. cont. w.r.t. the Haar measure m and let $\rho := \frac{d\mu}{dm}$.

The differential entropy is

$$h_{diff}(\mu) := - \int_G \rho(g) \log \rho(g) dm(g)$$

Entropies

Let μ be a measure on G , abs. cont. w.r.t. the Haar measure m and let $\rho := \frac{d\mu}{dm}$.

The differential entropy is

$$h_{diff}(\mu) := - \int_G \rho(g) \log \rho(g) dm(g)$$

Note: it can be negative!

Entropies

Let μ be a measure on G , abs. cont. w.r.t. the Haar measure m and let $\rho := \frac{d\mu}{dm}$.

The **differential entropy** is

$$h_{\text{diff}}(\mu) := - \int_G \rho(g) \log \rho(g) dm(g)$$

Note: it can be negative!

The **Furstenberg entropy** of the μ -boundary (B, λ) is

$$h(\lambda) := \int_{G \times B} \log \frac{dg\lambda}{d\lambda}(\gamma) dg\lambda(\gamma) d\mu(g)$$

Poisson boundary

Let $(\partial G, \nu)$ be the Poisson boundary of (G, μ) .

Poisson boundary

Let $(\partial G, \nu)$ be the Poisson boundary of (G, μ) .

Theorem (Derriennic)

For any μ -boundary (B, λ) , we have

Poisson boundary

Let $(\partial G, \nu)$ be the Poisson boundary of (G, μ) .

Theorem (Derriennic)

For any μ -boundary (B, λ) , we have

$$h(\lambda) \leq h(\nu).$$

Poisson boundary

Let $(\partial G, \nu)$ be the Poisson boundary of (G, μ) .

Theorem (Derriennic)

For any μ -boundary (B, λ) , we have

$$h(\lambda) \leq h(\nu).$$

If $h(\nu)$ is finite,

Poisson boundary

Let $(\partial G, \nu)$ be the Poisson boundary of (G, μ) .

Theorem (Derriennic)

For any μ -boundary (B, λ) , we have

$$h(\lambda) \leq h(\nu).$$

If $h(\nu)$ is finite, then

$$h(\nu) = h(\lambda)$$

Poisson boundary

Let $(\partial G, \nu)$ be the Poisson boundary of (G, μ) .

Theorem (Derriennic)

For any μ -boundary (B, λ) , we have

$$h(\lambda) \leq h(\nu).$$

If $h(\nu)$ is finite, then

$$h(\nu) = h(\lambda) \Leftrightarrow (B, \lambda) \text{ is the Poisson boundary}$$

The Shannon-McMillan-Breiman theorem - locally compact case

Let G be a lcsc group,

The Shannon-McMillan-Breiman theorem - locally compact case

Let G be a lcsc group, μ a prob. measure on G ,

The Shannon-McMillan-Breiman theorem - locally compact case

Let G be a lcsc group, μ a prob. measure on G , $\mu \ll m$,

The Shannon-McMillan-Breiman theorem - locally compact case

Let G be a lcsc group, μ a prob. measure on G , $\mu \ll m$,
 $\rho := \frac{d\mu}{dm}$.

The Shannon-McMillan-Breiman theorem - locally compact case

Let G be a lcsc group, μ a prob. measure on G , $\mu \ll m$,
 $\rho := \frac{d\mu}{dm}$.

Theorem (Forghani-T. '19)

Let $(\partial G, \nu)$ be the Poisson boundary of (G, μ) .

The Shannon-McMillan-Breiman theorem - locally compact case

Let G be a lcsc group, μ a prob. measure on G , $\mu \ll m$,
 $\rho := \frac{d\mu}{dm}$.

Theorem (Forghani-T. '19)

Let $(\partial G, \nu)$ be the Poisson boundary of (G, μ) . If ρ is bounded and the differential entropy $H_n < \infty$ for any n ,

The Shannon-McMillan-Breiman theorem - locally compact case

Let G be a lcsc group, μ a prob. measure on G , $\mu \ll m$,
 $\rho := \frac{d\mu}{dm}$.

Theorem (Forghani-T. '19)

Let $(\partial G, \nu)$ be the Poisson boundary of (G, μ) . If ρ is bounded and the differential entropy $H_n < \infty$ for any n , then

$$\liminf_{n \rightarrow \infty} \left(-\frac{1}{n} \log \frac{d\mu_n}{dm}(w_n) \right) = h(\nu)$$

almost surely.

The Shannon-McMillan-Breiman theorem - locally compact case

Let G be a lcsc group, μ a prob. measure on G , $\mu \ll m$,
 $\rho := \frac{d\mu}{dm}$.

Theorem (Forghani-T. '19)

Let $(\partial G, \nu)$ be the Poisson boundary of (G, μ) . If ρ is bounded and the differential entropy $H_n < \infty$ for any n , then

$$\liminf_{n \rightarrow \infty} \left(-\frac{1}{n} \log \frac{d\mu_n}{dm}(w_n) \right) = h(\nu)$$

almost surely.

Recall $h(\nu)$ = Furstenberg entropy.

Proof of Shannon theorem

Let (B, λ) be a μ -boundary.

Proof of Shannon theorem

Let (B, λ) be a μ -boundary. Suppose $\rho \leq C$.

Proof of Shannon theorem

Let (B, λ) be a μ -boundary. Suppose $\rho \leq C$.

Let $\rho_n := \frac{d\mu_n}{d\mu}$,

Proof of Shannon theorem

Let (B, λ) be a μ -boundary. Suppose $\rho \leq C$.

Let $\rho_n := \frac{d\mu_n}{dm}$, $w_\infty := \text{bnd}(\omega)$.

Proof of Shannon theorem

Let (B, λ) be a μ -boundary. Suppose $\rho \leq C$.

Let $\rho_n := \frac{d\mu_n}{dm}$, $w_\infty := \text{bnd}(\omega)$.

1. By the ergodic theorem,

Proof of Shannon theorem

Let (B, λ) be a μ -boundary. Suppose $\rho \leq C$.

Let $\rho_n := \frac{d\mu_n}{d\mu}$, $w_\infty := \text{bnd}(\omega)$.

1. By the ergodic theorem, for \mathbb{P} -almost every $\omega = (w_n)$

$$\lim_n \frac{1}{n} \log \frac{dw_n \lambda}{d\lambda}(w_\infty) = h(\lambda).$$

Proof of Shannon theorem

Let (B, λ) be a μ -boundary. Suppose $\rho \leq C$.

Let $\rho_n := \frac{d\mu_n}{dm}$, $w_\infty := \text{bnd}(\omega)$.

1. By the ergodic theorem, for \mathbb{P} -almost every $\omega = (w_n)$

$$\lim_n \frac{1}{n} \log \frac{dw_n \lambda}{d\lambda}(w_\infty) = h(\lambda).$$

2. Let $f_n : (M, \kappa) \rightarrow \mathbb{R}^{\geq 0}$ with $\int f_n d\kappa \leq C$.

Proof of Shannon theorem

Let (B, λ) be a μ -boundary. Suppose $\rho \leq C$.

Let $\rho_n := \frac{d\mu_n}{dm}$, $w_\infty := \text{bnd}(\omega)$.

1. By the ergodic theorem, for \mathbb{P} -almost every $\omega = (w_n)$

$$\lim_n \frac{1}{n} \log \frac{dw_n \lambda}{d\lambda}(w_\infty) = h(\lambda).$$

2. Let $f_n : (M, \kappa) \rightarrow \mathbb{R}^{\geq 0}$ with $\int f_n d\kappa \leq C$. Then

$$\liminf_n \left(-\frac{1}{n} \log f_n(x) \right) \geq 0 \quad \kappa\text{-a.e.}$$

Proof of Shannon theorem

Let (B, λ) be a μ -boundary. Suppose $\rho \leq C$.

Let $\rho_n := \frac{d\mu_n}{d\mu}$, $w_\infty := \text{bnd}(\omega)$.

1. By the ergodic theorem, for \mathbb{P} -almost every $\omega = (w_n)$

$$\lim_n \frac{1}{n} \log \frac{dw_n \lambda}{d\lambda}(w_\infty) = h(\lambda).$$

2. Let $f_n : (M, \kappa) \rightarrow \mathbb{R}^{\geq 0}$ with $\int f_n d\kappa \leq C$. Then

$$\liminf_n \left(-\frac{1}{n} \log f_n(x) \right) \geq 0 \quad \kappa\text{-a.e.}$$

3. Check

$$\int_{\Omega} \frac{dw_{n-1} \lambda}{d\lambda}(w_\infty) \rho_n(w_n) d\mathbb{P} \leq C.$$

Proof of Shannon theorem - II

4. Hence

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) - \frac{1}{n} \log \frac{dw_n^\lambda}{d\lambda}(w_\infty) \right) \geq 0.$$

Proof of Shannon theorem - II

4. Hence

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) - \frac{1}{n} \log \frac{dw_n^\lambda}{d\lambda}(w_\infty) \right) \geq 0.$$

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) \geq h(\lambda)$$

Proof of Shannon theorem - II

4. Hence

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) - \frac{1}{n} \log \frac{dw_n^\lambda}{d\lambda}(w_\infty) \right) \geq 0.$$

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) \geq h(\lambda)$$

5. (Derriennic)

$$\lim_{n \rightarrow \infty} \int_{\Omega} -\frac{1}{n} \log \rho_n(w_n) d\mathbb{P} = h(\nu)$$

Proof of Shannon theorem - II

4. Hence

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) - \frac{1}{n} \log \frac{dw_n^\lambda}{d\lambda}(w_\infty) \right) \geq 0.$$

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) \geq h(\lambda)$$

5. (Derriennic)

$$\lim_{n \rightarrow \infty} \int_{\Omega} -\frac{1}{n} \log \rho_n(w_n) d\mathbb{P} = h(\nu)$$

6. By Fatou's lemma

$$h(\lambda) \leq \int_{\Omega} \liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) d\mathbb{P} \leq \lim_n \int_{\Omega} \left(-\frac{1}{n} \log \rho_n(w_n) \right) d\mathbb{P} = h(\nu)$$

Proof of Shannon theorem - II

4. Hence

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) - \frac{1}{n} \log \frac{dw_n^\lambda}{d\lambda}(w_\infty) \right) \geq 0.$$

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) \geq h(\lambda)$$

5. (Derriennic)

$$\lim_{n \rightarrow \infty} \int_{\Omega} -\frac{1}{n} \log \rho_n(w_n) d\mathbb{P} = h(\nu)$$

6. By Fatou's lemma

$$h(\lambda) \leq \int_{\Omega} \liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) d\mathbb{P} \leq \lim_n \int_{\Omega} \left(-\frac{1}{n} \log \rho_n(w_n) \right) d\mathbb{P} = h(\nu)$$

7. Thus setting $\lambda = \nu$,

Proof of Shannon theorem - II

4. Hence

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) - \frac{1}{n} \log \frac{dw_n^\lambda}{d\lambda}(w_\infty) \right) \geq 0.$$

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) \geq h(\lambda)$$

5. (Derriennic)

$$\lim_{n \rightarrow \infty} \int_{\Omega} -\frac{1}{n} \log \rho_n(w_n) d\mathbb{P} = h(\nu)$$

6. By Fatou's lemma

$$h(\lambda) \leq \int_{\Omega} \liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) d\mathbb{P} \leq \lim_n \int_{\Omega} \left(-\frac{1}{n} \log \rho_n(w_n) \right) d\mathbb{P} = h(\nu)$$

7. Thus setting $\lambda = \nu$, a.s.

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) = h(\nu)$$

Proof of Shannon theorem - II

4. Hence

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) - \frac{1}{n} \log \frac{dw_n^\lambda}{d\lambda}(w_\infty) \right) \geq 0.$$

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) \geq h(\lambda)$$

5. (Derriennic)

$$\lim_{n \rightarrow \infty} \int_{\Omega} -\frac{1}{n} \log \rho_n(w_n) d\mathbb{P} = h(\nu)$$

6. By Fatou's lemma

$$h(\lambda) \leq \int_{\Omega} \liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) d\mathbb{P} \leq \lim_n \int_{\Omega} \left(-\frac{1}{n} \log \rho_n(w_n) \right) d\mathbb{P} = h(\nu)$$

7. Thus setting $\lambda = \nu$, a.s.

$$\liminf_n \left(-\frac{1}{n} \log \rho_n(w_n) \right) = h(\nu)$$

QED

The end



Thank you!!!